

10

by Dr. Triawanti

Submission date: 27-Sep-2019 10:45AM (UTC+0700)

Submission ID: 1181037971

File name: 10._reduction_of_oxidative_stress.pdf (1.39M)

Word count: 4185

Character count: 22226

6
Reduction of Oxidative Stress by Seluang Fish
(*Rasbora spp.*) in Brain of Malnourished Rats
(*Rattus norvegicus*)

3
Triawanti and Didik Dwi Sanyoto
Biochemistry Department, Faculty of Medicine, the Medical Chemistry, Lambung Mangkurat University, Banjarmasin,
Indonesia
3
Email: tria_fkunlam@yahoo.co.id, didikdwisanyoto@yahoo.com

33
Hendra Wana Nur'amin
Faculty of Medicine, Lambung Mangkurat University, the Pharmacology Department, Banjarmasin, Indonesia
Email: hendranuramin@gmail.com

Abstract—Malnutrition children have lower intelligence due to brain cells damage caused by oxidative stress. Seluang (*Rasbora spp.*) has the potential to reduce this problem. This study aimed to analyze the correlation between hydrogen peroxide (H_2O_2) levels with malondialdehyde (MDA) levels in the brain of malnourished rats after administration of seluang. This study used a posttest only with control group design. 30 white rats (*Rattus norvegicus*) made into malnutrition and divided into three groups, M: malnutrition group, P1: malnourished rats fed standard diet, P2: malnourished rats fed seluang for 4 weeks. The results showed that malnourished rats had the highest levels of H_2O_2 . ANOVA statistical showed a significant differences between H_2O_2 levels ($p = 0.001$) and MDA levels ($p = 0.001$) among malnutrition group with the other groups Pearson correlation test results concluded that there was a correlation H_2O_2 levels with MDA levels ($p = 0.001$, $r = 0.684$) as indicated by decrease of H_2O_2 levels followed by decrease in MDA levels. This study showed a correlation between H_2O_2 levels and MDA levels in the of malnourished rat brain fed seluang. Thus seluang can improve the condition of oxidative stress in malnourished rat brain.

Index Terms— H_2O_2 , malnutrition, MDA, oxidative stress, seluang

I. INTRODUCTION

Malnutrition is still a serious problem in the health sector in both of developed and developing countries. In Indonesia, the prevalence of malnutrition remains high. Basic Health Research [Risikesdas] in 2013 reported that the prevalence of malnutrition in children under five at 5.7%, up from 5.4% in 2010. The prevalence of stunting increased to be 13.9% from 13.0% in 2010. In addition, of the 33 provinces in Indonesia, 20 provinces have prevalence of malnutrition above the national average. South Kalimantan was ranked 5th out of 18 provinces have toddlers weighing less than the national average [1].

Children who have less or poor nutritional status and short or very short at risk of losing intelligence quotient [IQ] by 10-15 points [2]. Nutrients, especially protein, minerals, vitamins and essential fatty acids are needed in the development of brain cells. Malnutrition due to protein deficiency would interfere with the synthesis of enzymes that act as antioxidants and reducing the concentration of antioxidants in the tissue, causing a state of deficiency of antioxidants and oxidative stress in the brain [3]. Excessive levels of oxidative stress [free radicals] will damage many cellular components, including cellular proteins, DNA, phospholipid membranes, and enzymes [4], [5]. Cellular component disruption have an impact on overall brain damage and intelligence [6].

Endogenous enzyme that works as antioxidant such as superoxide dismutase (SOD) which catalyze superoxide radicals to be H_2O_2 . Furthermore, H_2O_2 will be catalyzed by peroxidase (GPx) and catalase into water (H_2O) and oxygen (O_2). Thus, H_2O_2 can be used as one of the parameters of oxidative stress [5]. In addition, one of the most stable markers of oxidative stress is malondialdehyde (MDA) [7], [8]. MDA levels may indicate oxidative reactions, thus MDA levels rise in various diseases accompanied by excessive oxygen free radicals, so the relationship with the MDA levels of free radical damage is quite often researched [9], [10]. In a previous study, new malnutrition protein leads to oxidative stress in the brain of rats [11], [12]. There was an increase in lipid peroxidation accompanied by increased of MDA levels in the group of malnourished rats when compared to control rats [11].

Free radicals or other reactive oxygen species increase should be counterbalanced by raising synthesis of enzymatic antioxidants. However, this may not be sufficiently resolved by endogenous antioxidants only, thus exogenous antioxidants are needed. Endogenous antioxidant defenses in the brain are not adequate, can be reduced by administering exogenous antioxidants, so the balance of reactive oxygen species and antioxidant will be corrected [3]. These exogenous antioxidants can be obtained from natural components in food stuffs. One of

Manuscript received March 22, 2017; revised August 2, 2017.

them can be obtained from fish. Fish is a good source of nutrients; rich in protein and essential fatty acids [13]. Saluang [*Rasbora spp.*] is a river fish, known and widely consumed by the people of South Kalimantan. Based on research conducted by Yunanto *et al* [14], saluang contains amino acids cysteine (0.32 mg) and Docosahexaenoic acid (DHA) (1.04 grams) per 100 grams. Cysteine and DHA has a role in lowering the binding of various types of free radicals. DHA has an ability to inhibit the oxidative stress and protect the brain from lipids and proteins peroxidative damage in the brain [15], [16]. This is an opportunity to do research on potency of Seluang in the response to nutritional problems in Indonesia.

II. MATERIAL AND METHODS

A. Material

30 white male rats weighing 80-100 grams with an age range of 4 weeks, rice crust, seluang, standard rat feed (protein 30%, fat 5%, 8% fiber, calcium 9%, phosphorus 1%, and 3.8% amino acids), ether, phosphate buffer pH 7.4, 200 mL TCA 100%, 100 mL Na-Tiobarbiturat 1%, 250 mL HCl 1 N, H₂O₂, dichromate, acetic acid glacial, adrenalin, Na₂CO₃, EDTA, and distilled water.

B. Methods

This study used male white rats (*Rattus norvegicus*) as the models. This work was approved by the Animal Care and Experimentation Committee (Ethical Committee), Faculty of Medicine Lambung Mangkurat University).

1) Making seluang fish meal

Fresh seluang fish was milled and obtained raw fish porridge. Then steamed with hot steam for 1 hour. Then steaming results were dried in an oven until the moisture content reached 8%. Next step, grinding back to destroy clots or bone particles. After becoming dry grain, seluang fish meal was made into pellets and used as rat feed.

2) Giving treatment

All rats fed a low protein and fat for 8 weeks. Low-protein feed such as nasi karak (dry spoiled rice) was used as the equivalent of a low-protein feed 4% [13]. After 8 weeks (after rats are malnourished), 10 rats from malnutrition group were sacrificed and brains were taken and then measured the levels of H₂O₂ and MDA in the rat of brain. The P1 group fed with standard diet and P2 group fed seluang formula for 4 weeks. After that, all groups were terminated for the measure H₂O₂ and MDA levels in the rat brain.

3) Making brain homogenization

Brains were crushed with a mortar at room temperature and added 1 mL of phosphate buffer at pH 7.4 until it turned into a liquid. Then take 5 mL and centrifuged at a speed of 8000 rpm for 20 minutes. Furthermore, the supernatant was taken to measure levels of SOD and H₂O₂.

4) Measurement of rat brain MDA

Brains had just taken were crushed with a mortar at room temperature and added 1 mL of phosphate buffer at pH 7.4 until it turned into a liquid. Take 5 mL and centrifuged at a speed of 8000 rpm for 20 minutes. Furthermore, 200 mL of the supernatant was taken to measure the levels of MDA. Making the MDA standard

curve. A total of 0.05 μM MDA standard was added 1 mL aquadest, then accommodated in an Eppendorf tube. After that, successively added 100 mL TCA 100%, 100 mL Na-Tiobarbiturat 28%, and 250 mL of HCl 1 N. Then heated at 100°C for 20 minutes, then centrifuged at 3500 rpm for 10 minutes. After that, 450 mL of the supernatant was taken and added to distilled water up to 3500 μL. Then read with a spectrophotometer with a wavelength of at most 540 nm. The same procedures were done to 0.025, 0.0125, 0.00625, 0.003125 and 1.56 x 10⁻⁵ μM MDA. Then graphed the relationship between the absorbance on the Y axis with MDA on the X axis to obtain a linear equation.

5) Examination of hydrogen peroxide

Examination of the samples to measure the levels of H₂O₂ using a spectrophotometer.

6) Making the standard curve

A total of 20 mol H₂O₂ was added 2 ml of a mixture of dichromate: glacial acetic acid (1:3). Then the mixture was heated in boiling water for 10 minutes. The mixture was cooled and then measured the absorbance at 570 nm wavelength. The same procedures were done to 40,60,80,100,120, 140,160 and 180 mol H₂O₂. Then graphed the relationship between the absorbance on the Y axis with the levels of H₂O₂ on the X axis to obtain a linear equation.

7) Making the test solution

1 ml of brain homogenates was added 5 ml phosphate buffer at pH 7.4. A total of 1 ml of the mixture was taken and added to a mixture of 2 ml of dichromate: acetate (1: 3) and then wrapped in aluminum foil for 30 minutes. The mixed solution is heated using a water bath for 10 minutes at 100°C. The solution was cooled at room temperature. The solution was then transferred into a cuvette and its absorbance was measured using a UV-VIS at a wavelength of 570 nm.

C. Data Analysis

The data obtained were tested using normality and homogeneity test. Normal and homogeneously distributed data were tested by one way Anova test followed by post hoc test (Tukey HSD) and Pearson correlation test at 95% confidence level.

III. RESULTS

This study used malnourished white rats by providing low protein fat for 8 weeks. After 8 weeks, 10 rats were sacrificed to measure serum protein levels and brain were taken to measure the levels of H₂O₂ and MDA. Furthermore, on the other malnourished rats were divided into two groups and fed in accordance with their respective groups for 4 weeks. After 4 weeks, rats were terminated and brains were taken. The results of H₂O₂ and MDA measurements were presented respectively in Fig. 1 and Fig. 2.

Fig. 1 showed that group of malnourished rats without other feeding had the highest H₂O₂ levels. While groups of malnourished rats fed seluang had the lowest levels of H₂O₂. The statistical test result with one-way ANOVA on the data of H₂O₂ showed p = 0.001 (p <0.05), which

showed a significant difference at a minimum of one group. These results indicated that the condition of malnutrition increased oxidative stress which marked the levels of H₂O₂ was high. Then after rats were fed standard diet (P1) and seluang (P2), levels of H₂O₂ tended to be lower. This proved that the feed able to reduce oxidative stress in the brain of malnourished rats.




Figure 1. Average levels of H₂O₂ in the malnourished rat brain (M), standard diet (P1), and Seluang (P2) (p = 0.001).




Figure 2. Average levels of MDA in malnourished rat brain (M), after being given a standard diet (P1), and Seluang (P2) [p = 0.001]

Fig. 2 showed that malnutrition group without any other diet had the highest MDA levels. After given the standard diet (P1) and Seluang (P2) there were MDA levels reduction. One way Anova test results concluded that there were differences between groups [p = 0.001]. These results proved that seluang could reduce MDA levels of malnourished rat brains.

Pearson correlation test with the 95% confidence level was done to determine a correlation between H₂O₂ levels and MDA levels in malnourished rat brain fed seluang. Fig. 3 showed the correlation between H₂O₂ levels and MDA levels of malnourished rat brain.




Figure 3. Correlation between H₂O₂ levels and MDA levels of malnourished rat brain after feeding seluang

Based on correlation test, there was significant relationship [p = 0.001] with moderate positive correlation [r = 0.684]. It meant increased H₂O₂ levels followed by elevated MDA levels and vice versa. This proved that seluang able to reduce H₂O₂ levels of malnourished rat brain followed by decreased MDA levels.

IV. DISCUSSION

Malnutrition is a condition caused by an imbalance between the supply of energy and protein to fulfill body needs to function and grow optimally. Children who are malnourished have more oxidant damage products, antioxidant levels are lower, and decreased antioxidant defense system [3]. This is due to a deficiency of the protein in the diet interfere with the synthesis of antioxidant enzymes and reduce the concentration of antioxidants in the tissue, causing a state of antioxidants deficiency [3].

Brain is well protected from free radicals damage by the balance between prooxidant and antioxidant mechanisms, including antioxidant enzymes and free chemical radical scavenger [5], [9]. This balance tends to be impaired in protein energy malnutrition (PEM). Many clinical and pathologic manifestations in the PEM alleged to be the result of an imbalance between the defense of free radicals and free radical production [17], [18]. Pathological hallmark of protein deficiency include shrinkage of brain size and an increase in lipid peroxidation in the brain [19].

Oxidative stress may occur when there is an imbalance between the reactive oxygen species [ROS], consisting of superoxide, hydrogen peroxide, and hydroxyl radicals, with defensive oxidative inadequate, including superoxide dismutase [SOD], catalase [CAT], and peroxidase [POX] [5], [11]. This condition is very harmful for cells. ROS is the result of normal aerobic metabolism that can not be avoided. However, ROS production can be increased also by the amount of stress that disturb cellular homeostasis [5]. ROS will react with lipids, proteins, pigments, and nucleic acids, causing lipid peroxidation [20], [21].

This study demonstrated that H₂O₂ levels of in the malnutrition group were very high. This indicated that under malnutrition conditions would increase the formation of H₂O₂. H₂O₂ is derived from superoxide radicals and converted by the SOD enzyme into H₂O₂. Furthermore, the H₂O₂ levels should be counterbalanced by an increase in the activity of GPx enzyme and catalase which converts H₂O₂ to H₂O and O₂. When this enzyme or its activity slightly lower than H₂O₂, it would be decomposed spontaneously into hydroxyl radicals (OH•) that more dangerous than H₂O₂ [5], [6].

This research showed that in the malnutrition group, levels of malondialdehyde (MDA) was higher. One of the markers of oxidative stress is the MDA. Malondialdehyde is a three-carbon component formed from peroxidation of polyunsaturated fatty acids, particularly arachidonic acid and is one of the most stable end product of lipid peroxidation. Oxidative stress can damage the central nervous system because it is very susceptible induced by a variety of biological agents. Due to the high metabolic

activity, brain requires molecular oxygen in large quantities, which is then followed by the formation of free radicals in high levels. The brain also contains polyunsaturated fatty acids and oxidized easily [22], [23]. In addition, the total capacity of antioxidants found in the central nervous system are relatively few [22], [24]. Increased oxidative stress may also be a result of the bad effects of calorie deficiency and intake of micronutrients [3].

Excessive levels of free radicals will damage many cellular components, including cellular proteins, DNA, phospholipids membrane and enzyme inactivation [4], [5]. It also causes a loss of function of specific proteins, clearance of abnormal proteins, cellular redox balance depletion and disruption of the cell cycle, and ultimately cause death of nerve [22], [23].

This study denoted the impact of protein malnutrition to levels of free radicals in experimental animals. Other study found malnourished children have more oxidant damage products, while lower antioxidant levels with decreased antioxidant defense system [3]. This was due to a deficiency of the protein in the diet interfere with the synthesis of antioxidant enzymes and reduce the concentration of antioxidants in the tissue, causing a state of antioxidants deficiency [3].

In this study, malnourished rats were given treatment in the form of standard diet [P1] and seluang [P2]. After the diet was given for 4 weeks, the rats were sacrificed and the measurement of the brain levels of H₂O₂ and MDA. The results showed all groups had decreased H₂O₂ levels and MDA levels significantly. The decline that occurred in the group fed seluang larger than the group fed standard. This proves that seluang was able to improve the condition of brain oxidative stress better due to malnutrition. The results of Pearson correlation test also proved that the decreased levels of H₂O₂ in the malnourished rat brain by seluang correlated with decreased MDA significantly. This proved that seluang able to improve the condition of the brain oxidative stress due to malnutrition better than regular feed.

Seluang had been shown to contain cysteine amino acids and DHA [14]. Both of these nutrients have some roles as antioxidant [16], [25]. Amino acids are known to have significant use as an antioxidant. The reaction between oxidized lipids with amino acids produce a lot of non-enzymatic reaction components, which have antioxidant effects [16], [25]. One of the 20 amino acids for basic human needs is cysteine. Cysteine is a natural amino acid that contains sulfur and unique because it contains thiol, and can be found in most of the protein, although only in very small amounts. The natural form of cysteine is L-cysteine. This amino acid is a precursor of glutathione, which its sulfhydryl group (*thiol*) acts as a proton donor and responsible for the biological activity of glutathione. Glutathione is an antioxidant tripeptide made of three amino acids, cysteine, glycine, and glutamate. Glutathione is an important antioxidant contained in vital organs and bone marrow of live animals, and often referred to as primary antioxidants [16]. Glutathione works on the peroxide substrate. Seluang contains cysteine

which act as antioxidants with levels of about 0.32 mg/100g [14]. Cysteine plays a role in binding various types of free radicals *in vitro*, one of them is OH•, whereas *in vivo*, cysteine together with glycine and glutamate is a glutathione precursor enzyme, an enzyme which plays a role in the ROS oxidation reduction [16].

In addition, seluang also contains DHA levels of 1.04 mg/100 g. Docosahexaenoic acid [DHA, 22; 6n-3] is a 22 unsaturated fatty acid chains [LCPs], which is an important substance for the development of the central nervous system of mammals [14]. DHA plays an important role in neurocognitive development [26] and suggested to have a relationship with malnutrition incident [27].

DHA has an important role in inhibiting oxidative stress and protect from damage lipid peroxidation and protein in the brains of children who are still developing and adult brain, thus avoiding the loss of neurons, cognitive deficits, and locomotor. In this study we demonstrated that DHA and amino acids in the seluang [*Rasbora spp.*] was better than standard diet against to oxidative stress on malnourished rat brain.

2 CONCLUSION

21
Based on the results, it can be concluded that there was a significant correlation between H₂O₂ levels and MDA levels in the brain of malnourished rats fed seluang. Seluang [*Rasbora spp.*] can improve oxidative stress profile as indicated by decrease of H₂O₂ levels and decrease in MDA levels. Therefore, seluang [*Rasbora spp.*] have potency to be a source of nutrient for malnutrition.

ACKNOWLEDGMENT

This study was funded by Research Grant from Faculty of Medicine, Lambung Mangkurat University.

REFERENCES

- [1] Kementerian Kesehatan RI. Profil kesehatan Indonesia 2014. Departemen Kesehatan RI, 2015.
- [2] Bappenas. Rencana Aksi Nasional Pangan dan Gizi 2011-2015. 2011, pp. 1-63.
- [3] M. Khare, C. Mohanty, B. K. Das, A. Jyoti, B. Mukhopadhyay, and S. P. Mishra, "Free radicals and antioxidant status in protein energy malnutrition," *Int. J. Pediatr.*, 2014, 254396.
- [4] Z. Radak, O. Marton, E. Nagy, E. Koltai, and S. Goto, "The complex role of physical exercise and reactive oxygen species on brain," *J. Sport Heal Sci.*, vol. 2, no. 2, pp. 87-93, 2013.
- [5] S. M. Al-Dalaen, "Review article: Oxidative stress versus antioxidants," *Am J. Biosci Bioeng.* vol. 2, no. 5, p. 60, 2014.
- [6] J. A. Hernández, R. C. López-Sánchez, and A. Rendón-Ramírez, "Lipids and oxidative stress associated with ethanol-induced neurological damage," *Oxid Med Cell Longev.* 2016.
- [7] Q. Yin, H. Ge, C. C. Liao, D. Liu, S. Zhang, and Y. H. Pan, "Antioxidant defenses in the brains of bats during hibernation," *PLoS One.*, vol. 11, no. 3, pp. 1-17, 2016.
- [8] R. Mancinelli, E. Barlocchi, S. Palminiello, and L. Saso, "Oxidative stress and brain diseases: Biomarkers and analytical methodologies," *Indian J. Biotechnol.*, vol. 10, no. 4, pp. 395-403, 2011.
- [9] M. Victoria Urquiza-Martínez and B. Fenton Navarro, "Antioxidant capacity of food," *Free Radicals Antioxidants*, vol. 6, no. 1, pp. 1-12, 2016.
- [10] L. F. Wei, H. M. Zhang, S. Wang, et al., "Changes of MDA and SOD in brain tissue after secondary brain injury with seawater

immersion in rats," *Turk Neurosurg*, vol. 26, no. 3, pp. 384–8, 2016.

[11] R. A. Ghone, A. N. Suryakar, *et al.*, "A study of oxidative stress biomarkers and effect of oral antioxidant supplementation in severe acute malnutrition," *J. Clin Diagnostic Res.*, vol. 7, no. 10, pp. 2146–2148, 2013.

[12] A. M. Feoli, I. R. Siqueira, L. Almeida, A. C. Tramontina, C. Vanzella, S. Sbaraini, *et al.*, "Effects of protein malnutrition on oxidative status in rat brain," *Nutrition*, vol. 22, no. 2, pp. 160–165, 2006.

[13] M. F. Laus, L. D. M. F. Vales, T. M. B. Costa, and S. Almeida, "Early postnatal protein-calorie malnutrition and cognition: A review of human and animal studies," *Int. J. Environ Res Public Health*, vol. 8, no. 2, pp. 590–612, 2011.

[14] A. Yunanto, D. D. Sanyoto, M. S. Noor, and I. Kustiyah, *Pengembangan potensi ikan saluang (Rasbora spp.) sebagai sumber nutrisi yang meningkatkan memori spasial dan ekspresi peroxisome proliferator activated receptor (ppar) otak: studi eksperimental pada tikus Rattus norvegicus*, 2014.

[15] X. Yu and Y. C. Long, "Crosstalk between cysteine and glutathione is critical for the regulation of amino acid signaling pathways and ferroptosis," *Sci Rep*, 2016.

[16] P. Piste, "Cysteine-master antioxidant," *Int. J. Pharm, Chem Biol. Sci.*, vol. 3, no. 1, pp. 143–149, 2013.

[17] D. Lee, W. Hwang, M. Artan, D. E. Jeong, and S. J. Lee, "Effects of nutritional components on aging," *Aging Cell.*, vol. 14, no. 1, pp. 8–16, 2015.

[18] G. S. Aly, A. H. Shaalan, M. K. Mattar, H. H. Ahmed, M. E. Zaki, and H. R. Abdallah, "Oxidative stress status in nutritionally stunted children," *Egypt Pediatr Assoc Gaz.*, vol. 62, no. 1, pp. 28–33, 2014.

[19] E. Vásquez-Garibay, K. Stein, C. Gómez, E. Romero, G. Hernández-flores, and A. B. Cuellar, "Effect of nutritional recovery on the serum concentration of lipid peroxide in children with primary and severe protein energy," *JONNPR*, vol. 1, no. 4, pp. 149–155, 2016.

[20] A. Catalá and M. Díaz, "Editorial: Impact of lipid peroxidation on the physiology and pathophysiology of cell membranes," *Front Physiol.*, vol. 7, pp. 1–3, 2016.

[21] R. Shirley, E. Ord, and L. Work, "Oxidative stress and the use of antioxidants in stroke," *Antioxidants*, vol. 3, no. 3, pp. 472–501, 2014.

[22] A. Rahal, A. Kumar, V. Singh, *et al.*, "Oxidative stress, prooxidants, and antioxidants: The interplay," *Biomed Res Int.*, 2014.

[23] D. A. Butterfield, M. Perluigi, T. Reed, *et al.*, "Redox proteomics in selected neurodegenerative disorders: From its infancy to future applications," *Antioxid Redox Signal.*, vol. 17, no. 11, pp. 1610–1655, 2012.

[24] P. S. Baxter and G. E. Hardingham, "Adaptive regulation of the brain's antioxidant defences by neurons and astrocytes," *Free Radic Biol Med.*, vol. 100, pp. 1–6, 2016.

[25] V. Casañas-Sánchez, J. A. Pérez, N. Fabelo, *et al.*, "Addition of docosahexaenoic acid, but not arachidonic acid, activates glutathione and thioredoxin antioxidant systems in murine hippocampal HT22 cells: Potential implications in neuroprotection," *J. Neurochem.*, vol. 131, no. 4, pp. 470–483, 2014.

[26] E. L. Prado and K. G. Dewey, "Nutrition and brain development in early life," *Nutr Rev.* vol. 72, no. 4, pp. 267–284, 2014.

[27] R. H. Glew and D. J. Vanderjagt, "Does docosahexaenoic acid play a role in infant malnutrition in the children of Fulani nomads in Northern Nigeria?" *Journal of Medicine in the Tropics*, vol. 15, no. 2, pp. 69–75, 2013.



Triawanti was born in Surabaya, Indonesia, in September 1971. She was graduated as physician in 1998 from Lambung Mangkurat University, Banjarmasin, Indonesia. She was graduated master degree 2004 from Airlangga University, Surabaya, Indonesia, PhD 2012 from Brawijaya University, Malang, Indonesia. She currently works in Medical Biochemistry Department, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia.



Didik Dwi Sanyoto was born in Pelaihari, Indonesia, in March 1972. He was graduated as physician in 1998 from Lambung Mangkurat University, Banjarmasin, Indonesia. He was graduated master degree 2003 from Airlangga University, Surabaya, Indonesia. He currently works in Medical Anatomy Department, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia.



Hendra Wana Nur'amin was born in Kandangan, Indonesia in February 1991. He was graduated as physician in 2013 from Lambung Mangkurat University, Banjarmasin, Indonesia. He got his master degree from Gadjah Mada University, Yogyakarta, Indonesia in 2016. He currently works in Medical Pharmacology Department, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia.

ORIGINALITY REPORT

13%

SIMILARITY INDEX

5%

INTERNET SOURCES

8%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	Telma Magaia, Amália Uamusse, Ingegerd Sjöholm, Kerstin Skog. "Proximate Analysis of Five Wild Fruits of Mozambique", The Scientific World Journal, 2013 Publication	1%
2	repo.unand.ac.id Internet Source	1%
3	Submitted to Lambung Mangkurat University Student Paper	1%
4	Submitted to University of Bath Student Paper	1%
5	www.freepatentsonline.com Internet Source	<1%
6	www.liebertpub.com Internet Source	<1%
7	Ricielle Lopes Augusto, Alinny Rosendo Isaac, Ivanildo Inácio da Silva-Júnior, David Filipe de Santana et al. "Fighting Oxidative Stress: Increased Resistance of Male Rat Cerebellum	<1%

at Weaning Induced by Low Omega 6/Omega 3 Ratio in a Protein-Deficient Diet", The Cerebellum, 2016

Publication

-
- | | | |
|----|---|-----|
| 8 | D. Calderon-Guzman. "Oxidative stress induced by morphine in brain of rats fed with a protein deficient diet", Human & Experimental Toxicology, 09/01/2009
Publication | <1% |
| 9 | m.blog.daum.net
Internet Source | <1% |
| 10 | baadalsg.inflibnet.ac.in
Internet Source | <1% |
| 11 | www.mdpi.com
Internet Source | <1% |
| 12 | www.sfrr-thai.net
Internet Source | <1% |
| 13 | media.neliti.com
Internet Source | <1% |
| 14 | www.tandfonline.com
Internet Source | <1% |
| 15 | Ayokunle O. Ademosun, Ganiyu Oboh, Fatai Bello, Peluola O. Ayeni. "Antioxidative Properties and Effect of Quercetin and Its Glycosylated Form (Rutin) on | <1% |

Acetylcholinesterase and Butyrylcholinesterase Activities", Journal of Evidence-Based Complementary & Alternative Medicine, 2016

Publication

16

nardus.mpn.gov.rs

Internet Source

<1%

17

Graciela Gavia-García, Haydeé González-Martínez, Ángel Miliar-García, Edmundo Bonilla-González et al. "Oxidative damage and antioxidant defense in thymus of malnourished lactating rats", Nutrition, 2015

Publication

<1%

18

onlinelibrary.wiley.com

Internet Source

<1%

19

Masoud Amiri. "Oxidative stress and free radicals in liver and kidney diseases; an updated short-review", Journal of Nephropathology, 2017

Publication

<1%

20

B R Cardoso, D J Hare, A I Bush, B R Roberts. "Glutathione peroxidase 4: a new player in neurodegeneration?", Molecular Psychiatry, 2016

Publication

<1%

21

Nesma A.M. Ghonimi, Mohamed E. Mahdy, Osama A. Abdel Salam. "Total Antioxidant Capacity Predicts Outcome in Acute Ischemic Stroke Subtypes in Egyptian Patients", Journal

<1%

22

Xintian Yan, Xinzhi Zhao, Juxue Li, Lin He, Mingqing Xu. "Effects of early-life malnutrition on neurodevelopment and neuropsychiatric disorders and the potential mechanisms", Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2018

Publication

<1%

23

Mayara Queiroga Barbosa, Rita de Cássia Ramos Egypto Queiroga, Camila Carolina de Menezes Santos Bertozzo, Daline Fernandes de Souza Araújo et al. "Effect of diets with goat milk fat supplemented with exercise on anxiety and oxidative stress in the brains of adult rats", Food & Function, 2018

Publication

<1%

24

Terez Boshra Kamel, Tharwat Ezzat Deraz, Rasha H. Elkabarity, Rasha K. Ahmed. "Protein energy malnutrition associates with different types of hearing impairments in toddlers: Anemia increases cochlear dysfunction", International Journal of Pediatric Otorhinolaryngology, 2016

Publication

<1%

25

Janina Galler, Danielle Galler Rabinowitz. "The Intergenerational Effects of Early Adversity",

<1%

26

Lina Lu, Zongli Li, Yanhua Zuo, Libo Zhao, Bin Liu. "Radioprotective activity of glutathione on cognitive ability in X-ray radiated tumor-bearing mice", Neurological Research, 2018

Publication

<1%

27

Di Domenico, Fabio, Raffaella Coccia, Annalisa Coccio, M. Paul Murphy, Giovanna Cenini, Elizabeth Head, D. Allan Butterfield, Alessandra Giorgi, Maria Eugenia Schinina, Cesare Mancuso, Chiara Cini, and Marzia Perluigi. "Impairment of proteostasis network in Down syndrome prior to the development of Alzheimer's disease neuropathology: Redox proteomics analysis of human brain", Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease, 2013.

Publication

<1%

28

www.karger.com

Internet Source

<1%

29

Submitted to University of Edinburgh

Student Paper

<1%

30

James Nathan Cobley, Maria Luisa Fiorello, Damian Miles Bailey. "13 reasons why the brain is susceptible to oxidative stress", Redox Biology, 2018

<1%

31

Patricia T. Alpert. "New and Emerging Theories of Cardiovascular Disease: Infection and Elevated Iron", Biological Research For Nursing, 2016

Publication

<1%

32

www.frontiersin.org

Internet Source

<1%

33

Maria Filkova. "MiRs in RA: possible biomarkers and therapeutic targets", Arthritis Research & Therapy, 2012

Publication

<1%

34

Jintanaporn Wattanathorn, Wipawee Thukhammee, Supaporn Muchimapura, Panakaporn Wannanon, Terdthai Tong-un, Somsak Tiamkao. "Preventive Effect of Cashew-Derived Protein Hydrolysate with High Fiber on Cerebral Ischemia", BioMed Research International, 2017

Publication

<1%

35

Olusegun L. Adebayo, Gbenga A. Adenuga, Rajat Sandhir. "Selenium and zinc protect brain mitochondrial antioxidants and electron transport chain enzymes following postnatal protein malnutrition", Life Sciences, 2016

Publication

<1%

36

Submitted to University of Southampton

Student Paper

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On