

11

by Dr. Triawanti

Submission date: 27-Sep-2019 10:47AM (UTC+0700)

Submission ID: 1181040051

File name: 11._seluang_and_PPAR.pdf (1.62M)

Word count: 3596

Character count: 19670

The Quality of Rat Brain Spatial Memory and Expression of Peroxisome Proliferator Activated Receptor (PPAR) Which Fed with Seluang (*Rasbora spp.*)

Ari Yunanto¹, Didik Dwi Sanyoto², Meitria Syahadatina Noor³, Ika Kustiyah⁴, and Triawanti⁵

¹Pediatric Department Faculty of Medicine Lambung Mangkurat University/Ulin General Hospital, Banjarmasin, Indonesia

²Medical Anatomy Department Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

³Public Health Department, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

⁴Pathology of Anatomy Department, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

⁵Medical Chemistry/Biochemistry Department, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

Email: {ariefkaunlam, ikaoktaviyanti}@gmail.com, {didikdwisanyoto, drmeitria}@yahoo.com; tria_fkunlam@yahoo.co.id

Abstract—Seluang fish (*Rasbora spp.*) besides contain high protein also contain an essential fatty acids (PUFA), which can increase intelligence. The objective of this research was to prove that Seluang fish is one of the nutrient source to increase the spatial memory and expression of PPAR brain tissue by using *Rattus norvegicus* as a model. Rats were divided into 3 groups: control group were fed standard (C), Treatment Group 1 were fed low-protein and fat (P1), Group Treatment 2 (P2) were fed standard with Seluang Formula, for 8 weeks. The first phase was to testing spatial memory using the Morris Water Maze. The second phase, was the making brain tissue's slide preparation, immunohistochemical stained, calculated the percentage of PPAR α and PPAR γ expression. The data analysed using ANOVA and Kruskal Wallis test with a 95% confidence level. The first phase results showed the percentage of period time in the target area for the P2 group is the highest compared to the C and P1 group, respectively 33.52%; 29.9%; and 23.15% ($p = 0.021$). On PPAR α and PPAR γ examination showed the average percentage of PPAR α neurons expression in group P2 is higher than the C and P1 group, respectively 18.5%, 3.75% and 2.5% ($p = 0.002$). Average percentage of total PPAR γ neuronan cells expression in P2 group also higher than the C and P1 group, respectively 35.41%, 6.25% and 5.83% ($p = 0.001$). The study shows that rats fed with Seluang Fish Formula able to improved the brain tissue's spatial memory and PPAR expression.

Index Terms—seluang fish (*Rasbora Spp*), spatial memory, PPAR expression.

I. INTRODUCTION

South Kalimantan included in 18 provinces that have low weight toddler, above the national rank. In 2010, prevalence of bad nutrition was 6%, low nutrition was

16.8%, stunting toddler was 35.3%. It still above the RPJMN target in 2014 (32%) [1]. It is a bad news, because South Kalimantan has a lot of natural resources, like freshwater fish. The freshwater fish consumption level of South Kalimantan community was 36.84 kg/period [2]. This amount is still higher than the National freshwater fish consumption level that around 33.89 kg/period, but it still lower than Malaysia (55.4 kg/period) and Singapore.

Seluang fish (*Rasbora spp.*) is a freshwater fish which famous and common consumed among South Kalimantan community. This is a potential opportunity to do research of South Kalimantan Seluang fish to response the nutritional problems in South Kalimantan in particular and Indonesia in general.

The role of nutrition in infancy greatly affect intelligence. Children who have less or poor nutritional status and short or very short children are at risk of losing intelligence or Intelligence Quotient (IQ) by 10-15 points [3]. Nutrients, especially protein, minerals, vitamins and essential fatty acids are needed in the brain cells development. Studies showed that by giving dietary omega-3 fatty acids in rats can increase PPAR gene expression which furthermore improves the learning and spatial memory. Omega-3 fatty acids increase the membrane fluidity by replacing the omega 6 and cholesterol from the cell membrane, maintaining membrane fluidity optimally is necessary to neurotransmitter binding and signaling in cells. Omega 3 incorporation in the nerve cells membranes also increased the synaptic proteins expression and strengthen plasticity synapses in hippocampus. This process is modulated by transcription factors which is peroxisome proliferator-activated receptors (PPARs) [4].

Omega 3 of fatty acid commonly found in fish as in Seluang fish from South Kalimantan. Furthermore

Manuscript received March 11, 2015; revised August 12, 2015.

research is need to recognize all its potentials as omega 3 fatty acid resource that can increase intelligence. The model in this study are rats (*Rattus norvegicus*). This study aims to determine the rats spatial memory quality that fed with Seluang fish compared to standard and low protein feed.

II. MATERIAL AND METHOD

A. Material

Material in this study were 30 male rats within age of 4 weeks old (*Rattus norvegicus*), rat's blood, aquades, deionized water, standard rats feed, rat feed with low fat and protein, Seluang fish formula, antibodi PPAR γ (rat), antibodi PPAR α (rat), kit imunostaining, formalin, eter, DAB chromogen, alcohol, formalin 10%, eter, Morris Water Maze (MWM) device.

B. Procedure

Rats divided into 3 groups, control (C) that were given by standard feed, treatment 1 (P1) that given low protein feed (dry rice that was made into pellet), it is equivalent to 4% low protein [5]. Treatment 2 (P2) that given Seluang fish extract powder feed, consist of 25% standard feed and 75% Seluang fish extract powder, made into pellet. The feed is given to rats from the age 4 weeks old until 12 weeks old, everyday ad libitum.

TABLE I. NUTRITIONS IN 100 GRAMS SELUANG FISH EXTRACT POWDER

Nutrition	Concentration (%)
Ca	1,6
Fe	19,9 ppm
P	1,67
Zn	122,7 ppm
Oleat acid	11,77
Linoleat acid	7,33
14-HA	1,04
Aspartat	3,71
Glutamat	4,98
Serin	1,94
Glisin	3,06
Histidin	1,82
Arginin	3,05
Threonin	2,15
Alanin	2,83
Prolin	1,92
Valin	2,41
Metionin	0,48
Isoleusin	2,07
Leusin	3,62
Phenilalanin	2,37
Lisin	4,81
Sistin	0,32
Tirosin	1,62

LSIH Malang and Farnasi Unair Surabaya

1) The first phase of the study: Measuring the rats memory special using MWM

After 8 weeks of treatment, all of the rats models were examined by MWM to measure their spati⁶ memory, in the form of cylindrical drum-shaped pool measuring 1.8 m by 0.5 m high¹⁰ ed with water to a depth of 0.3 m. A circular platform wi¹⁰ diameter of 13 cm and a height of 28 cm was placed 2 cm below the water surface. To make the platform invi²¹ e, coconut milk is added to the water. Drum surface is divided into four quadrants A, B, C, and

D. The platform is placed on a particular quadrant area (and not changed its location during the experiment) (Fig. 1).



Figure 1. MWM.

Spatial memory was compared among control and treatment group. All rats were trained (escape latency) by MWM method hidden platform test for 8 days to calculated its time to reaches the platform. Each rat got trained twice a day [6].

MWM measurement system for the escape latency: Everytime when the experiment started set the starting point when rat is placed in the pool (e.g in position A). The platform has been placed in position between A and D quadrant area. Then the rats released at the specific¹³ starting point, facing the drum wall, rats were left to swim to reach the platform and climb onto the platform. The time that needed to reach the platform (escape latency) was recorded. After the rats reached the platform, the rats were given time to rest for 30 seconds.

After that, rats are dried and rested to prepare for the next experiments. Everyday, the rat was trained twice a day with duration 90 seconds. If it failed in 90 seconds, it would be directed into platform and let it took a rest for 30 seconds. The failed rat has its escape latency noted as 90 seconds. After that, rats are dried and rested to prepare for the next experiments. The second examination, it would be placed at the different location (e.g potition C). Rats put in place and it swimmmed to the platform and up onto the platform. Time to reached the platform was noted (escape latency). After reaching platform, the rat was rested for 30 seconds.




Figure 2. Probe test

⁶ To measure the rats spatial memory, it examined by probe test. Platform was removed from the water, the other components were allowed like before. Starting place was chosen randomly The rat swimmmed for 90

seconds and the percentation for swimming to target quadrans was calculated (quadrant previously laid the platform) compared to overall time taken by the rat through the entire quadrant (performance in this case 90 seconds).

The drum was given rope that connected each position (A-C) and (13-D). 2 observer observed the length of time (seconds) that rats spent in the target quadrant (in this case between quadrant A and D). Observations using the MWM probe test is performed one time in morning at 9:00 AM to finish.

2) The second phase of the study: Calculating the PPAR expression

After completing the rat spatial memory measurement, in order to make the brain tissue preparations rats is killed. The sample that cut then stained by immunohistochemical staining method. The percentage of PPAR α and PPAR γ expression is calculated by counting the number of cells which given a brown color to the core compared to the number of cells in each field of view. From the five visual fields were taken and then calculated the average number of cells expressing the brown color

3) Data analysis

Data was tested using Shapiro Wilk. If data in normal distribution, continued by one way Anova test with 95% significances post hoc test (LSD).

III. RESULT

The first phase study showed the percentage of the time length in target area from the P2 group the highest compared to the K and P1 group, respectively 33.52%, 29.9%, and 23.15%

The result showed that there were significant differences for rats to stay at platform among control and treatment group (p=0,021).




Figure 3. The average percentage of time in the target quadrant (probe test) after treatment for 8 weeks.

Advanced analysis test used post hoc test (LSD), it showed significant differences between P1 and P2 (p=0,006). There was no significant differences between K and P1. It means that Seluang fish formula increasing rat's memory showed with the highest time length in target area.

On the PPAR α and PPAR γ examination showed the average percentage of neuron cells that express PPAR α in P2 group is higher than K and P1 group, respectively 18.5%, 3.75% and 2.5%. The mean percentage of neuron cells that express PPAR γ in P2 group also higher than K and P1 group, respectively 35.41%, 6.2% and 5.83%.

The Kruskal Wallis Statistical test results showed that there were significant differences in average percentage of neurons that express PPAR α were examined between 3 groups (p = 0.002). The same test results the percentage of neuronal cells that express PPAR γ showed significantly different in average percentage of neuron cells that express PPAR γ were examined between 3 groups (p = 0.001)




Figure 4. Neuron cells expressing PPAR α .




Figure 5. Neuron cells expressing PPAR γ




Figure 6. The average percentage of neuron cells expressing PPAR α and PPAR γ

IV. DISCUSSION

Hajja *et al* [4] reported with administrating higher dietary omega-3 fatty acids from the omega-6 (ratio of omega-6 : omega-3 low) improving spatial memory and

up-regulation of PPAR α and PPAR γ gene expression in the hippocampus. This suggests that PPAR α and PPAR γ as transcription factors can influence the synaptic proteins expression that trigger the synaptic plasticity and improves spatial memory in animals that fed high dietary in omega-3 fatty acids. Peroxisome Proliferator-Activated Receptors (PPARs) are a subset of the nuclear hormone receptor super family that has transcriptional activity, modulated by ligand-receptor interaction. There are three families of PPARs known as PPAR α , PPAR γ and PPAR β / δ . They bind to the peroxisome proliferator response elements are similar but show differences in transactivating function, which is partially mediated by the distribution network, and coactivator ligand specificity [5]-[7]. Peroxisome proliferator-activated receptors expressed in various tissues, among others, liver cells, kidney, and brain. In brain tissue PPARs are expressed in a number of brain cells, especially in the hippocampus [8].

In this study, groups of rats were fed Seluang Formula showed higher neurogenesis in hippocampus that express PPAR α and PPAR γ compared to the control group and the low-protein group. It is proven that the nutrients contained in Seluang fish especially unsaturated fatty acids such as essential fatty acids linoleic, oleic and DHA can trigger the PPAR α and PPAR γ expression. PPAR α plays an important role in the biosynthesis regulation of acetylcholine that contribute to cognitive function, while PPAR γ has a prominent role in the central nervous system inflammatory regulation and neuroprotective, triggering the cognitive performance. Several studies showed that cognitive performance can be improved through the PPAR nuclear receptors [4]. Kuang *et al* (2012) reported that preactivation of PPAR β can improve spatial memory and cognitive functions as well as changes cytomorphologi neuron cells hippocampus after sustained global cerebral ischemia-reperfusion injury (GCIRI). Meanwhile another study reported that PPAR γ able to up-regulation of Bcl-2 anti-apoptotic protein in neurons and induces mitochondrial stabilization also protected from oxidative stress and apoptosis in cultured neuron cells [7].

Other studies have reported that chronic deficiency of alpha linolenic fatty acids can cause changes in cholinergic parameters by increasing spontaneous release of acetylcholine (ACh) in the hippocampus and lead to a lower release which associated with less binding to the muscarinic receptors. The responses can be different based on the excessive loss of neuronal DHA and n-6 PUFA, particularly by the septo-hippocampal cholinergic system which plays a role in the cognitive function regulation. Such modifications may contribute to cognitive impairment caused by the deficiency of n-3 PUFAs [8].

In a other study about the gyrus dentatus granule cells and the hippocampus CA3 pyramidal cells, suggesting that protein malnutrition effected on all of these cells, the dominant effect is on the gyrus dentatus cells [9]. In the same result associated with protein malnutrition showed a decrease in prenatal neurogenesis in the hippocampus

granule cells dentatus formation [10].

Formation hippocampus is involved in learning and memory which are ruled the construction of spatial cognitive map to surrounding environment. This map is used by the animal to identify the differences in place (situation). The information that viable will be organized into working memory as short-term memory. In the spatial cognition study, Morris water maze test is often used as a tool [11]. If an interruption occurred in the hippocampus formation caused by malnutrition, can impact on spatial memory. Another study found that malnutrition in early life caused a decrease in hippocampus neural progenitor and also deficiency recognition object in adulthood [12]. Malnutrition adverse effects on cognitive development can occur as a result of neuron structural and functional maturation disruption such as myelination delayed and dendrite arborization decreased [13]. One theory about the involvement of protein synthesis in learning and memory is the involvement of cyclic adenosine monophosphate / Protein Kinase A (cAMP / PKA) that mediate the signaling of learning and memory process. The connection between protein synthesis and memory formation depends on synaptic facilitation. The important role of cAMP factor response binding protein (CREB) in the form of plasticity and sensitive to apamin forming protein, is also an important role of NMDA receptors in the signaling and stimulation of gene expression to form protein will produce synaptic plasticity and long-term memory [14].

V. CONCLUSION

The conclusion of this study is Seluang fish has the complete nutrition composition that could increase the spatial memory, expression of PPAR α and PPAR γ that lead to intelligence increased.

ACKNOWLEDGMENT

We would like to thank to Biochemistry Laboratory, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia, and Faculty of Medicine Brawijaya University, Malang, Indonesia for all of support and good teamwork for this research.

REFERENCES

- [1] Riset Kesehatan Dasar, Kementerian Kesehatan RI, Jakarta, 2010.
- [2] Tingkat konsumsi ikan di Kalimantan Selatan, Dinas Perikanan dan Kelautan Provinsi Kalimantan Selatan, Banjarmasin, 2010.
- [3] Rencana Aksi Nasional Pangan dan Gizi 2011-2015, Kementerian Perencanaan Pembangunan Nasional/Badan Perencanaan Pembangunan Nasional (BAPPENAS), Jakarta, 2011.
- [4] T. Hajjar, G. Y. Meng, M. A. Rajio, S. Vidyadaran, F. Othman, A. S. Farjam, *et al.*, "Omega 3 polyunsaturated fatty acid improves spatial learning and hippocampal peroxisome proliferator activated receptors (PPAR α and PPAR γ) gene expression in rats," *BMC Neuroscience*, vol. 13, pp. 109, 2012.
- [5] R. F. Morrison and S. R. Farmer, "Hormonal signaling and transcriptional control of adipocyte differentiation," *Journal of Nutrition*, vol. 22, pp. 3116S-321S, 2000.
- [6] H. Wang, R. Jiang, Q. He, Y. Zhang, Y. Zhang, Y. Li, *et al.*, "Expression pattern of peroxisome proliferator-activated receptors

in rat hippocampus following cerebral ischemia and reperfusion injury," *PPAR Research*, vol. 2012, no. 11, pp. 239-261, 2012.

- [7] G. Kuang, Q. He, Y. Zhang, R. Zhuang, A. Xiang, Q. Jiang, *et al.*, "Modulation of preactivation of PPAR- β on memory and Learning dysfunction and inflammatory response in the hippocampus in rats exposed to global cerebral ischemia/reperfusion," *PPAR Research*, vol. 2012, 2012.
- [8] S. Aid, S. Vancassel, C. Poumes-Ballihaut, S. Chalon, P. Guesnet, and M. Lavielle, "Effect of a diet-induced n-3 PUFA depletion on cholinergic parameters in the rat hippocampus," *J. Lipid Res*, vol. 44, pp. 1545-1551, 2003.
- [9] L. Cintra, A. Aguilar, L. Granados, A. Galvan, T. Kemper, W. DeBassio, *et al.*, "Effects of prenatal protein malnutrition on hippocampal CA1 pyramidal cells in rats of four age groups," *Hippocampus*, vol. 7, pp. 192-203, 1997.
- [10] W. A. Debassio, T. L. Kemper, J. R. Galler, and J. Tonkiss, "Prenatal malnutrition effect on pyramidal and granule cell generation in the hippocampal formation," *Brain Research Bulletin*, vol. 35, no. 1, pp. 57-61, 1994.
- [11] C. T. Valadares, M. T. H. Fukuda, A. L. Francolin-Silva, A. S. Hernandesdan, and S. S. Almeida, "Effects of post natal protein malnutrition on learning and memory procedures," *Nutritional Neuroscience*, vol. 13, no. 6, pp. 274-282, 2010.
- [12] M. A. De Godoy, A. S. de Souza, M. A. Lobo, O. V. K. Sampaio, L. Moraesa, M. R. Balanza, *et al.*, "Effects of protein restriction during gestation and lactation on cell proliferation in the hippocampus and subventricular zone: Functional implications. Protein restriction alters hippocampal/SVZ cell proliferation," *Brain Research*, vol. 1496, pp. 10-27, 2013.
- [13] B. R. Kar, S. L. Rao, and B. A. Chandramouli, "Cognitive development in children with chronic protein energy malnutrition," *Behavioral and Brain Functions*, vol. 4, pp. 31, 2008.
- [14] P. J. Hernandez and T. Abel, "The role of protein synthesis in memory consolidation: Progress amid decades of debate," *Neurobiol Learn Mem*, vol. 89, no. 3, pp. 293-311, 2008.



Ary Yunanto was born in Salatiga, 24th November 1952. He was graduated from Medicine Faculty of Diponegoro University, Semarang, Indonesia, in 1978. He studied as Pediatrician also in Medicine Faculty of Diponegoro University, Semarang, Indonesia, and graduated in 1989. He continued to study doctoral degree of Brawijaya University, Malang, Indonesia, in 2009 and graduated in 2012.

He worked as Pediatrician in Ulin Hospital, Banjarmasin, Indonesia, and Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia, until now. He ever published his research in IOSR Journal of dental and medical sciences (JDSM) in 2012. He also ever presented his research in Manado (Indonesia) in 2012, in 14th pediatric congress in Singapore in 2011, and Perinasia Congress in Balikpapan in 2009. He wrote some books such as "Kapita Selektta Biokimia: Peran Radikal Bebas Pada Intoksikasi dan Patobiologi Penyakit" in 2009, "Pedoman Penatalaksanaan Medis Neonatologi, Sepsis neonatal in 2011, tinjauan

dari aspek biomolekuler" in 2012, and "Saliva Sebagai dasar diagnosis sepsis neonatal" in 2012.



Lambung Mangkurat University, Banjarmasin, Indonesia.

Didik Dwi Sanyoto was born in Pelaihari, Indonesia, in March 1972. He was graduated as physician in 1998 from Lambung Mangkurat University, Banjarmasin, Indonesia. He was graduated master degree 2003 from Airlangga University, Surabaya, Indonesia. He currently work in Medical Anatomy Department, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia. Now, he is studying doctoral



Meitria Syahadatina Noor was born in Surabaya, Indonesia, on 19th May 1979. She was graduated as physician in 2005 from Lambung Mangkurat University, Banjarmasin, Indonesia. She was graduated master degree 2010 from Airlangga University, Surabaya, Indonesia, shee currently work in Public Health Department, Faculty of Medicine, degra-e in Airlangga University, Surabaya, Indonesia.



University, Malang, Indonesia. She currently work in Pathology of Anatomy Department, Faculty of Medicine, Lambung Mangkurat University/Ulin Hospital, Banjarmasin, Indonesia.

Ika Kustiyah, was born in Banjarmasin, Indonesia, on 12th October 1968. She was graduated as physician in 1994 from Diponegoro University, Semarang, Indonesia. She was graduated master degree in 2003 and Pathologist of Anatomy in 2004 from Diponegoro University, Semarang, Indonesia. She was graduated from doctoral degree in Biomedicine in 2012 from Brawijaya



Department, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia.

Triawanti was born in Surabaya, Indonesia, in September 1971. She was graduated as physician in 1998 from Lambung Mangkurat University, Banjarmasin, Indonesia. She was graduated master degree in 2002 from Airlangga University, Surabaya, Indonesia, PhD 2012 from Brawijaya University, Malang, Indonesia. She currently work in Medical Biochemistry

ORIGINALITY REPORT

17%

SIMILARITY INDEX

7%

INTERNET SOURCES

13%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

-
- | | | |
|---|--|----|
| 1 | Submitted to Academia Sinica Life Science Library
Student Paper | 2% |
| 2 | "20th International Congress of Nutrition: Granada, Spain, September 15 20, 2013", Annals of Nutrition and Metabolism, 2013
Publication | 1% |
| 3 | Hernandez, P.J.. "The role of protein synthesis in memory consolidation: Progress amid decades of debate", Neurobiology of Learning and Memory, 200803
Publication | 1% |
| 4 | www.jlr.org
Internet Source | 1% |
| 5 | Submitted to Udayana University
Student Paper | 1% |
| 6 | Fitting, S.. "Neonatal intrahippocampal injection of the HIV-1 proteins gp120 and Tat: Differential effects on behavior and the relationship to stereological hippocampal measures", Brain | 1% |
-

Research, 20080926

Publication

-
- | | | |
|---|---|----|
| 7 | Submitted to Ramapo College
Student Paper | 1% |
|---|---|----|
-
- | | | |
|---|---|----|
| 8 | Submitted to Concordia University Wisconsin
Student Paper | 1% |
|---|---|----|
-
- | | | |
|---|--|----|
| 9 | Francisco J. Torres-Espínola, Signe Altmäe, Maria Teresa Segura, Antonio Jerez et al. "Maternal PPARG Pro12Ala polymorphism is associated with infant's neurodevelopmental outcomes at 18months of age", Early Human Development, 2015
Publication | 1% |
|---|--|----|
-
- | | | |
|----|--|----|
| 10 | Fitting, S.. "Effects of chronic adult dietary restriction on spatial learning in the aged F344xBN hybrid F1 rat", Physiology & Behavior, 20080227
Publication | 1% |
|----|--|----|
-
- | | | |
|----|---------------------------------------|----|
| 11 | univmed.org
Internet Source | 1% |
|----|---------------------------------------|----|
-
- | | | |
|----|---|----|
| 12 | www.frontiersin.org
Internet Source | 1% |
|----|---|----|
-
- | | | |
|----|---|----|
| 13 | Submitted to King's College
Student Paper | 1% |
|----|---|----|
-
- | | | |
|----|--|----|
| 14 | jatp.ift.or.id
Internet Source | 1% |
|----|--|----|

15

Submitted to Universiti Putra Malaysia

Student Paper

1%

16

www.cambridge.org

Internet Source

1%

17

Submitted to Universitas Airlangga

Student Paper

<1%

18

paradisevillagechiro.com

Internet Source

<1%

19

www.ijcea.org

Internet Source

<1%

20

Norio Amizuka. "Assessment of histological alterations in cartilage and extracellular matrix driven by collagen-induced arthritis in *Macaca fascicularis*", *Arthritis Research & Therapy*, 2012

Publication

<1%

21

Tianda Feng, Yunhui Liu, Chao Li, Zhen Li. "Protective Effects of Nigranoic Acid on Cerebral Ischemia–Reperfusion Injury and its Mechanism Involving Apoptotic Signaling Pathway", *Cell Biochemistry and Biophysics*, 2014

Publication

<1%

22

www.physiology.org

Internet Source

<1%

Submitted to University of New England

23

Student Paper

<1%

24

www.jneurosci.org

Internet Source

<1%

25

www.jourlib.org

Internet Source

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On