

# Detection and genetic characterization of human enterovirus A71 from hand, foot and mouth disease cases in Banjarmasin, Indonesia. Tahun Terbit 2018

*by* Edi Hartoyo

---

**Submission date:** 15-Nov-2020 10:03AM (UTC+0700)

**Submission ID:** 1446236370

**File name:** h\_disease\_cases\_in\_Banjarmasin,\_Indonesia.\_Tahun\_Terbit\_2018.pdf (249.92K)

**Word count:** 2578

**Character count:** 13806

## RESEARCH NOTE

# 4 DETECTION AND GENETIC CHARACTERIZATION OF HUMAN ENTEROVIRUS A71 FROM HAND, FOOT AND MOUTH DISEASE CASES IN BANJARMASIN, INDONESIA

Ageng Wiyatno<sup>1</sup>, Edi Hartoyo<sup>2,3</sup>, Ungke Anton Jaya<sup>4</sup>, Chairin Nisa Ma'roef<sup>1</sup>, Khin Saw Myint<sup>1</sup>, Amethyst Gillis<sup>5</sup>, Corina Monagin<sup>5,6</sup> and Dodi Safari<sup>1</sup>

<sup>1</sup>Eijkman Institute for Molecular Biology, Jakarta; <sup>2</sup>Faculty of Medicine, Lambung Mangkurat University South, Kalimantan; <sup>3</sup>Ulin General Hospital, Banjarmasin, South Kalimantan; <sup>4</sup>Eijkman <sup>12</sup>xford Clinical Research Unit, Jakarta, Indonesia; <sup>5</sup>Metabiota, San Francisco; <sup>6</sup>One Health Institute, School of Veterinary Medicine, University of California, Davis, CA, USA

**Abstract.** Enterovirus A71 (EV-A71) infection associated with <sup>10</sup>hand, foot and mouth disease (HFMD) has been recognized as a major health threat especially in children. The aim of this study was to identify and genetically characterize the viral etiological agent associated with HFMD. Using molecular methods, EV-A71 sub-genogroup B5 was detected as the causative agent in 7 out of 13 archived throat swab specimens collected during the 2016 HFMD outbreak in Banjarmasin, South Kalimantan, Indonesia.

**Keywords:** enterovirus A71, hand, foot and mouth disease, Indonesia

### INTRODUCTION

<sup>11</sup>Enterovirus 71 (EV-A71), an RNA virus of family Picornaviridae, is a common cause of hand, foot and mouth disease (HFMD) (Solomon *et al*, 2010). The infection usually manifests with fever, rash, blisters and oral lesions, and in severe cases may proceed to encephalitis, aseptic meningitis, pancreatitis, cardiopulmonary complications and poliomyelitis-like

paralysis (Wu *et al*, <sup>17</sup>2; Ooi *et al*, 2010; Zhang *et al*, 2016). The virus is highly transmissible via a fecal-oral route and respiratory droplets and is recognized as a global public health threat with a potential to cause pandemics similar to poliovirus (Yip *et al*, 201<sup>3</sup>). EV-A71, together with coxsackievirus CV-A6 and CV-A16, attract the most public health because these three viruses are the major <sup>9</sup> common causes of HFMD. However, compared to CV-A6 and CV-A16, EV-A71 has a stronger association with CNS involvement, such as acute flaccid paralysis (Long *et al*, 2016; Suresh *et al*, 2017).

EV-A71 epidemics have occurred with increasing numbers in the Asia-Pa-

Correspondence: Ageng Wiyatno, Eijkman Institute for Molecular Biology, Jakarta 10430, Indonesia.

Fax: +62213147982

Email: ageng@eijkman.go.id

16  
cific region, including Hong Kong, Japan, mainland China, Malaysia, Singapore, Taiwan, and Thailand for the past two decades (Shimizu *et al*, 1999; Fujimoto *et al*, 2002; Lin *et al*, 2003; Chua *et al*, 2007; Wu *et al*, 2010; Mauleekoonphairoj *et al*, 2015). In the Southeast Asian region, HFMD usually occur during the transition from rainy to dry season and the numbers have increased with large temperature fluctuations (Hii *et al*, 2011). A study in Indonesia between 2008-2012 detected EV-A71 as a cause of mild HFMD across the country with a prevalence of 6.5% (Susanti *et al*, 2014).

Virulence of EV-A71 depends on multiple factors, with host and environmental factors appearing to be the most significant (Shih *et al*, 2011; Chang *et al*, 2012). Mutations in VP1 gene, which encodes the capsid protein has reported to be associated with EV-A71 virulence, resistance and adaptation in *in-vitro* studies (Shih *et al*, 2004; Chen *et al*, 2008; Kelly *et al*, 2015). Despite being endemic, studies on the etiological agent of HFMD and genetic characterization of the virus are limited in Indonesia. This study aimed to describe the investigation on HFMD cases caused by EV-A71 in Banjarmasin, Indonesia in 2016.

## MATERIALS AND METHODS

### Samples

This study was conducted retrospectively on specimens from HFMD cases collected between November 2015 to February 2016 at pediatric wards in Ulin General Hospital and Islamic Hospital, Banjarmasin, South Kalimantan, Indonesia. Throat swab specimens were from 13 hospitalized patients presenting with fever ( $\geq 38.5^{\circ}\text{C}$ ) and typical HFMD symptoms, viz. rash and blisters in the

mouth, hands and feet. Samples were eluted from swabs using Viral Transport medium prepared from brain heart infusion (Oxoid, Hampshire, UK) and penicillin-streptomycin (Gibco, Carlsbad, CA) and stored at  $-80^{\circ}\text{C}$  until shipped to the Eijkman Institute, Jakarta for viral molecular testing.

The study was approved by the Eijkman Institute for Molecular Biology Research Ethics Commission (no. 66, 18 November 2013). Residential locations of the cases were based on street addresses provided by the patients.

### PCR-based virus genotyping assay

5  
Viral RNA was extracted using QIAamp Viral RNA Mini Kit (Qiagen, Hilden, Germany) to obtain 60  $\mu\text{l}$  of RNA, from which 5  $\mu\text{l}$  aliquot was taken for cDNA synthesis employing Go Script Reverse Transcription System (PROMEGA, Madison, WI) followed by PCR with Go Taq Green Master Mix (PROMEGA) pan-enterovirus primers targeting the 5'-untranslated region (Wiyatno *et al*, 2016) and primers targeting  $\beta$ -actin gene as internal control. In brief, the reaction mixture (25  $\mu\text{l}$ ) contained 2  $\mu\text{l}$  cDNA template, 400  $\mu\text{M}$  forward and reverse primers, 1.5 mM  $\text{MgCl}_2$ , 200  $\mu\text{M}$  dNTPs, and buffer containing DNA polymerase (PROMEGA, Madison, WI). Thermocycling was conducted in Proflex PCR instrument (Applied Biosystems, Carlsbad, CA) as follows: 95 $^{\circ}\text{C}$  for 15 minutes; 16 cycles of 94 $^{\circ}\text{C}$  for 45 seconds, 65 $^{\circ}\text{C}$  for 45 seconds (-1 $^{\circ}\text{C}$ /cycle) and 72 $^{\circ}\text{C}$  for 45 seconds; 32 cycles of 98 $^{\circ}\text{C}$  for 45 seconds, 48 $^{\circ}\text{C}$  for 45 seconds, and 72 $^{\circ}\text{C}$  for 45 seconds; and a final step at 72 $^{\circ}\text{C}$  for 5 minutes. Amplicons (~400 bp) were detected by 1.5% agarose gel-electrophoresis and staining with SYBR Safe (Invitrogen, Carlsbad, CA), directly sequenced at Eijkman Institute,

Jakarta and analyzed using Geneious Software V.8. (Biomatters, Auckland, NZ). The presence of EV-A71 was confirmed by amplifying a 891-bp region of the VP1 gene followed by sequencing for phylogenetic analysis (Ap *et al*, 2011; Ling *et al*, 2014). In short, the PCR mixture (25  $\mu$ l) contained 2  $\mu$ l cDNA template, 400  $\mu$ M upstream and downstream primers, 1.5 mM MgCl<sub>2</sub>, 200  $\mu$ M dNTPs and buffer containing DNA polymerase (PROMEGA, Madison, WI), and amplified using the following thermocycling conditions: 95°C for 10 minutes; 44 cycles of 94°C for 45 seconds, 50°C for 45 seconds and 72°C for 1 minute; with a final step at 72°C for 5 minutes. Amplicons were processed for sequencing as described above.

All sequences were submitted to GenBank database with accession numbers KY041858, KY041859, KY041860, KY041861, KY041862, KY041863, and KY041864. All enterovirus negative samples were also tested using consensus PCR for other viral causes, such as herpesvirus (terminase gene; 419 bp) and paramyxoviruses (polymerase gene; 561 bp) (Chmielewicz *et al*, 2001; Tong *et al*, 2008).

## RESULTS

Throat swab specimens came from six male and seven female children with median age of 16 months (range 8-37 months). In addition to the characteristic HFMD lesions, other symptoms, such as cough, conjunctivitis, diarrhea, seizure and vomiting were also observed in some patients. There were no CNS complications or mortality. Nine HFMD cases resided in southeastern Banjarmasin, with the remaining in the northwestern region of the city.

Seven HFMD patients were positive by pan-enterovirus PCR and one by

paramyxovirus PCR (data not shown). Sequence analysis of the enterovirus-specific amplicons revealed all seven positives were EV-A71. Further amplification and sequencing of VP1 region demonstrated that all EV-A71 sequences had 96-98% similarity with strains from Malaysia, Taiwan, Thailand, and Vietnam, with all isolates belonging to the subgenogroup B5. Intratypic similarity among the sequences ranged from 93.7% to 99.9%. A dendrogram constructed from 891 nt fragments of the VP1 region showed five isolates clustered with strains from mainland China and Taiwan, and the remaining with strains from Malaysia, Thailand and Vietnam (Fig 1). No co-detection with other enteroviruses were found.

## DISCUSSION

In this study, we identified EV-A71 associated with a 2016 outbreak of HFMD in Banjarmasin, South Kalimantan. Although both hospitals where the patients were admitted were located in northeast Banjarmasin, cases came from other parts of the city. Human EV-A71 is considered endemic in Southeast Asia, with outbreaks occurring in cycles every 2-3 years in Japan, Malaysia and Taiwan and annually in China (Sabanathan *et al*, 2014). CV-A6 and CV-A16 are known to co-exist with EV-A71 especially in cases with CNS manifestations (Solomon *et al*, 2010; Ling *et al*, 2014; Gaunt *et al*, 2015) and recombination between EV-A71 and CV-A16 is linked with pathogenicity of the virus (Zhang *et al*, 2010). Only a single enterovirus strain was detected from swab of each HFMD patient.

The EV-A71 strains isolated in this study belonged to subgenogroup B5, the most prevalent strain in the region (Ooi *et al*, 2007; Ap *et al*, 2011; Hassel *et al*, 2015).

HUMAN ENTEROVIRUS A71 IN BANJARMASIN, INDONESIA




Fig 1-Phylogenetic tree of 891-bp enterovirus (EV)-A71 VP1 gene fragments isolated from seven hand, foot and mouth disease (HFMD) samples from Banjarmasin, South Kalimantan, Indonesia. Human coxsackievirus (CVA)A16 is used as the outgroup. GenBank accession number is indicated for each EV strain. Genetic relationship was analyzed using Molecular Evolutionary Genetics Analysis (MEGA) version 6.06 (<http://www.megasoftware.net/mega.php>) using neighbor-joining method and applying Kimura 2-parameter model with 1,000 bootstraps. Number at branch point denotes percent similarity. Scale bar indicates percent nucleotide substitution change.

In the Asia-Pacific region, subgenogroup B5 together with C4 have recently replaced subgenogroups B3 and B4 (Chang *et al*, 2016). Subgenogroup C4 is now recognized as the major subgenogroup causing severe illness particularly associated with CNS manifestations and high mortality.

To the best of our knowledge, this study is the first genetic characterization of EV-A71 from a HFMD outbreak in Banjarmasin, South Kalimantan, Indonesia. Our study should increase the awareness of EV-A71 as an etiological agent of HFMD in Indonesia. Further studies of genetic characterization are important to identify additional causative agents of HFMD and subgenotypes to understand the epidemiology of outbreaks.

#### ACKNOWLEDGEMENTS

The authors thank Ms Aghnianditya Dewantari for assistance in molecular testing of viral panels.

#### REFERENCES

- Ap M, Yusof I, Rais F, *et al*. Molecular epidemiology of human enterovirus 71 (HEV71) strains isolated in Peninsular Malaysia and Sabah from year 2001 to 2009. *J Gen Mol Virol* 2011; 3: 18-26.
- Chang H-L, Chio C-P, Su H-J, *et al*. The association between enterovirus 71 infections and meteorological parameters in Taiwan. *PLOS One* 2012; 7: e46845.
- Chang P-C, Chen S-C, Chen K-T. The current status of the disease caused by enterovirus 71 infections: epidemiology, pathogenesis, molecular epidemiology, and vaccine development. *Int J Environ Res Public Health* 2016 Sep 9; 13 (9). pii:E890
- Chen T-C, Liu S-C, Huang P-N, Chang H-Y, Chern J-H, Shih S-R. Antiviral activity of pyridyl imidazolidinones against enterovirus 71 variants. *J Biomed Sci* 2008; 15: 291-300.
- Chmielewicz B, Goltz M, Ehlers B. Detection and multigenic characterization of a novel gammaherpesvirus in goats. *Virus Res* 2001; 75: 87-94.
- Chua KB, Chua BH, Lee CSM, *et al*. Genetic diversity of enterovirus 71 isolated from cases of hand, foot and mouth disease in the 1997, 2000 and 2005 outbreaks, Peninsular Malaysia. *Malays J Pathol* 2007; 29: 69-78.
- Fujimoto T, Chikahira M, Yoshida S, *et al*. Outbreak of central nervous system disease associated with hand, foot, and mouth disease in Japan during the summer of 2000: detection and molecular epidemiology of enterovirus 71. *Microbiol Immunol* 2002; 46: 621-7.
- Gaunt E, Harvala H, Österback R, *et al*. Genetic characterization of human coxsackievirus A6 variants associated with atypical hand, foot and mouth disease: a potential role of recombination in emergence and pathogenicity. *J Gen Virol* 2015; 96, 1067-79.
- Hassel C, Mirand A, Lukashev A, *et al*. Transmission patterns of human enterovirus 71 to, from and among European countries, 2003 to 2013. *Euro Surveill* 2015; 20: 30005.
- Hii YL, Rocklöv J, Ng N. Short term effects of weather on hand, foot and mouth disease. *PLOS One* 2011; 6: e16796.
- Kelly JT, De Colibus L, Elliott L, *et al*. Potent antiviral agents fail to elicit genetically-stable resistance mutations in either enterovirus 71 or coxsackievirus A16. *Antiviral Res* 2015; 124: 77-82.
- Lin T-Y, Twu S-J, Ho M-S, Chang L-Y, Lee C-Y. Enterovirus 71 outbreaks, Taiwan: occurrence and recognition. *Emerg Infect Dis* 2003; 9: 291-3.
- Ling BP, Jalilian FA, Harmal NS, Yubbu P, Sekawi Z. Detection and characterization of viruses causing hand, foot and mouth disease from children in Seri Kembangan, Malaysia. *Trop Biomed* 2014; 31: 654-62.
- Long L, Xu L, Xiao Z, *et al*. Neurological complications and risk factors of cardiopulmo-

- nary failure of EV-A71-related hand, foot and mouth disease. *Sci Rep* 2016; 6: 23444.
- Mauleekoonphairoj J, Puenpa J, Korkong S, Vongpunsawad S, Poovorawan Y. Prevalence of human enterovirus among patients with hand, foot, and mouth disease and herpangina in Thailand, 2013. *Southeast Asian J Trop Med Public Health* 2015; 46: 1013-20.
- Ooi MH, Wong SC, Podin Y, *et al.* Human enterovirus 71 disease in Sarawak, Malaysia: a prospective clinical, virological, and molecular epidemiological study. *Clin Infect Dis* 2007; 44: 646-56.
- Ooi MH, Wong SC, Lewthwaite P, Cardoso MJ, Solomon T. Clinical features, diagnosis, and management of enterovirus 71. *Lancet Neurol* 2010; 9: 1097-105.
- Sabanathan S, Tan LV, Thwaites L, Wills B, Qui PT, Rogier van Doorn H. Enterovirus 71 related severe hand, foot and mouth disease outbreaks in South-East Asia: current situation and ongoing challenges. *J Epidemiol Community Health* 2014; 68: 500-2.
- Shih S-R, Tsai M-C, Tseng S-N, *et al.* Mutation in enterovirus 71 capsid protein VP1 confers resistance to the inhibitory effects of pyridyl imidazolidinone. *Antimicrob Agents Chemother* 2004; 48: 3523-9.
- Shih S-R, Stollar V, Li M-L. Host factors in enterovirus 71 replication. *J Virol* 2011; 85: 9658-66.
- Shimizu H, Utama A, Yoshii K, *et al.* Enterovirus 71 from fatal and nonfatal cases of hand, foot and mouth disease epidemics in Malaysia, Japan and Taiwan in 1997-1998. *Jpn J Infect Dis* 1999; 52: 12-5.
- Solomon T, Lewthwaite P, Perera D, Cardoso MJ, McMinn P, Ooi MH. Virology, epidemiology, pathogenesis, and control of enterovirus 71. *Lancet Infect Dis* 2010; 10: 778-90.
- Suresh S, Forgie S, Robinson J. Non polio enterovirus detection with acute flaccid paralysis: a systematic review. *J Med Virol* 2018; 90: 3-7.
- Susanti N, Herna, Purnamawati S, Setiawaty V. Detection of cause and distribution of hand foot and mouth diseases (HFMD) outbreaks in 2008-2012. *J Biotek Med Indones* 2014; 3: 77-84.
- Tong S, Chern S-WW, Li Y, Pallansch MA, Anderson LJ. Sensitive and broadly reactive reverse transcription-PCR assays to detect novel paramyxoviruses. *J Clin Microbiol* 2008; 46: 2652-8.
- Wiyatno A, Antonjaya U, Ma'roef CN, *et al.* Detection and identification of coxsackievirus B3 from sera of an Indonesian patient with undifferentiated febrile illness. *J Infect Dev Ctries* 2016; 10: 880-3.
- Wu J-M, Wang J-N, Tsai Y-C, *et al.* Cardiopulmonary manifestations of fulminant enterovirus 71 infection. *Pediatrics* 2002; 109: E26-.
- Wu Y, Yeo A, Phoon MC, *et al.* The largest outbreak of hand, foot and mouth disease in Singapore in 2008: the role of enterovirus 71 and coxsackievirus A strains. *Int J Infect Dis* 2010; 14: e1076-81.
- Yip CCY, Lau SKP, Woo PCY, Yuen K-Y. Human enterovirus 71 epidemics: what's next? *Emerg Health Threats J* 2013; 6: 10.3402/ehtj.v6i0.19780.
- Zhang Y-F, Deng H-L, Fu J, Zhang Y, Wei J-Q. Pancreatitis in hand-foot-and-mouth disease caused by enterovirus 71. *World J Gastroenterol* 2016; 22: 2149-52.

# Detection and genetic characterization of human enterovirus A71 from hand, foot and mouth disease cases in Banjarmasin, Indonesia. Tahun Terbit 2018

## ORIGINALITY REPORT

13%

SIMILARITY INDEX

10%

INTERNET SOURCES

8%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://rupress.org">rupress.org</a> Internet Source	1%
2	Yu-Wei Chang, Wei-Lun Chiang, Wen-Hung Wang, Chun-Yu Lin, Ling-Chien Hung, Yi-Chang Tsai, Jau-Ling Suen, Yen-Hsu Chen. "Google Trends-based non-English language query data and epidemic diseases: a cross-sectional study of the popular search behaviour in Taiwan", BMJ Open, 2020 Publication	1%
3	Submitted to essex Student Paper	1%
4	<a href="http://jbiomedsci.biomedcentral.com">jbiomedsci.biomedcentral.com</a> Internet Source	1%
5	<a href="http://article.sapub.org">article.sapub.org</a> Internet Source	1%
6	M. Arita. "Temperature-sensitive mutants of enterovirus 71 show attenuation in cynomolgus monkeys", Journal of General	1%



---

7	<a href="http://nthur.lib.nthu.edu.tw">nthur.lib.nthu.edu.tw</a> Internet Source	1%
8	Meng, Tao, Tanja K Kiener, and Jimmy Kwang. "RNA polymerase I-driven reverse genetics system for enterovirus 71 and its implications for vaccine production", <i>Virology Journal</i> , 2012. Publication	1%
9	Ang, Li Wei, Joanne Tay, Meng Chee Phoon, Jung Pu Hsu, Jeffery Cutter, Lyn James, Kee Tai Goh, and Vincent Tak-Kwong Chow. "Seroepidemiology of Coxsackievirus A6, Coxsackievirus A16, and Enterovirus 71 Infections among Children and Adolescents in Singapore, 2008-2010", <i>PLoS ONE</i> , 2015. Publication	1%
10	<a href="http://bmcpublichealth.biomedcentral.com">bmcpublichealth.biomedcentral.com</a> Internet Source	1%
11	<a href="http://www.labome.org">www.labome.org</a> Internet Source	1%
12	<a href="http://rsj-prod.literatumonline.com">rsj-prod.literatumonline.com</a> Internet Source	1%
13	<a href="http://www.thieme-connect.com">www.thieme-connect.com</a> Internet Source	1%
14	Qian Zhang, Gabriel A. Al-Ghalith, Mayumi	1%

Kobayashi, Takahiro Segawa, Mitsuto Maeda, Satoshi Okabe, Dan Knights, Satoshi Ishii. "High-Throughput flaA Short Variable Region Sequencing to Assess Campylobacter Diversity in Fecal Samples From Birds", *Frontiers in Microbiology*, 2018

Publication

---

15	<a href="#">zdoc.site</a> Internet Source	1%
16	<a href="#">www.cifa-net.org</a> Internet Source	1%
17	<a href="#">eprints.um.edu.my</a> Internet Source	1%

---

Exclude quotes  On

Exclude matches  < 10 words

Exclude bibliography  On