

FROZEN-THAWED QUALITY OF SWAMP BUFFALO SPERMATOZOA CRYOPRESERVED WITH SUGAR PALM JUICE EXTENDER

by Muhammad Riyadhhi

Submission date: 02-Feb-2021 09:43AM (UTC+0700)

Submission ID: 1499577719

File name: 2020,_Buffalo_Bulletin_September.pdf (452.65K)

Word count: 3374

Character count: 18191

FROZEN-THAWED QUALITY OF SWAMP BUFFALO SPERMATOZOA
CRYOPRESERVED WITH SUGAR PALM JUICE EXTENDERMuhammad Rizal^{1,*}, Muhammad Riyadhhi¹ and Yulnawati Yusnizar²

ABSTRACT

The objective of this research was to examine the effectivity of sugar palm juice as an alternative extender to cryopreserve buffalo. Moreover, we expected to obtain the best composition of sugar palm juice that can be used widely as an alternative extender for buffalo semen cryopreservation. Semen was collected from swamp buffalo bull using artificial vagina. Fresh, diluted, equilibrated, and frozen-thawed was evaluated for general parameters, i.e spermatozoa progressive motility (M), viability (V) and membrane integrity (MI). Fresh semen that had good quality was diluted up to 1.10^8 sperm/ml in four different compositions of extenders, i.e lactose extender containing 20% egg yolk as control (C), 74% sugar palm juice + 20% egg yolk + 6% glycerol (SPG6), 73% sugar palm juice + 20% egg yolk + 7% glycerol (SPG7), and 72% sugar palm juice + 20% egg yolk + 8% glycerol (SPG8), respectively. Diluted-semen was equilibrated at 5°C for 4 h. It followed by frozen and stored in liquid nitrogen (-196°C). The results showed that the percentages of M, V, and MI of frozen-thawed spermatozoa in control (C) were 45.83, 50.37, and 57.83%, respectively. These results were significantly ($P < 0.05$) higher than

those parameters in SPG7 (30.83, 38.83, and 48.83%), SPG6 (18.33, 38.83, and 40.33%), and SPG8 (18.33, 39.17, and 39.33%). In conclusion, sugar palm juice might not be as ideal as control group in maintaining frozen-thawed swamp buffalo semen quality. But this inform us that in a very remote area where chemical or commercial extenders are not easy to find, sugar palm juice can be an alternative due to the fact that it did not affect the pregnancy rate of AI program. Moreover, the cryoprotectant (glycerol) concentration play important role in order to maintain the spermatozoa post thawed quality.

Keywords: *Bubalus bubalis*, buffaloes, sugar palm juice, cryopreservation, artificial insemination, swamp buffalo

INTRODUCTION

Artificial insemination (AI) is most common reproductive technologies that is very useful to increase farm animals population. It is also useful for genetic improvement from the male side (Harshan *et al.*, 2006). AI technology is an integrated process of semen processing

¹Department of Animal Science, Faculty of Agriculture, University of Lambung Mangkurat, Banjarbaru, Indonesia, *E-mail: icang65@yahoo.com

²Research Center for Biotechnology, Indonesian Institute of Sciences, Cibinong, Indonesia

(preservation or cryopreservation) and the artificial insemination itself. In most effective and efficient way, AI has been done using frozen-thawed semen. In the cryopreservation process, ice crystal formation can be harmful for sperm viability. Ice crystal formation occurs during the frozen stage, where the spermatozoa cells are exposed to very low temperature (-196°C) of liquid nitrogen.

Intracellular ice crystallization during cryopreservation is the main cause of damage to the cells. Due to ice crystal formation, normally about 50% of spermatozoa are damaged during cryopreservation (Watson, 2000). Therefore, extender composition and suitable cryoprotectants are important factors for successful semen cryopreservation (Hammerstedt *et al.*, 1990; Curry *et al.*, 1994). The cryoprotectant agent is essential for the viability of spermatozoa after thawing, as these agents minimize intracellular ice formation and restrict the solution effect (Medeiros *et al.*, 2002). Addition of intracellular cryoprotectant compounds, cause increased extender osmotic pressure, and potentially damage spermatozoa cells. Water will leave the cell so that spermatozoa shrink, then swell when cryoprotectant and water reenter the cell to maintain chemical equilibrium. At the time of cryoprotectant removal, the cell initially will swell due to the influx of water and then slowly return to the isoosmotic state when cryoprotectant and water leave the cell. Repeated changes in osmotic pressure can result in significant loss of spermatozoa functional integrity, such as motility, or even cell death without loss of plasma membrane integrity (Gao *et al.*, 1997).

Since the discovery of glycerol (Polge *et al.*, 1949), it has been used extensively for the cryopreservation of many types of cells, including mammalian spermatozoa (McGonagle *et al.*, 2002). However, glycerol has osmotic and toxic

effects on the plasma membrane and metabolism of cryopreserved cells (Hammerstedt *et al.*, 1990). It is responsible for the disorganization of spermatozoa plasma membrane (Hammerstedt and Graham, 1992; Buhr *et al.*, 2001) and reducing motility and fertilizing ability (Jeyendran *et al.*, 1985). Higher concentrations of glycerol lead to cell death (Wünderlich *et al.*, 2006).

MATERIALS AND METHODS

Preparation of extenders

The basic ingredients of lactose extender was 9.3% w/v lactose (Merck, Darmstadt, Germany) and 1.24% w/v fructose (Merck, Darmstadt, Germany) in bidistilled water. Meanwhile, 100 ml pure sugar palm juice was heated up to 100°C and then filtered using a filter paper. Later, those basic ingredients were mixed with egg yolk and glycerol, as it is explained in following table (1). Moreover, 1,000 µg/ml Penicillin and 1,000 IU/ml streptomycin sulphate were added into each group of extenders in order to avoid the contaminations.

Semen collection and processing

Semen from a single buffalo bull was collected at the Artificial Insemination Center of South Kalimantan and processed at the Laboratory of Reproduction and Breeding, Department of Animal Science, University of Lambung Mangkurat (ULM), Banjarbaru, South Kalimantan.

Semen was collected weekly during six weeks in a row. After collection, fresh semen was evaluated immediately for following parameters, including volume, degree of acidity (pH), spermatozoa concentration, mass movement, progressive motility, viability, abnormality, and membrane integrity. Spermatozoa concentration

11s counted using Neubauer chamber. While, progressive motility, viability, abnormality and membrane integrity were observed using 40x objective magnifications microscope.

The qualified fresh semen in this research had >2+ mass movement, $\geq 70\%$ progressive motility, $\geq 5 \times 10^8$ spermatozoa/ml concentration, <10% abnormality. In each replication, the qualified fresh semen was divided in equal volume into four tubes. They have already contained four different extenders in each tube. Final concentration in each tube was 1×10^8 spermatozoa/ml. Later on, diluted semen from each tube was packaged into 0.25 ml plastic straw (IMV France), and equilibrated at 5°C for 4 h.

Freezing process was started by placing the straws 10 cm above surface of liquid nitrogen vapours in styrofoam for 15 minutes, and then straws were plunged into liquid nitrogen (-196°C). Straws were inserted into a goblet and stored in a liquid nitrogen container. After storage for 4 days, random straw samples from each group were thawed at 37°C for 30 seconds to evaluate the sperm quality.

Sperm quality following cryopreservation process

24 Three parameters including progressive motility, viability and membrane integrity were used to evaluate the spermatozoa quality in each group at 17 dilution, equilibration and thawed process. A drop of semen sample was placed on object glass and covered by a thin cover glass to observe the progressive motility. The progressive motility was evaluated randomly in ten different locations under 40x objective magnifications microscope (Rasul *et al.*, 2001).

Spermatozoa viability was evaluated using eosin nigrosin staining (Felipe-Perez *et al.*, 2008).

21 The percentage viability was the number of live cells divided by the total number of the cells in several sites of observation. The dead cells would reserve the staining and the spermatozoa head would become red/dark, while it would not happen in the live cells. At least 200 cells were counted from the different 10 sites of observation under 40x objective magnifications light microscope.

Spermatozoa membrane integrity was 11 evaluated using osmotic resistance test (ORT) with hypo-osmotic swelling (HOS) solution (Revell and Mrode, 1994). About 20 μ l of spermatozoa sample was diluted in 200 μ l of HOS solution, and incubated at 37°C for 45 minutes. Hypo-osmotic swelling solution consists of 0.9 g fructose and 0.49 g natrium citrate in 100 ml bidistilled water. Intact plasma membrane would be swollen on the tail site, while damage plasma membrane would have the linear tail. At least 200 cells were counted from the different 10 sites of observation under 40x objective magnifications light microscope.

Statistical analysis

23 Obtained data of six repetitions were analyzed using analysis of variance (ANOVA) by the linear model using SAS statistical software (SAS 9.1, 2001). The comparative analysis of mean value was analyzed using least significant difference (LSD) test with significant level of 0.05. The results were presented as the means \pm standard error mean (SEM) on the table.

RESULTS AND DISCUSSION

Characteristics of fresh semen

In general, the mean fresh buffalo semen that was used in this research had a good quality. The important parameter such as the percentage of

progressive motility, abnormality, and membrane integrity were 74.17%, 5.17%, and 87.67%, respectively (Table 2). As standard quality said that fresh semen should have the percentage of progressive motility more than 70% and percentage of normal spermatozoa more than 80% (Swelum *et al.*, 2011), and the membrane integrity more than 60% (Revell and Mrode, 1994).

Quality of spermatozoa after semen processing

The results of this research showed that after dilution and equilibration, there was no significant difference ($P>0.05$) among control and treatments on the percentages of progressive motility. Meanwhile after thawing, the percentage of progressive motility of buffalo spermatozoa in control group (45.83%) was significantly ($P<0.05$) higher than SPG6 (18.33%), SPG7 (30.83%), and SPG8 (18.33%) (Table 3). Nevertheless among the treatments group, the percentage of progressive motility, viability, and membrane integrity after thawing of SPG7 was significantly ($P<0.05$) higher than SPG6 and SPG8.

Our data also showed that the frozen-thawed spermatozoa of control and SPG7 were suitable to be used in AI program, with progressive motility more than 30%. Due to Indonesian national standard value (SNI), the minimum percentage of progressive motility for buffalo frozen-thawed semen should be more than 30% (SNI 4869.2-2008).

Equal to that, the percentage of spermatozoa viability after dilution and equilibration was also similar ($P>0.05$) among the groups. While after thawing, the percentage of viability was significantly ($P<0.05$) higher in control (59.37%) group (Table 4). As the previous parameter, the percentage of viability in SPG7 (47.67%) was higher ($P<0.05$) than in SPG6 (38.83%) and SPG8

(39.17%) groups.

In addition, there was no significant difference on the percentage of membrane integrity after dilution and equilibration among the groups. Similar to other two parameters, the percentage of membrane integrity was higher ($P<0.05$) in control (57.83%) than other groups (Table 5). The percentage of membrane integrity is an important variable that strongly related to the motility and fertility of the spermatozoa. Furthermore, the percentage of membrane integrity in SPG7 (48.83%) was higher than SPG6 (40.33%) and SPG8 (39.33%) groups.

Swelum *et al.* (2011) reported that percentage of progressive motility, viability, and membrane integrity of buffalo frozen-thawed spermatozoa in Tris extender containing 7% glycerol were 50.25%, 61.75%, and 68.50%, respectively. El-Sisy *et al.* (2016) reported that the best spermatozoa motility, viability, normal morphology, cell membrane and DNA integrities of buffalo frozen-thawed spermatozoa occurred in Tris extender containing 7.3% glycerol. Abbas and Andrabi (2002) studied the effects of different concentrations of glycerol (2%, 3%, 4%, 5%, 6%, 7%, 8%, 10% or 12%) on post-thawed spermatozoa quality of Nili-Ravi buffalo bull. They reported that the spermatozoa frozen in 7% were significantly better to those in other concentrations of glycerol as judged by post-thawed motility, survivability, and plasma membrane integrity. While Barati *et al.* (2009) used 7% glycerol in cryopreservation of buffalo epididymal spermatozoa. Parallel to those previous results, SPG7 extender in our study that also contains 7% of glycerol showed a better results in three parameters compare to SPG6 and SPG8. Therefore, we assume that the glycerol concentration in the extender is the important key of high success rate of buffalo

Table 1. Extender compositions.

Extender group	Basic lactose (%)	Sugar palm juice (%)	Egg yolk (%)	Glycerol (%)
Control	73	-	20	7
SPG 6	-	74	20	6
SPG 7	-	73	20	7
SPG 8	-	72	20	8

Table 2. The mean characteristics of fresh swamp buffalo semen.

Variable	Mean±SD
Volume (ml)	1.83±0.41
Degree of acidity (pH)	6.90±0.09
Mass movement (1-3)	2.00±0.00
Spermatozoa concentration (million/ml)	1.295±143.07
Progressive motility (%)	74.17±2.04
Live spermatozoa (%)	88.33±1.50
Abnormality (%)	5.17±0.75
Membrane integrity (%)	87.67±0.82

Table 3. Mean percentage of progressive motility after diluting, equilibrating, and thawing.

Treatment	Stage of semen processing		
	After diluting	After equilibrating	After thawing
Lactose	74.17±2.04	68.33±2.58	45.83±4.91 ^a
SPG6	74.17±2.04	65.83±3.76	18.33±2.58 ^c
SPG7	74.17±2.04	68.33±2.58	30.83±2.04 ^b
SPG8	74.17±2.04	65.83±3.76	18.33±2.58 ^c

^{a,b,c} superscript in the same column showed that significantly different (P<0.05).

Table 4. Mean percentage of live spermatozoa after diluting, equilibrating, and thawing.

Treatment	Stage of semen processing		
	After diluting	After equilibrating	After thawing
Lactose	88.33±1.50	82.50±1.05	59.37±3.97 ^a
SPG6	88.33±1.50	80.67±1.63	38.83±2.56 ^c
SPG7	88.33±1.50	80.83±1.83	47.67±3.39 ^b
SPG8	88.33±1.50	80.33±1.86	39.17±1.17 ^c

^{a,b,c} superscript in the same column showed that significantly different (P<0.05).

Table 5. Mean percentage of membrane integrity after diluting, equilibrating, and thawing.

Treatment	Stage of semen processing		
	After diluting	After equilibrating	After thawing
Lactose	87.67±0.82	82.67±1.21	57.83±2.17 ^a
SPG6	87.67±0.82	80.50±1.38	40.33±5.28 ^c
SPG7	87.67±0.82	81.00±1.26	48.83±2.32 ^b
SPG8	87.67±0.82	81.67±1.94	39.33±2.06 ^c

^{a,b,c} superscript in the same column showed that significantly different (P<0.05).

semen cryopreservation. We can see that about 7% of glycerol in the basic extender is the best concentration that can maintain sperm quality during cryopreservation process. Spermatozoa are very sensitive to alterations in the osmolality of the surrounding solution. If the osmolality difference becomes intolerable, spermatozoa lose their motility, and it is an irreversible effect (Willoughby *et al.*, 1996; Gilmore *et al.*, 1998).

In conclusion, sugar palm juice might not as ideal as control group in maintaining frozen-thawed swamp buffalo semen quality. But this inform us that in a very remote area where chemical or commercial extenders are not easy to find, sugar palm juice can be an alternative due to the fact that it did not affect the pregnancy rate of AI program. Moreover, the cryoprotectant (glycerol) concentration play important role in order to maintain the post thawed quality. Further study to evaluate the consisting materials in sugar palm juice is needed in order to obtain stable results in future.

ACKNOWLEDGEMENT

This study was funded by Indonesian Ministry of Research, Technology, and Higher Education of The Republic of Indonesia No. 244/UN8.2/PL/2016. The authors wish to acknowledge the head and staff of Artificial Insemination Center of South Borneo for supporting this study and collecting semen.

REFERENCES

Abbas, A. and S.M.H. Andrabi. 2002. Effect of different glycerol concentrations on motility

before and after freezing, recovery rate, longevity and plasma membrane integrity of Nili-Ravi buffalo bull spermatozoa. *Pak. Vet. J.*, **22**: 1-4. Available on: http://www.pvj.com.pk/pdf-files/22_1/1-4.pdf

Barati, F., M. Khaksary and Gh. Mohammadi. 2009. Cryopreservation of in situ cool stored buffalo (*Bubalus bubalis*) epididymal sperm. *Iran. J. Vet. Res.*, **10**(29): 339-345. DOI: 10.22099/IJVR.2009.1718

Buhr, M.M., P. Fiser, J.L. Bailey and E.F. Curtis. 2001. Cryopreservation in different concentrations of glycerol alters boar sperm and their membranes. *J. Androl.*, **22**: 961-969. DOI: 10.1002/j.1939-4640.2001.tb03436.x

Curry, M.R., J.D. Millar and P.F. Watson. 1994. Calculated optimal cooling rates for ram and human sperm cryopreservation fail to conform with empirical observations. *Biol. Reprod.*, **51**(5): 1014-1021. DOI: 10.1095/biolreprod51.5.1014

El-Sisy, G.A., M.I. Shahba and R.I. El-Sheshtawy. 2016. Freezability of buffalo semen with TRIS extender enriched with disaccharides (trehalose or sucrose) and different glycerol concentrations. *Asian Pacific Journal of Reproduction*, **5**(5): 416-418. DOI: 10.1016/j.apjr.2016.07.007

Felipe-Perez, Y.E., M.L. Juarez-Mosqueda, E.O. Hernandez-Gonzalez and J.J. Valencia. 2008. Viability of fresh and frozen bull sperm compared by two staining techniques. *Acta Veterinaria Brasilica*, **2**(4): 123-130. DOI: 10.21708/AVB.2008.2.4.895

Gao, D.Y., P. Mazur and J.K. Critser. 1997. Fundamental cryobiology of mammalian spermatozoa. In Karow, A.M. and J.K. Critser. (ed.), p. 263-328. *Reproductive*

- Tissue Banking: Scientific Principles*, San Diego: Academic Press, USA.
- Gilmore, J.A., J. Liu, A.T. Peter and J.K. Critser. 1998. Determination of plasma membrane characteristics of boar spermatozoa and their relevance to cryopreservation. *Biol. Reprod.*, **58**(1): 28-56. DOI: 10.1095/biolreprod58.1.28
- Hammerstedt, R.H., J.K. Graham and J.P. Nolan. 1990. Cryopreservation of mammalian sperm: What we ask them to survive. *J. Androl.*, **11**(1): 73-88. Available on: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/j.1939-4640.1990.tb01583.x>
- Hammerstedt, R.H. and J.K. Graham. 1992. Cryopreservation of poultry sperm: The enigma of glycerol. *Cryobiology*, **29**: 26-38. DOI: 10.1016/0011-2240(92)90004-L
- Harshan, H.M., L.P. Singh, A. Arangasamy, M.R. Ansari and S. Kumar. 2006. Effect of buffalo seminal plasma heparin binding protein (HBP) on freezability and *in vitro* fertility of buffalo cauda spermatozoa. *Anim. Reprod. Sci.*, **93**(1-2): 124-133. DOI: 10.1016/j.anireprosci.2005.07.010
- Jeyendran, R.S., H.H. van der Ven, M. Perez-Pelaez and L.J. Zaneveld. 1985. Nonbeneficial effects of glycerol on the oocyte penetrating capacity of cryopreserved and incubated human spermatozoa. *Cryobiology*, **22**(5): 434-437. DOI: 10.1016/0011-2240(85)90154-3
- McGonagle, L.S., M. Goldstein, J. Feldschuh and R.H. Foote. 2002. The influence of cryoprotective media and processing procedures on motility and migration of frozen-thawed human sperm. *Asian J. Androl.*, **4**(2): 137-141.
- Medeiros, C.M., F. Forell, A.T. Oliveira and J.L. Rodrigues. 2002. Current status of sperm cryopreservation: Why isn't it better?. *Theriogenology*, **57**(1): 327-344. DOI: 10.1016/s0093-691x(01)00674-4
- Polge, C., A.U. Smith and A.S. Parkes. 1949. Revival of spermatozoa after vitrification and dehydration at low temperatures. *Nature*, **164**(4172): 666-666. DOI: 10.1038/164666a0
- Rasul, Z., N. Ahmad and M. Anzar. 2001. Changes in motion characteristics, plasma membrane integrity and acrosome morphology during cryopreservation of buffalo spermatozoa. *J. Androl.*, **22**(2): 278-283. DOI: 10.1002/j.1939-4640.2001.tb02181.x
- Revell, S.G. and R.A. Mrode. 1994. An osmotic resistance test for bovine semen. *Anim. Reprod. Sci.*, **36**(1-2): 77-86. DOI: 10.1016/0378-4320(94)90055-8
- SAS, Institute. SAS State Software. 2001. *Changes and Enhancement Through Release 9.1*. SAS Institute, Inc Cary, North Carolina, USA.
- Swellum, A.A., H.A. Mansour, A.A. Elsayed and H.A. Amer. 2011. Comparing ethylene glycol with glycerol for cryopreservation of buffalo bull semen in egg-yolk containing extenders. *Theriogenology*, **76**(5): 833-842. DOI: 10.1016/j.theriogenology.2011.04.015
- Watson, P.F. 2000. The causes of reduced fertility with cryopreserved semen. *Anim. Reprod. Sci.*, **2**(60-61): 481-492. DOI: 10.1016/s0378-4320(00)00099-3
- Willoughby, C.E., P. Mazur, A.T. Peter and J.K. Critser. 1996. Osmotic tolerance limits and properties of murine spermatozoa. *Biol. Reprod.*, **55**(3): 715-727. DOI: 10.1095/biolreprod55.3.715
- Wünderlich, K., U. Paasch, M. Leicht and H.J. Glander. 2006. Activation of caspases in human

spermatozoa during cryopreservation - An immunoblot study. *Cell Tissue Bank.*, 7(2): 81-90. DOI: 10.1007/s10561-005-0276-7

FROZEN-THAWED QUALITY OF SWAMP BUFFALO SPERMATOZOA CRYOPRESERVED WITH SUGAR PALM JUICE EXTENDER

ORIGINALITY REPORT

18%

SIMILARITY INDEX

%

INTERNET SOURCES

18%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Roghieh Sakooei-Vayghan, Seyed Hadi Peighambardoust, Javad Hesari, Maral Soltanzadeh, Donatella Peressini. "Properties of Dried Apricots Pretreated by Ultrasound-Assisted Osmotic Dehydration and Application of Active Coatings", Food Technology and Biotechnology, 2020
Publication 1%
- 2** R.S. JEYENDRAN. "Effect of Glycerol and Cryopreservation on Oocyte Penetration by Human Spermatozoa*", Andrologia, 04/24/2009
Publication 1%
- 3** Erik J. Woods, Sreedhar Thirumala, Sandhya S. Badhe-Buchanan, Dominic Clarke, Aby J. Mathew. "Off the shelf cellular therapeutics: Factors to consider during cryopreservation and storage of human cells for clinical use", Cytotherapy, 2016
Publication 1%

Umut Taşdemir, Serhat Büyükleblebici,

4 Pürhan Barbaros Tuncer, Erdem Coşkun et al. "Effects of various cryoprotectants on bull sperm quality, DNA integrity and oxidative stress parameters", Cryobiology, 2013 1%

Publication

5 M Gunawan, S Setiorini, H N Fitri, E M Kaiin. " The effect of siam orange juice Lour.) in extender on Garut Ram L.) spermatozoa quality post-cryopreservation ", Journal of Physics: Conference Series, 2020 1%

Publication

6 E Nakatsukasa. "Generation of live rat offspring by intrauterine insemination with epididymal spermatozoa cryopreserved at -196 degrees C", Reproduction, 09/01/2001 1%

Publication

7 Dorji, P, V Pattarajinda, and T Vongprolub. "Cryopreservation of Semen of Mithun and Siri Bulls", Bangladesh Journal of Veterinary Medicine, 2014. 1%

Publication

8 Hossein Salmani, Mohammad Mahdi Nabi, Hossein Vaseghi-Dodaran, Mohammad Bozlor Rahman et al. "Effect of glutathione in soybean lecithin-based semen extender on goat semen quality after freeze-thawing", Small Ruminant Research, 2013 1%

Publication

9 Ana J. Soler, Andrés J. García, María R.

Fernández-Santos, Milagros C. Estesó, José J. Garde. "Effects of Thawing Procedure on Postthawed In Vitro Viability and In Vivo Fertility of Red Deer Epididymal Spermatozoa Cryopreserved at -196°C ", Journal of Andrology, 2003

Publication

1%

10

KASHIWAZAKI, Naomi, Yasushi OKUDA, Yasunari SEITA, Shin HISAMATSU, Shigenori SONOKI, Masao SHINO, Toshio MASAOKA, and Tomo INOMATA. "Comparison of Glycerol, Lactamide, Acetamide and Dimethylsulfoxide as Cryoprotectants of Japanese White Rabbit Spermatozoa", Journal of Reproduction and Development, 2006.

Publication

1%

11

Larbi Allai, Xavier Druart, Jesus Contell, Nouredine Louanjli et al. "Effect of argan oil on liquid storage of ram semen in Tris or skim milk based extenders", Animal Reproduction Science, 2015

Publication

1%

12

Buranaamnuay, K.. "Influence of cryoprotectants glycerol and amides, combined with antioxidants on quality of frozen-thawed boar sperm", Animal Reproduction Science, 201108

Publication

1%

13

YULNAWATT, MUHAMMAD GUNAWAN, HERA MAHESHWARI, MUHAMMAD RIZAL, HERDIS, ARIEF BOEDIONO. "Quality of Epididymal and Ejaculated Sperms of Spotted Buffalo in Dextrose Supplemented Extender", HAYATI Journal of Biosciences, 2010

Publication

<1 %

14

CHANAPIWAT, Panida, Kampon KAEOKET, and Padet TUMMARUK. "Cryopreservation of Boar Semen by Egg Yolk-Based Extenders Containing Lactose or Fructose is Better Than Sorbitol", Journal of Veterinary Medical Science, 2012.

Publication

<1 %

15

Shiva Shankar Reddy, N.. "Effects of adding taurine and trehalose to a tris-based egg yolk extender on buffalo (*Bubalus bubalis*) sperm quality following cryopreservation", Animal Reproduction Science, 201006

Publication

<1 %

16

Marie Ide Ngaha Njila, Dieudonné Massoma Lembè, Brice Landry Koloko, Goh Yong Meng et al. " Sperm parameters quality and reproductive effects of methanolic extract of leaves on senescent male rats ", Andrologia, 2019

Publication

<1 %

17

Harshan, H.M.. "Effect of buffalo seminal plasma heparin binding protein (HBP) on

<1 %

freezability and in vitro fertility of buffalo cauda spermatozoa", *Animal Reproduction Science*, 200606

Publication

18

M Gunawan, E M Kaiin, G S Mudita, R R A Chaidir. "Soybean Phospholipids-Based Extender as an Alternative for Bull Sperm Cryopreservation", *IOP Conference Series: Earth and Environmental Science*, 2020

Publication

<1 %

19

Tomás, C., E. Blanch, A. Fazeli, and E. Moc. "Effect of a pre-freezing treatment with cholesterol-loaded cyclodextrins on boar sperm longevity, capacitation dynamics, ability to adhere to porcine oviductal epithelial cells in vitro and DNA fragmentation dynamics", *Reproduction Fertility and Development*, 2012.

Publication

<1 %

20

Joseph Saragusty. "Do physical forces contribute to cryodamage?", *Biotechnology and Bioengineering*, 2009

Publication

<1 %

21

Nabil Mansour. "Effect of extender composition and freezing rate on post-thaw motility and fertility of Arctic char, *Salvelinus alpinus* (L.), spermatozoa", *Aquaculture Research*, 6/2006

Publication

<1 %

22

N. Srivastava, S.K. Srivastava, S.K. Ghosh,

L.P. Singh, J.K. Prasad, Amit Kumar, P. Perumal, A. Jerome, A. Thamizharasan. "Sequestration of PDC-109 protein improves freezability of crossbred bull spermatozoa", *Animal Reproduction Science*, 2012

Publication

<1 %

23

B.M Eriksson, H Rodriguez-Martinez. "Effect of freezing and thawing rates on the post-thaw viability of boar spermatozoa frozen in FlatPacks and Maxi-straws", *Animal Reproduction Science*, 2000

Publication

<1 %

24

B. Khalili, M. Jafaroghli, A. Farshad, M. Paresh-Khiavi. "The Effects of Different Concentrations of Glycine and Cysteine on the Freezability of Moghani Ram Spermatozoa", *Asian-Australasian Journal of Animal Sciences*, 2010

Publication

<1 %

25

Mahdi Zhandi, Mohsen Sharafi. "Negative effect of combined cysteine and glutathione in soy lecithin-based extender on post-thawed ram spermatozoa", *Cell and Tissue Banking*, 2014

Publication

<1 %

26

Sieme, H.. "Cryobiological determinants of frozen semen quality, with special reference to stallion", *Animal Reproduction Science*, 200809

Publication

<1 %

27

S. Movassaghi, G. Saki, F. Javadnia, M. Panahi, M. Mahmoudi, F. Rhim. "Effects of Methyl-Beta-Cyclodextrin and Cholesterol on Cryosurvival of Spermatozoa from C57BL/6 Mouse", Pakistan Journal of Biological Sciences, 2009

Publication

<1%

28

M. García-Herreros. "Morphometric changes in boar spermatozoa induced by cryopreservation", International Journal of Andrology, 7/25/2007

Publication

<1%

29

R Kumar. "Effect of Incorporation of Additives in Tris-Based Egg Yolk Extender on Buffalo (*Bubalus bubalis*) Sperm Tyrosine Phosphorylation During Cryopreservation : Buffalo Sperm Cryopreservation and Tyrosine Phosphorylation", Reproduction in Domestic Animals, 06/2012

Publication

<1%

30

"Abstracts", Reproduction in Domestic Animals, 2014.

Publication

<1%

31

Dawar Hameed Mughal, Ahmad Ijaz, Muhammad Shahbaz Yousaf, Fazal Wadood, Umer Farooq, Syed Aamer Mahmood, Amjad Riaz. " Effect of osmotic pressure on spermatozoa characteristics of cryopreserved buffalo bull () semen ", Journal of Applied

<1%

32

Santiago-Moreno, J., C. Castano, A. Toledano-Diaz, M. A. Coloma, A. Lopez-Sebastian, M. T. Prieto, and J. L. Campo. "Semen cryopreservation for the creation of a Spanish poultry breeds cryobank: Optimization of freezing rate and equilibration time", Poultry Science, 2011.

Publication

<1%

33

Haugan, T.. "Effects of sperm concentration at semen collection and storage period of frozen semen on dairy cow conception", Animal Reproduction Science, 200701

Publication

<1%

34

Villaverde, Ana Izabel S. Balbin, Eduardo G. Fioratti, Marcimara Penitenti, Maura R.V. Ikoma, Miriam H. Tsunemi, Frederico O. Papa, and Maria D. Lopes. "Cryoprotective effect of different glycerol concentrations on domestic cat spermatozoa", Theriogenology, 2013.

Publication

<1%

Exclude quotes

Off

Exclude matches

< 2 words

Exclude bibliography

On