

Fish species composition and diversity in a river, a swamp, and a reservoir in Banjar District, South Kalimantan Province

by Rusmilyansar Rusmilyansar


Submission date: 01-Apr-2021 03:26PM (UTC+0700)

Submission ID: 1548064049

File name: 2021.412-423.pdf (577.42K)

Word count: 6237

Character count: 32718



Fish species composition and diversity in a river, a swamp, and a reservoir in Banjar District, South Kalimantan Province

¹Rusmilyansari, ¹Aulia A. Wahab, ²Wiryo, ¹Reika Cahyati

¹Study Program of Fisheries Resource Utilization, Faculty of Fisheries and Marine, Lambung Mangkurat University, South Kalimantan, Indonesia; ² Graduate program in Natural Resource Management, Faculty of Agriculture, Universitas Bengkulu, Bengkulu, Indonesia. Corresponding author: Rusmilyansari, r_melyan@yahoo.com

Abstract. Freshwater fish is a component of biodiversity in river, swamp, and lake ecosystems. Over-exploitation of fish can change the characteristics of waters and result in the decrease in the abundance and diversity of fish. This study aimed to assess the species composition and diversity of fish community in three mainland water systems, i.e., river, swamp and reservoir. Purposive sampling of fish was done in three stations for each type of water. The data were analyzed descriptively and quantitatively, using the Shannon-Wiener index of diversity, evenness index, Simpson index of dominance (C) and similarity index of Sørensen (ISs) and Bray-Curtis (C_{bc}). A total of 35 species belonging to 18 families were found in the three water types, where the river had the highest species richness (17 species), followed by the swamp (15) and the reservoir (3). The family Cyprinidae had the highest number of species. The index value of fish species diversity ranged from 1.20 to 2.55, considered as medium, the evenness index 0.5-0.9, considered medium to high, and the dominance index 0.09-0.39 considered as low. The Sørensen similarity index ranged from 0.12 to 0.29, and the Bray-Curtis similarity index 0.05-0.38, considered as low. The river and swamp fish community had higher similarity index than the reservoir and the river, and the reservoir and the swamp. No common species was found in all the three water types. The water physico-chemical parameters met the quality standards to support fish life, namely pH 6.0-7.88, temperature 26.4-30.1°C, DO 6.2-8.1 mg L⁻¹, transparency 97-130 cm.

Key Words: freshwater fish, habitat types, species diversity.

Introduction. Fish constitutes a large portion of the biodiversity of aquatic ecosystems, and consequently, play significant roles in maintaining the aquatic ecosystem services of which benefit the human communities (Arthington et al 2016). Approximately 43% or 11,952 species of fish occupy freshwater ecosystems (Nelson 2006), such as rivers, lakes, and swamps (Kottelat et al 1993). These freshwater fish contribute significantly to the livelihood of the people, especially in the developing world (Sayer et al 2018). However, the water where fish live is needed by many sectors, such as agriculture, energy, industries and domestic purposes (Taylor et al 2016), which lead to the competition for water uses among sectors, threatening the freshwater ecosystems which cause the decline of freshwater fish diversity (Dudgeon 2019).

Pollution from industrial, agricultural and domestic uses have detrimental effects on both, marine and freshwater fish (Ganguly 2013). Freshwater swamps are rich in fish diversity in Southeast Asia have been threatened due to conversion to other landuses (Ho et al 2018). River systems are also severely threatened ecosystems on the earth due to several activities, one of which is the construction of dam which alters the natural flow of rivers (Schneider et al 2017). Since many species require migration to complete their life cycles, the construction of dams impede the migration of fish (Leirmann et al 2012). In Yangtze River basin of China, spawning sites of some migration fish has been lost due to the construction of a dam, resulting in the decline of growth and reproduction (Yi et al 2010). Another study in the upstream from the upper Parana River floodplain, Brazil,

showed that the construction of a dam has simplified the the functional diversity of fish (Oliveira et al 2018).

The threats to freshwater ecosystems has reduced the diversity of freshwater fish, nearly a third of which are in danger of extinction (Dudgeon 2012), and many species have been extinct already (Baillie & Butcher 2012). With the loss of freshwater biodiversity, their essential support to human well-being will be lost too (Albert et al 2020). It is, therefore, urgent to protect the freshwater ecosystems to ensure the sustainability of freshwater fish consumption for the welfare of human communities. The first step to do is to improve the assessment of the production in order to have science-based fishery management, because the accurate and complete information of freshwater fish is not sufficient at local, national and global level (Taylor et al 2016).

In Kalimantan there are about 350 species of freshwater fish, 135 of which are endemic to this island. One of the districts in South Kalimantan Province which has large production of freshwater fish is Banjar District, with an area of +466,850 ha, producing 4,053.3 tonnes of freshwater fish in 2018 (Office of fisheries Banjar District 2020). The freshwater ecosystem types in Banjar District are swamps, rivers and dams or reservoirs (Office of fisheries Banjar District 2020). The Ir. P. M. Noor Riam Kanan Dam (hereafter referred to as Riam Kanan Dam), located in Banjar District, is a multi-use dam with an area of 9,730 ha, used for hydroelectric power, agricultural irrigation, raw water for local drinking water company, tourism and fisheries.

The construction of Riam Kanan Dam certainly has impacts on fish diversity like what happened in other places (Yi et al 2010; Leirmann et al 2012; Schneider et al 2017; Oliveira et al 2018). The impact of Riam Kanan Dam construction on fish diversity in the river has not been studied. Meanwhile, the dam itself has been used by the regional government for the development of fishery. The first step to be taken in making management plan of the Riam Kanan Reservoir fishery and other waters is to gather information of fish diversity and abundance, in order to avoid over exploitation that will deplete the fish resource.

The objective of this study was to assess the species composition and diversity of fish in three mainland water systems, i.e., Martapura river, Lebak swamp and Riam Kanan reservoir. We hypothesized that the species composition among the three water systems would be different due to differences in the habitat's characteristics including the water quality.

Material and Method

Study site. The research was conducted two months from April to June 2020 in Riam Kanan reservoir, Martapura river and Lebak swamp in Banjar District, South Kalimantan Province. The study sites were selected purposively by considering the characteristics of the population based following Usman & Purnomo (2008). Sampling points were selected systematically with a random starting point at 9 points in different water systems (Figure 1) based on information from fishermen who usually catch at that locations. In the river, the government has introduced non native fish species, i.e., *Oreochromis niloticus* and *Cyprinus carpio*, while in the reservoir, *C. carpio*, *Colossoma macropomum*, and *Pangasianodon hypophthalmus*.




Figure 1. The sites of study at a Lebak swamp, Martapura river and Riam Kanan reservoir in Banjar District, South Kalimantan Province, Indonesia.

Data collection. Fish samples were taken using fishing gill net, trammel net, portable lift net, hand line, stage trap and cast net, fish pots and scoop net, hand line, bamboo's split 34th one day trip catching time. The fish caught were collected by separating each species of catch and counting the number of each species of fish. Identification of fish species was carried out by observing their morphometric and meristic characteristics by referring to the fish identification key book (Saainin 1984). Sampling at each sample point was carried out 3 times with a time interval between sampling of 1 week to obtain an accurate sample. Each fish caught was identified and counted. The number of sampling was considered sufficient because at the third sampling, no more new species was found in each water system (Table 1).

Table 1

Total species found in each sampling period according water system

Sampling period	Total species		
	River	Swamp	Reservoir
First sampling	14	10	16
Second sampling	17	11	16
Third sampling	17	11	16

The water quality parameters, namely brightness, temperature and pH were measured in situ as supporting data. In addition, water samples were taken for dissolved oxyeg (DO) measurement in the laboratory.

Data analysis. The data were analyzed using the diversity Shannon-Wiener index (H'), the evenness index (E), the dominance index (C), and Sørensen similarity index (Krebs 1972) and Bray-Curtis similarity index (Chao et al 2006).

Shannon-Wiener diversity index was calculated using the following equation:

$$H' = - \sum_{i=1}^s (p_i) (\log p_i)$$

Where:

H' - species diversity index

s - number of species

p_i - proportion of species i = n_i/N = (number of individual i/number of individuals of all species).

Evennes index was calculated using formula:

$$E = \frac{H'}{\ln(s)}$$

Where:

E - evennes index

H' - species diversity index

s - number of species

Simpson dominance index was calculated with formula:

$$C = \sum_{i=1}^s (p_i)^2$$

Where:

C - Simpson dominance index

p_i - proportion of species i = n_i/N

Similarity index of species composition was determined using Sørensen index and Bray-Curtis index, with the following formulas:

$$\text{Sørensen index} = IS_s = \frac{2c}{A + B}$$

Where:

A - number of all fish species in community A

B - number of all fish species in community B

c - number of common species (present in both communities A and B)

(Ludwig & Reynolds 1988)

$$\hat{C}_{BC} = \frac{2 \sum_{i=1}^{D_{12}} \min(X_i, Y_i)}{\sum_{i=1}^{D_1} X_i + \sum_{i=1}^{D_2} Y_i}$$

CBC - Bray-Curtis similarity index

Min(X_i, Y_i) - the minimum count of the ith species in the first community (X) and second community (Y).

D₁ - species found in the first community (X)

D₂ - species found in the second community (Y)

D₁₂ - common species found in the first and second community

(Chao et al 2006)

Results and Discussion

Species composition. The fish community in the sampling sites in Martapura river was composed of 17 species from 13 families, in Lebak swamp 11 species from 8 families, and in Riam Kanan reservoir 15 species from 6 families, with a total of 35 species from 18 families in all the three water systems (Table 2). The species composition differed among the water systems. The Sørensen similarity index of species composition among the water systems ranged from 0.12 to 0.29, while the Bray-Curtis index 0.05 to 0.38 (Table 2). No single common species was found in all the three water systems, but there were four common species found the river and swamp waters, two common species

between the river and reservoir, and two common between the swamp and the reservoir (Table 2). The low similarity index of species composition indicates that the environmental factors or human disturbances among the three water systems were different.

Table 2
The list of species and family of fish caught in Martapura river, Lebak swamp and Riam Kanan reservoir in Banjar District, South Kalimantan Province

Species	Family	Number of individuals			Total
		River	Swamp	Reservoir	
<i>Trichopodus trichopterus</i>	Anabantidae	265	740		1005
<i>Trichopodus pectoralis</i>	Anabantidae	216	500		716
<i>Helostoma temminckii</i>	Helostomatodae	220	380		600
<i>Anabas testudineus</i>	Anabantidae	141	325		566
<i>Oxyeleotris marmorata</i>	Eleotridae			396	396
<i>Channa striata</i>	Channidae		175	81	256
<i>Barbonymus schwanenfeldii</i>	Cyprinidae			241	241
<i>Osteochilus vittatus</i>	Cyprinidae	26		84	184
<i>Mystacoleucus padangensis</i>	Cyprinidae		170		175
<i>Mystus nigriceps</i>	Bagridae		148	27	175
<i>Rasbora argyrotaenia</i>	Cyprinidae			168	168
<i>Clarias batrachus</i>	Clariidae		168		168
<i>Pangasianodon hypophthalmus</i>	Pangasidae	166			166
<i>Pangasius nasutus</i>	Pangasidae	152			152
<i>Oreochromis niloticus</i>	Cichidae	108			108
<i>Osphronemus goramy</i>	Osphronemidae	46		45	91
<i>Kryptopterus macrocephalus</i>	Siluridae	89			89
<i>Hampala macrolepidota</i>	Cyprinidae			87	87
<i>Channa micropeltes</i>	Channidae			82	82
<i>Cyprinus carpio</i>	Cyprinidae	74			74
<i>Belontia hasselti</i>	Osphronemidae		61		61
<i>Hemisilurus heterorhynchus</i>	Siluridae	56			56
<i>Osteochilus vittatus</i>	Cyprinidae		49		49
<i>Mystus gulio</i>	Bagridae	46			46
<i>Macrobrachium rosenbergii</i>	Palaemonidae	45			45
<i>Barbonymus gonionotus</i>	Cyprinidae	25			25
<i>Monopterus albus</i>	Synbranchidae		21		21
<i>Pterygoplichthys</i> sp.	Loricaridae	15			15
<i>Oxygaster anomalura</i>	Cyprinidae			14	13
<i>Chitala lopis</i>	Notopteridae	8			8
<i>Cychoheilichthys apogon</i>	Cyprinidae			7	7
<i>Mastacembelus erythrotaenia</i>	Mastacembelidae			7	7
<i>Hemibagrus nemurus</i>	Bagridae			6	6
<i>Mystacoleucus marginatus</i>	Cyprinidae			5	5
<i>Osteochilus repang</i>	Cyprinidae			3	3
35 Total individuals		1698	2737	1253	5698
Number of families		13	8	6	
Number of species		17	11	15	

Family Cyprinidae constituted the largest portion of the fish community by having the highest number of species, i.e., twelve, followed by family Anabantidae, three species. Each of the following families, i.e., Channidae, Bagridae, Osphronemidae and Siluridae was represented by two species, while each of other families had only one species. The four most abundant species were *Trichopodus trichopterus*, *Trichopodus pectoralis*, *Helostoma temminckii* and *Anabas testudineus* (Table 3).

Table 3
Common species and similarity index of species composition among water systems

Species	Family	Water systems			Similarity index	
		River	Swamp	Reservoir	Sørensen	Bray-Curtis
<i>Trichopodus trichopterus</i>	Anabantidae	265	740		River and swamp	
<i>Trichopodus pectoralis</i>	Anabantidae	216	500			
<i>Helostoma temminckii</i>	Helastomatodae	220	380		0.29	0.38
<i>Anabas testudineus</i>	Anabantidae	141	325			
<i>Osphronemus goramy</i>	Osphronemidae	46		45	River and reservoir	
<i>Osteochilus vittatus</i>	Cyprinidae	26		84	0.12	0.05
<i>Channa striata</i>	Channidae		175	81	Swamp and reservoir	
<i>Mystus nigriceps</i>	Bagridae		148	27	0.15	0.05

Species diversity. In addition to number of species or species richness, diversity index, which integrate the species richness and abundance, has been used by ecologists to determine the species diversity. In the river of Martapura the Shannon-Wiener index of species diversity was 2.57, in Lebak swamp 2.09 and in Riam Kanan reservoir 2.16, which could be considered medium. The evenness index was high in all three water systems, i.e., 0.91 in Martapura river, 0.84 in Lebak swamp and 0.78 in Riam Kanan reservoir, indicating that the abundance of each species was relatively equal. This was evident from the low value of Simpson dominance index, i.e., 0.09 in Martapura river, 0.15 in Lebak swamp and 0.09 in Riam Kanan Reservoir.

In Martapura river, all the community indexes had the same value among the sampling sites, while in Lebak swamp and Riam Kanan the values varied among sampling sites (Table 4).

Table 4
Indexes of diversity, evenness and dominance of fish community in the nine sampling sites in three water systems

Community indexes	Habitat type / Sampling site									
	9	River			Swamp			Reservoir		
	Site 1	Site 2	Site 3	Site 1	Site 2	Site 3	Site 1	Site 2	Site 3	
14 Species diversity (H')	2.55	2.54	2.55	2.23	1.98	1.86	1.69	1.43	1.20	
Evenness index (E)	0.90	0.89	0.90	0.89	0.80	0.75	0.77	0.56	0.5	
Dominance index (C)	0.09	0.09	0.09	0.12	0.18	0.22	0.15	0.38	0.39	
Species diversity (H')		2.57			2.09			2.16		
Evenness index (E)		0.91			0.84			0.78		
Dominance index (C)		0.09			0.15			0.09		

Water quality. The parameters of water quality are shown in Table 5. The pH of water in the study sites ranged from 6.0 to 7.88, the lowest of which was found in Lebak swamp and the highest in Martapura river. The lowest temperature was also found in the swamp, i.e., 26.4°C while the highest in the reservoir, i.e., 30.1°C. The reservoir also had the highest dissolved oxygen, i.e., 8.1 mg L⁻¹, while the lowest in the river, i.e., 6.2 mg L⁻¹. Meanwhile, the highest clarity was found in the river, and the lowest in the swamp. The salinity was zero in all water systems. Overall, the water quality in all the tree habitats

was within the normal range for fish habitat, except that the water clarity was too high, because its optimum value for fish growth is 30-80 cm (Bhatnagar & Devi 2013).

Table 5
The water quality in Martapura river, Lebak swamp and Riam Kanan reservoir

Water quality	River			Swamp			Reservoir		
pH	7.88	7.78	7.16	6.39	6.17	6.0	7	6.9	6.8
Temperature (°C)	28.5	28.4	28.7	26.4	26.3	27.1	30.0	30.1	29.9
Salinity (‰)	0	0	0	0	0	0	0	0	0
Dissolved oxygen (mg L ⁻¹)	6.8	7.2	6.2	7.45	7.65	7.6	8.1	7.8	7.5
Clarity (cm)	130	124	130	98	97	99	102	102.4	102.2

Species composition. The species composition in the river, swamp and dam in this study was different from each other as indicated by low similarity indexes, both the Sørensen and Bray-Curtis indexes. The Bray-Curtis Index is more accurate because it is based not only on the presence or absence of common species but also on the abundance of those species, while the Sørensen index was based only on the absence and presence of species (Cao et al 2006). In this study, Bray-Curtis index showed the pronounced difference in species composition of fish community between the dam and the two other water types, i.e., 0.05. Meanwhile, the similarity of species composition between the fish community in the river and in the swamp was higher, i.e., 0.38 because they had not only higher number of common species but also the higher abundance of those common species. The four common species between the river and swamp also became the most abundant species in the study sites. i.e., *Tr. trichopterus*, *T.s pectoralis*, *H. temminckii* and *A. testudineus*. Two of these four species, i.e., *T. trichopterus* and *T. pectoralis* were also found as the most dominant species in Bankau swamp of Hulu Sungai Selatan District, South Kalimantan Province (Irhamsyah et al 2017). On the contrary, these four species were not found in the Riam Kanan dam. The most abundant species found in the dam was *Oxyeleotris marmorata* (Table 2).

Differences in species composition among the water systems in this study may reflect the differences in the habitat's characteristics. In Colorado river basin of the United States, the fish community composition was associated with differences in water temperature, concentrations of dissolved solids and suspended sediment, elevation, habitat, and land use which in turn was the effects of natural and human factors (Deacon & Mize 1997). In this study sites, the water of dam had higher temperature (29.9-30.1°C) than the water in the river (28.3-28.7°C) and swamp (26.3-27.1°C) (Table 5). The optimum temperature for fish is 20-30°C (Effendi 2003), so the temperature in the dam was on the higher end of range of the optimum. The depth was also different among the three water systems, i.e., Riam Kanan dam 52 m, Martapura river 4-8 m, and Lebak swamp 1-4 m. Another difference was the water clarity, with the river having the highest clarity while the swamp the lowest.

In a stable small stream, the fish species found are those which have evolved various morphological and behavioral adaptations to exploit that habitat, while over geographical and habitat scales, the species found are determined by environmental filtering and anthropogenic effects (Gebrekios 2016). In our study sites, the anthropogenic factor might have affected the species composition. In the river and swamp, fishermen sometime used poison and electric current to catch the fish, while in dam these activities were absent.

In this study the Cyprinidae was the most often found family, similar with the finding of Rashid et al (2015) in Tembeling and Pahang rivers, Pahang, Malaysia, and that of Samitra & Rozi (2019) in Lakitan river, South Sumatera. In Brunei Darussalam, among 104 freshwater species found, 40 of which belong to famili Cyprinidae (Sulaiman et al 2018). These results are in accordance with the statement of Lowe-McConnell (1987) that freshwater fish in tropic Asia are dominated by the family Cyprinidae. The family Cyprinidae can live in both river areas that have strong currents and weak currents with good water quality (Nikolsky 1963). Buwono et al (2017) stated that the Cyprinidae is a

family with a relatively large number of species in freshwater. Cyprinidae is a very large family and occurs almost everywhere except in Australia, Madagascar, New Zealand, and South America; to date, 2420 species in 220 genera in the fish family Cyprinidae have been found (Karahana & Ergenen 2010). The very large number of species in the family Cyprinidae shows good adaptability and fast breeding processes, so species from this family are always found in freshwater almost all parts of the world (Beamis et al 2006). Cyprinidae dominate 20% of the world's freshwater fish and 8% of all fish (Berra 1997).

Three of the four most abundant species found both Martapura river and Lebak swamp, i.e., *A. testudineus*, *T. pectoralis*, *T. trichopterus* belong to the family Anabantidae which have additional respiration in the form of grooved bony bones like coral waste called a labyrinth which is used to take oxygen directly from the air (Muflikhah 2007). *Channa striata* which is ranked 5th in abundance also has labyrinth. Meanwhile, the most abundant species in Riam Kanan dam was *O. marmorata*, the most expensive fish in Banjar district.

Three of the introduced species, i.e., *P. hypophthalmus*, *O. niloticus* and *C. carpio*, were found in Martapura river, but not in Riam Kanan dam although those species were also introduced in the dam. The introduction of exotic species was intended to enrich freshwater fish diversity, because according to Fisheries Office of Banjar District, several freshwater fish had begun to disappear, namely *Osteochilus melanopleurus*, which is the mascot of Banjarmasin City, *Eleutheronema tetradactylum*, *Channa pleurophthalma*, *Channa lucius*, *Leptobarbus hoevenii*, *Barbonymus schwanenfeldii* and *Wallago attu*. Despite the good intention of the government, introduction of alien species may result in the demise of native species of fish, and even jeopardize the ecosystem service of the 28th habitat (Kiruba-Sankar et al 2018). Introduced species may also reduce the functional diversity of the invaded fish community (Milardi et al 2019). In Martapura river, *P. hypophthalmus*, *O. niloticus* and *C. carpio* ranked 4th, 7th and 9th in abundance respectively. Thus, the detrimental effect of exotic species in the fish community in dam was not evident in Martapura river. In Riam Kanan dam the introduced species did not even survived.

Species diversity and fish abundance. The Martapura river had 27th species richness and diversity index than Lebak swamp and Riam Kanan reservoir. The number of fish species in Martapura river (17) was lower than that in Lakitan river (20) in S 25th Sumatra (Samitra & Rozi 2019), in Trembeling river (47) also in Malaysia (Rashid et al 2015). The high number of species found in Trembeling river (47 species), however, was the result of two years of sampling (Rashid et al 2015), while the sampling in our study was done only in three weeks. Within two years of sampling there were several seasons 29th and each season might have different species composition as reported in the study in Ayeerwady River in Sagaing Region, Upper Myanmar (Win & Myint 2015), so the total species diversity of two year-sampling would be higher 23th than that of three week-sampling.

Fish density in a river is affected by several factors, such as substrate type, stream type and disturbance (Ahmad et al 2011). Martapura river has slow current, sandy and stony bottom and experiences disturbance from human activities such as the passing of boats and the fishing activities, sometimes using the destructive methods, i.e., electrofishing and poisoning.

Lebak swamp had the lowest number of species, i.e., 11, lower than Nee Soon freshwater swamp forest of Singapore, i.e., 21 species (Ho et al 2018), Shadegan marshland in Iran, i.e., 26 species (Hashemi et al 2015), and Beriah Kanan River (21 species), but higher than that in Ulu Sedili (10 species) in Malaysia (Zakaria et al 1999). Both Beria Kanan and Ulu Sedili rivers are mostly freshwater sw 30th mp. The study in peat swamp forest even found much higher number of species, i.e., 55 species in Sebangau watershed, Central Kalimantan (Thornton et al 2018), and 73 in species Tripa peat swamp forest in Aceh (Muchlisin et al 2015). This comparison, however, does not give the comprehensive information because of differences in the areas and duration of sampling.

Although Lebak swamp had the lowest diversity of fish, it had the highest fish abundance, presumably due to the presence of aquatic plants such as parupuk

(*Phragmites karka*), *Hydrilla (Hydrilla verticillata)*, *mendong (Fimbristylis globulosa)*, common susum (*Hanguana malayana*), common water hyacinth (*Eichhornia crassipes*), water snowflake (*Nymphoides indica*) and spiny false fiddleleaf (*Hydrolea spinosa*). Grown et al (2003) stated that aquatic plants in the swamp support fish stocks. According to Welcomme (1985) the lowland swamp fish community is influenced by three key factors, namely the inundation cycle, the pressure of fishing activities and a decrease in environmental quality.

Water quality. Fish has tolerable range of the water quality in which they live. Physical, chemical and biological factors determine the water quality which in turn influence the survival and growth of fish, and consequently the fish production (Bhatnagar & Devi 2013). The quality of water in three water systems in this study was within the range of tolerance for the survival of fish, except the water clarity which was too high. The pH in this study sites was 6.0–7.88, with the lowest one being found in the swamp due to the presence of abundant aquatic plants. Also, a freshwater swamp may contain several inches of peat layer (Whitten et al 1984). However, the pH in the swamp was still within normal range for fish, i.e., 6–9, and certain fish may even survive in more acidic water (Effendi 2003). However, in acidic water the release of heavy metals from waste into water is high (Li et al 2013) which may have detrimental effect on fish. The temperature in the study sites was 26.4–30.1°C, while the optimum temperature for fish according to Effendi (2003) is 20–30°C. Some fish may tolerate higher temperature, i.e., 30–35°C (Bhatnagar & Devi 2013), but at higher temperature (30–35°C) the release of heavy metals from waste into water is high (Li et al 2013). The dissolved oxygen in the study site was 6.2–8.1 mg L⁻¹, while the optimum DO for fish is >5 mg L⁻¹ (Bhatnagar & Devi 2013). The water clarity in the study sites was too high because the optimum range for fish health is 30–80 cm (Bhatnagar & Devi 2013). At higher clarity light penetrates deeper encouraging the growth of macrophytes, reducing the growth of plankton as fish food.

Conclusions. In conclusion, this study has shown that Martapura river, Lebak swamp and Riam Kanan reservoir had clear distinction in species composition. The fish species diversity in the reservoir was lower than that of the other water systems, but the diversity between the river and swamp did not differ significantly. However, since the time period of sampling was only three weeks and the number locations of sampling was only three for each water system, many parts of each habitat has not been adequately sampled. Therefore, further studies covering wider area and longer sampling period should be conducted to get comprehensive data of fish community as the basis for the fishery management.

Acknowledgement. We express our gratitude to University of Lambung Mngkurat which has given us research grant through the LPPM (Institution for Research and Community Service).

References

- Ahmad A. K., Mohd Sham O., Shukor M. N., Aweng Eh-R., Shuhaimi-Othman O., 2011 Ecological factors affecting fish diversity and density in Sungkai Wildlife Reserve, Perak, Malaysia. *The Zoologist* 9:78–84.
- Albert J. S., Destouni G., Duke-Sylvester S. M., Magurran A. E., Oberdorff T., Reis R. E., Winemiller K. O., Ripple W. J., 2020 Scientists' warning to humanity on the freshwater biodiversity crisis *Ambio* doi:10.1007/s13280-020-01318-8
- Arthington A. H., Dulvy N. K., Gladstone W., Winfield I. J., 2016 Fish conservation in freshwater and marine realms: Status, threats and management. *Aquatic Conserv: Marine and Freshwater Ecosystems* 26(5):838–857.
- Baillie J. E. M., Butcher E. R., 2012 Priceless or worthless? The world's most threatened species. *Zoological Society of London, United Kingdom*, 123 p.

- Bhatnagar A., Devi P., 2013 Water quality guidelines for the management of pond fish culture. *International Journal of Environmental Sciences* 3(6):1980-2009.
- Beamish F. W. H., Sa-Ardrit P., Tongnunui S., 2006 Habitat characteristics of the Cyprinidae in small rivers in central Thailand. *Environmental Biology of Fishes*, 76(2-4), 237. DOI: 10.1007/S10641-006-9029-0
- Berra T. M., 1997 Some 20th century fish discoveries. *Environmental Biology of Fishes* 50:1-12.
- Buwono N. R., Fariedah F., Anestyningrum R. E., 2017 Komunitas Ikan di Sungai Jerowan Kabupaten Madiun. *Journal of Aquaculture and Fish Health* 6(2):81-88.
- Chao A., Chazdon R. L., Colwell R. K., Shen T. J., 2006 Abundance-based similarity indices and their estimation when there are unseen species in samples. *Biometrics* 62(2):361-371.
- Deacon J. R., Mize C. V., 1997 Effects of water quality and habitat on composition of fish communities in the upper Colorado River Basin. *Upper Colorado River Basin NAWQA Fact Sheet FS-122-97*
- Dudgeon D., 2019 Multiple threats imperil freshwater biodiversity in the Anthropocene. *Current Biology* 29(19):960-967.
- Dudgeon D., 2012 Threats to freshwater biodiversity globally and in the Indo-Burma Biodiversity Hotspot In: *The status and distribution of freshwater biodiversity in Indo-Burma*. Allen D. J., Smith K. G., Darwall W. R. T. (eds), pp 1-28, IUCN, Cambridge, UK and Gland, Switzerland.
- Effendi H., 2003 Telaah kualitas air bagi pengelolaan sumberdaya dan lingkungan perairan. Kanisius, Yogyakarta, 145 p.
- Gebrekiros S. T., 2016 Factors affecting stream fish community composition and habitat suitability. *Journal of Aquaculture and Marine Biology* 4(2):00076. DOI: 10.15406/jamb.2016.04.00076
- Ganguly S., 2013 Water pollution from various sources and human infringements: an editorial. *Indian Journal of Scientific Research and Technology* 1(1):54-55.
- Growns I., Gerke P. C., Astles K. L., Pollard D. A., 2003 A comparison of fish assemblages associated with different riparian vegetation types in the Hawkesbury-Nepean River System. *Fisheries Management & Ecology* 10:209-220.
- Hashemi S. A., Ghorbani R., Kymaram F., Hossini S. A., Eskandari G., Hedayati A., 2015 Fish species composition, distribution and abundance in Shadegan Wetland. *Fisheries and Aquaculture Journal* 6: 128. doi:10.4172/2150-3508.1000128
- Ho J. K. I., Ramchunder S. J., Memory A., Tan H. H., Yeo D. C. J., 2018 Native and introduced fish community structure in a freshwater swamp forest: Implications for conservation and management. *Aquatic Conserv: Marine and Freshwater Ecosystems* 29(1):47-58.
- Irhamyah, Ahmadi, Rusmilyansari, 2017 Fish and fishing gears of the Bangkai Swamp, Indonesia. *Journal of Fisheries* 5:489-496.
- Karahan A., Ergenen S., 2010 Cytogenetic analysis of *Garra variabilis* (Heckel, 1843) (Pisces, Cyprinidae) from Savur Stream (Mardin), Turkey. *Turkish Journal of Fisheries and Aquatic Sciences* 10(4):483-489.
- Kiruba-Sankar R., Raj P. J., Saravanan K., Kumar L., Angel R., Murugan A. V., Roy S. D., 2018 Invasive species in freshwater ecosystems. In: *Threats to ecosystem services. Biodiversity and climate change adaptation in tropical islands*. Sivaperuman C., Velmurugan A., Singh A., Jaisankar I. (eds), Academic Press. London.
- Kottelat M., Whitten A. J., Kartikasari S. N., Wirjoatmojo S., 1993 *Freshwater fishes of Western Indonesia and Sulawesi*. Periplus Edition-EMDI Project, Jakarta.
- Krebs C. J., 1972 *Ecology: The experimental analysis of distribution and abundance*. Harper and Row Publishing, New York.
- Li H., Shi A., Li M., Zhang X., 2013 Effect of pH, temperature, dissolved oxygen, and flow rate of overlying water on heavy metals release from storm sewer sediments. *Journal of Chemistry*, Article ID 434012, <https://doi.org/10.1155/2013/434012>
- Liermann C. R., Nilsson C. J., Ng R. Y., 2012 Implications of dam obstruction for global freshwater fish diversity. *BioScience* 62:539-548.

- Lowe-McConnell R. H., 1987 Ecological studies in tropical fish communities Melbourne: Cambridge University Press, London, pp. 159-173.
- Ludwig J. A., Reynolds J. F., 1988 Statistical ecology. John Wiley and Sons, New York.
- Milardi M., Gavioli A., Soininen J., Castaldelli G., 2019 Exotic species invasions undermine regional functional diversity of freshwater fish. *Scientific Reports* 9:1–10.
- Muchlisin Z. A., Akyun Q., Rizka S., Fadli N., Sugiantp S., Halim A., Siti-AZizah M. N., 2015 Ichthyofauna of Tripa Peat Swamp Forest, Aceh province, Indonesia. *Check List* 11(2): 1560, doi: <http://dx.doi.org/10.15560/11.2.1560>
- Muflikhah N., 2007 Domestikasi Ikan Gabus (*Channa striata*). Prosiding Seminar Nasional Tahunan IV Hasil Penelitian Perikanan dan Kelautan. Jurusan Perikanan dan Kelautan Universitas Gadjah Mada, pp. 1-10.
- Nelson J. S., 2006 Fishes of the world. 4rd edition, Jhon Wiley & Sons, New York.
- Nikolsky G. V., 1963 The ecology of fishes. Translated from Russian by L. Birkett. Academic Press, London, 352 p.
- Oliveira A. G., Dias R. M., Baumgartner M. T., Gomes L. C., Agostinho A. A., 2018 Long-term effects of flow regulation by dams simplify fish functional diversity. *Freshwater Biology* 63(3):293–305.
- Rashid Z. A., Asmuni M., Amal M. N., 2015 Fish diversity of Tembeling and Pahang Rivers, Pahang Malaysia. *Checklist* 11(5): 1753, DOI: <https://doi.org/10.15560/11.5.1753>
- Saanin H., 1984 Taksonomi dan Kunci Identifikasi Ikan (Jilid 1 dan 2). Binacipta: Bogor.
- Samitra D., Rozi Z. F., 2019 The fish fauna in Lakitan River, Musi Rawas Regency, South Sumatra. *Jurnal Biodjati* 4(1):11-20.
- Sayer C. A., Máiz-Tomé L., Darwall W. R. T., 2018 The importance of freshwater species to livelihoods in the Lake Victoria Basin. In: *Freshwater biodiversity in the Lake Victoria Basin: guidance for species conservation, site protection, climate resilience and sustainable livelihoods*. IUCN UK, IUCN Gland, Switzerland, Gland, pp. 136–151.
- Schneider C., Flörke M., De Stefano L., Petersen-Perlman J. D., 2017 Hydrological threats to riparian wetlands of international importance – a global quantitative and qualitative analysis. *Hydrology and Earth System Sciences* 21:2799–2815.
- Sulaiman Z., Hu T. H., Lim K. K. P., 2018 Annotated checklist of freshwater fishes from Brunei Darussalam, Borneo. *Zootaxa* 4379(1):24–46.
- Taylor W. W., Bartley D. M., Goddard C. I., Leonard N. J., Welcomme R. (eds), 2016 *Freshwater, fish and the future: proceedings of the global cross-sectoral conference*. Food and Agriculture Organization of the United Nations, Rome; Michigan State University, East Lansing; American Fisheries Society, Bethesda, Maryland.
- Thornton S. A., Dudin, Page S. E., Upton C., Harrison M. E., 2018 Peatland fish of Sebangau, Borneo: diversity, monitoring and conservation. *Mires and Peat*, Volume 22, DOI: 10.19189/MaP.2017.OMB.313
- Usman H., Purnomo S. A., 2008 Metode penelitian sosial. Bumi angkasa, Jakarta, 110 p.
- Welcomme R. L., 1985 River Fisheries. FAO Fish Tech. Pap, (262), Rome.
- Whitten A. J., Damanik S. J., Anwar J., Hisyam N., 1984 The ecology of Sumatra. Gadjah Mada University Press, Yogyakarta, Indonesia.
- Win M. L., Myint T. T., 2015 Species composition, seasonal occurrence and abundance of freshwater fishes in Ayeyarwady River Segment, Sagaing Region, Upper Myanmar. *IOP Conf. Series: Earth and Environmental Science* 416 (2020) 012015. doi:10.1088/1755-1315/416/1/012015
- Yi Y., Yang Z., Zhang S., 2010 Ecological influence of dam construction and river-lake connectivity on migration fish habitat in the Yangtze River basin, China. *Procedia Environmental Sciences* 2:1942–1954.
- Zakaria R., Mansor M., Ali A. B., 1999 Swamp-riverine tropical fish population: a comparative study of two spatially isolated freshwater ecosystems in Peninsular Malaysia. *Wetlands Ecology and Management* 6:261–268.
- *** Office of fisheries of Banjar District, 2020 Annual statistics of fiseheries in public water. (Dinas Perikanan dan Kelautan. 2020. Laporan Tahunan Statistik Perikanan Di Perairan Umum.) Kabupaten Banjar. Martapura.

Received: 18 January 2021. Accepted: 19 February 2021. Published online: 26 February 2021.

Authors:

Rusmilyansari, Lambung Mangkurat University, Faculty of Fisheries and Marine, Study Program of Fisheries Resource Utilization, Indonesia, South Kalimantan, Simpang Empat Banjarbaru 70714, Jl. Achmad Yani, e-mail: r_melyan@yahoo.com

Aulia Azhar Wahab, Lambung Mangkurat University, Faculty of Fisheries and Marine, Study Program of Fisheries Resource Utilization, Indonesia, South Kalimantan, Simpang Empat Banjarbaru 70714, Jl. Achmad Yani, e-mail: auliaazharwahab@ulm.ac.id

Wiryono, Bengkulu University, Faculty of Agriculture, Study program in Natural Resource Management, Indonesia, Bengkulu, Kec. Muara Bangka Hulu 38371, Kandang Limun, Jl. WR. Supratman, e-mail: wiryonogood@yahoo.com

Reika Cahyati, Lambung Mangkurat University, Faculty of Fisheries and Marine, Study Program of Fisheries Resource Utilization, Indonesia, South Kalimantan, Simpang Empat Banjarbaru 70714, Jl. Achmad Yani, e-mail: r2kacahyati@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Rusmilyansari, Wahab A. A., Wiryono, Cahyati R., 2021 Fish species composition and diversity in a river, a swamp, and a reservoir in Banjar District, South Kalimantan Province. AACL Bioflux 14(1):412-423.

Fish species composition and diversity in a river, a swamp, and a reservoir in Banjar District, South Kalimantan Province

ORIGINALITY REPORT

13%

SIMILARITY INDEX

11%

INTERNET SOURCES

9%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Bethune Cookman University Student Paper	2%
2	www.bioflux.com.ro Internet Source	2%
3	biodiversitas.mipa.uns.ac.id Internet Source	1%
4	eia.emb.gov.ph Internet Source	1%
5	Pathy Kibungu kembelo, Favien Nzuki Bakwaye, Honoré Belesi Katula, Wouter Vanhove, Patrick Van Damme. "Ethnobotanical characterization of medicinal plants used in Kisantu and Mbanza-Ngungu territories, Kongo-Central Province in DR Congo", Research Square, 2020 Publication	1%
6	pubs.usgs.gov Internet Source	<1%
7	link.springer.com	

<1%

8

F Rahmayanti, F K Nazira, A K Dewi, D F Oktaviani et al. "Biodiversity of gastropod in the Sombu Beach, Wakatobi, Indonesia", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1%

9

laptop.deh.gov.au

Internet Source

<1%

10

onlinelibrary.wiley.com

Internet Source

<1%

11

medcraveonline.com

Internet Source

<1%

12

"Freshwater Fisheries Ecology", Wiley, 2015

Publication

<1%

13

www.fao.org

Internet Source

<1%

14

Desrita, I S Tamba, A Muhtadi, J Ariyanti, R Leidonald. " Diversity and habitat condition of Tor Fish (spp.) in the upstream of Wampu Waters, North Sumatra, Indonesia ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1%

15

W Subchan, V E Susilo, Y Khairiyah, G Wahyudewantoro, S Ariyunita, A Rohman. "The diversity of freshwater fish in sanenrejo and wonoasri river resorts from meru betiri national park", Journal of Physics: Conference Series, 2021

Publication

<1%

16

Rajeev Raghavan, Siby Philip, Neelesh Dahanukar, Anvar Ali. "Freshwater biodiversity of India: a response to Sarkar et al. (2013)", Reviews in Fish Biology and Fisheries, 2013

Publication

<1%

17

ijcrbp.com

Internet Source

<1%

18

issuu.com

Internet Source

<1%

19

Andrew P. Goyke, Anne E. Hershey. "Effects of fish predation on larval chironomid (Diptera: Chironomidae) communities in an arctic ecosystem", Hydrobiologia, 1992

Publication

<1%

20

S Irhami, A Fithri, A S Batubara, Z A Muchlisin. "Fish fauna of Meureubo River, Aceh Barat District, Indonesia", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1%

21	jurnal-iktiologi.org Internet Source	<1%
22	idoc.pub Internet Source	<1%
23	www.ajol.info Internet Source	<1%
24	"Fishes and Forestry", Wiley, 2004 Publication	<1%
25	"Life on Land", Springer Science and Business Media LLC, 2021 Publication	<1%
26	Bowman, William D.. "Ecology", Oxford University Press Publication	<1%
27	M Dekar, M A Sarong, A S Batubara, Z A Muchlisin. "Ichthyofauna of Aceh River, Aceh Province, Indonesia", IOP Conference Series: Earth and Environmental Science, 2018 Publication	<1%
28	Matteo Gentilucci, Costantino Parisi, Maria Rosaria Coppola, Fatima-Zahra Majdoubi, Adriano Madonna, Giulia Guerriero. "Influence of Mediterranean Sea Temperature Increase on Gaeta Gulf (Tyrrhenian Sea) Biodiversity", Proceedings of the Zoological Society, 2020 Publication	<1%

29 May Lei Win, Than Than Myint. "Species Composition, Seasonal Occurrence and Abundance of Freshwater Fishes in Ayeyarwady River Segment, Sagaing Region, Upper Myanmar", IOP Conference Series: Earth and Environmental Science, 2020
Publication <1%

30 Sara A. Thornton, Erna Setiana, Kris Yoyo, Dudin, Yulintine, Mark E. Harrison, Susan E. Page, Caroline Upton. "Towards biocultural approaches to peatland conservation: The case for fish and livelihoods in Indonesia", Environmental Science & Policy, 2020
Publication <1%

31 biotaxa.org
Internet Source <1%

32 journalarticle.ukm.my
Internet Source <1%

33 jurnal.untidar.ac.id
Internet Source <1%

34 oro.open.ac.uk
Internet Source <1%

35 wwfke.awsassets.panda.org
Internet Source <1%

36 www.hindawi.com
Internet Source <1%

37

www.ukm.my

Internet Source

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On