



The fertility of South Kalimantan buffalo spermatozoa after cold preservation and cryopreservation

Muhammad Rizal^{1*}, Muhammad Riyadhi¹, Nursyam Andi Syarifuddin¹, Muhammad Thahir²

¹*Department of Animal Science, Faculty of Agriculture, Lambung Mangkurat University, Jl. Jenderal Ahmad Yani Km. 36 Banjarbaru, Indonesia*

²*Agriculture Service of Enrekang District Government, Jl. Poros Pinang–Rappang Km. 3 Enrekang, Indonesia*

Key words: Sugar palm juice, Preservation, Cryopreservation, Artificial insemination, South Kalimantan buffalo.

<http://dx.doi.org/10.12692/ijb/17.5.85-95>

Article published on November 12, 2020

Abstract

Sugar palm juice can be an alternative extender of buffalo semen because it contains nutrients needed by spermatozoa during preservation. The objective of this study was to examine the fertility of South Kalimantan buffalo spermatozoa after the preservation and cryopreservation process. Semen was collected by the artificial vagina. Fresh semen was divided into two tubes in equal volume and then diluted with sugar palm juice extender with a different composition. Semen in the first tube was diluted with 80% sugar palm juice + 20% egg yolk, and then stored in a refrigerator at 5°C (chilled-semen) for three days. Semen in the second tube was diluted with 73% sugar palm juice + 20% egg yolk + 7% glycerol. Diluted-semen was loaded in a mini straw and then stored in a liquid nitrogen container (-196°C) for seven days. A total of 22 adult female buffaloes aged 3–5 years were selected as acceptors and estrous synchronization by PGF_{2α}. Thirteen females were inseminated with chilled-semen, and nine females were inseminated with frozen semen. The results of this study showed that motility of spermatozoa after preservation for three days was an average of 45.83% and 38.33% for frozen-thawed semen. The percentage of pregnancy and birth for chilled-semen was 69.23% and 61.54%, higher than for frozen-thawed semen (66.67% and 55.55%), but there was no significant difference between treatment. It can be concluded that sugar palm juice is suitable as an alternative extender for South Kalimantan buffalo semen because it is able to maintain the quality and fertility of spermatozoa after preservation and cryopreservation.

* **Corresponding Author:** Muhammad Rizal ✉ mrizal@ulm.ac.id

Introduction

Swamp buffalo (*Bubalus bubalis*) in South Kalimantan Province through the Minister of Agriculture's decree in 2012 has been designated as a separate family and named the South Kalimantan buffalo. The decree also emphasized that the South Kalimantan buffalo is a wealth of genetic resources of Indonesian local livestock that must be protected and preserved. One effective form of conserving buffaloes in South Kalimantan is to increase the population, so as to improve the welfare of buffalo breeders.

The application of artificial insemination (AI) technology is one alternative method to accelerate population increase and improve buffalo genetic quality (Morrell, 2006). Preservation and cryopreservation of semen are an integral and inseparable part of AI technology (Sansone *et al.*, 2000), and it is an essential method for conservation of germ cells and is suitable for the proper genetic management (Singh *et al.*, 2013). Semen cryopreservation is an effective method in the preservation of superior male genetic resources directly related to AI technology so that its application will enhance the genetic quality improvement of livestock (Martinez *et al.*, 2007). According to Hafez and Hafez (2000), the main purpose of processing semen is to increase the capacity of semen to serve more female. To achieve this goal, semen is diluted with certain extenders, which meet conditions such as energy source, buffer, non-toxic, prevent damage to spermatozoa, do not reduce spermatozoa fertility, do not impede the evaluation of spermatozoa after dilution, inexpensive, and easily obtained.

Extracts of natural ingredients from vegetables and fruits and their seeds contain an important compounds for health, so it is able to maintain life (Aviram *et al.*, 2000; Aviram *et al.*, 2004). These natural extracts and infusions are used in semen extenders for preserving animal sperm (Sansone *et al.*, 2000). Some natural ingredients commonly used as semen extender contains strong antioxidant compounds, thus protecting spermatozoa from

oxidative damage during the cryopreservation process (Seeram *et al.*, 2005; Tezcan *et al.*, 2009). Utilization of natural materials such as coconut water in semen cryopreservation has been reported in buck (Daramola *et al.*, 2016) and cattle (El-Sheshtawy, 2017; Tarig *et al.*, 2017), with good results.

Sugar palm juice can be an alternative extender of semen because it contains nutrients needed by spermatozoa during preservation (Rizal and Riyadhi, 2016), but do not have protective compounds of spermatozoa during the process of semen preservation at 5°C and cryopreservation at -196°C. In the process of preservation and cryopreservation of semen, extender must be added a special compound in the form of cryoprotectant which functions to protect spermatozoa from damage (Andrabi *et al.*, 2008). Egg yolk is one component that is commonly used in the preservation and cryopreservation of semen. Egg yolk lipoprotein, cholesterol, and lecithin are the function to maintain the integrity of the plasma membrane lipoprotein of spermatozoa during preservation at low temperatures (Kumar *et al.*, 1992). Specifically, in the cryopreservation process of semen, other than egg yolk, extender should also be added to intracellular cryoprotectant compounds such as glycerol which functions to minimize cell damage during freezing (Leboeuf *et al.*, 2000). Through this treatment, it is expected that the quality of spermatozoa during preservation and cryopreservation of semen can be maintained, and does not reduce fertility when applied in AI technology. The objective of this study was to examine the fertility of South Kalimantan buffalo spermatozoa after the preservation and cryopreservation process.

Materials and methods

Collection, evaluation, preservation, and cryopreservation of semen

Semen was collected from two adult South Kalimantan buffaloes using artificial vagina at The Center for Artificial Insemination, South Kalimantan Province, Banjarbaru. Semen was collected 12 times as a replication. Quality of fresh semen was evaluated,

including volume, consistency, pH, mass movement, spermatozoa concentration, spermatozoa motility, and spermatozoa abnormality. Fresh semen that meets the quality requirements was diluted according to treatment. The good fresh semen must have spermatozoa motility > 70% (Barati *et al.*, 2009; Akhter *et al.*, 2011; Wadood *et al.*, 2016; Almadaly *et al.*, 2019), mass movements > ++, spermatozoa concentration > 1,000 million cells/mL, and spermatozoa abnormality < 15% (Almadaly *et al.*, 2019). Preservation, cryopreservation, and evaluation of spermatozoa quality were carried out at The Animal Production Laboratory, Department of Animal Science, Lambung Mangkurat University, Banjarbaru.

Fresh semen was divided into two tubes in equal volume and then diluted with sugar palm juice extender with a different composition. Semen in the first tube was diluted with 80% sugar palm juice + 20% egg yolk, and then stored in a refrigerator at 5°C (chilled-semen) for three days. Chilled-semen was diluted to a concentration of 10 million motile per mini straw (0.25 mL). The extender was added with an antibiotic, penicillin 1,000 µg/mL and streptomycin 1,000 IU/mL extender. Sugar palm juice was prepared by heating to boiling and then filtered with filter paper.

Semen in the second tube was diluted with 73% sugar palm juice + 20% egg yolk + 7% glycerol. Diluted-semen was loaded in a mini straw (0.25 mL), and then equilibrated in a refrigerator at 5°C for four hours. Semen was frozen by placing straw 10 cm above the surface of liquid nitrogen in closed styrofoam (temperature around -130°C) for 15 minutes. Straw was put into liquid nitrogen (temperature around -196°C) and then stored in a liquid nitrogen container for seven days. Frozen semen was diluted to a concentration of 30 million motile spermatozoa per mini straw.

Estrous synchronization, artificial insemination, and pregnancy evaluation

A total of 22 adult female buffaloes aged 3–5 years

were selected as acceptors in the Maju Jaya 1 Farmers Group in Benua Raya Village, Tanah Laut District. Acceptors were estrous synchronization by injecting 2 ml (0.5 mg cloprostenol sodium) prostaglandin hormone (PGF_{2α}, BoVet). Each acceptor was injected with PGF_{2α} twice with an injection period of 12 days. The symptom of estrous was observed 2 days after the second injection of PGF_{2α}.

Artificial insemination was carried out on females who show symptoms of estrous 3 days (72 hours) after the second injection of PGF_{2α}. Thirteen females were inseminated with chilled-semen, and nine females were inseminated with frozen semen. Artificial insemination was using rectovaginal methods. Semen was deposited in the cervical lumen and the uterine corpus. Each buffalo was inseminated with one straw. Motility of spermatozoa was evaluated before using an AI application. Thawing of frozen semen was by putting a straw into 37°C water for 30 seconds.

Pregnancy was evaluated by observation of estrus 20–22 days (one estrous cycle) and 41–43 days (two estrous cycles) after insemination. Females who showed no symptoms of estrus were diagnosed pregnant, and others were not pregnant. The percentage of birth was obtained after the buffaloes give birth.

Statistical analysis

Data about the percentage of pregnancy and birth were analyzed using the Chi-squared test.

Results and discussion

Characteristics of fresh semen and semen quality after preservation and cryopreservation

The results obtained that fresh semen characteristics of South Kalimantan buffalo were volume 1.97 mL, concentration of spermatozoa 1,066.5 million/mL, and spermatozoa motility 71.67% (Table 1). In previous studies, it was reported that the volume of swamp buffalo semen was 2.02 mL (Amin *et al.*, 1999), 0.225 mL (Yulnawati *et al.*, 2010), and 1.44–1.76 mL (Sianturi *et al.*, 2012).

Table 1. Characteristics of South Kalimantan buffalo fresh semen.

Semen attributes	Mean \pm SD
Volume (mL)	1.97 \pm 0.35
Consistency	Moderate
Degree of acidity (pH)	6.9 \pm 0.42
Mass movement (1 – 3)	2 \pm 0.14
Concentration of spermatozoa (million/mL)	1,066.5 \pm 98.23
Spermatozoa motility (%)	71.67 \pm 3.27
Spermatozoa abnormality (%)	8.67 \pm 0.49

The consistency and mass movement of swamp buffalo semen were thick and 2–3 (Amin *et al.*, 1999), thin to thick and 2–3 (Sianturi *et al.*, 2012), respectively. Spermatozoa concentrations of swamp buffalo was 1,447.14 million/mL (Amin *et al.*, 1999), 2,695 million/mL (Yulnawati *et al.*, 2010), 1,070–1,983 million/mL (Sianturi *et al.*, 2012), and 1,600 million/mL (Almadaly *et al.*, 2019). The percentage of spermatozoa motility of swamp buffalo was 76.43%

(Rizal *et al.*, 1999), 70% (Yulnawati *et al.*, 2010), and 78.7% (Almadaly *et al.*, 2019). The percentage of spermatozoa abnormality of swamp buffalo was 12.14% (Rizal *et al.*, 1999) and 78.7% (Almadaly *et al.*, 2019). According to Mahmoud *et al.* (2013) that semen volume, spermatozoa concentration, and spermatozoa motility of buffalo in Iran were 2.9 mL, 1,079.2 million/mL, and 65.8%, respectively.

Table 2. Percentage of pregnancy and birth of South Kalimantan buffalo.

Treatment	Pregnancy (%)	Birth (%)
Chilled-semen (5°C)	69.23 (9/13)	61.54 (8/13)
Frozen-thawed semen (-196°C)	66.67 (6/9)	55.55 (5/9)

The results of this study showed that there was a decrease in spermatozoa motility with increasing storage time of semen at 5°C. The motility of spermatozoa after preservation for three days was an average of 45.83%. Decreased spermatozoa motility caused by the decreasing quality of the extender which becomes more acidic. The acidic condition of semen extender is caused by a buildup of metabolic lactic acid. The end result of the metabolic process of spermatozoa that preserved in the refrigerator is lactic acid because the storage takes place anaerobically (without oxygen). According to Stryer (1995), the end result of cell metabolism under anaerobic conditions is lactic acid.

The main principle of preservation of semen associated with efforts to extend the life of spermatozoa is to reduce the degree of metabolism through storage at low temperatures (Lemma, 2011). Preservation of semen at 5°C causes changes in the

spermatozoa cell membrane from liquid to gel. Metabolism of the cell is maximum at body temperature, and optimal when stored at room temperature (24–29°C). McKinnon (1999) reported that every drop in temperature of 10°C, resulting from a decrease in metabolism up to 50% so that preservation at 5°C metabolism of spermatozoa only about 10%.

Spermatozoa motility after thawing was obtained an average of 38.33%. The results showed that a combination of 20% egg yolk and 7% glycerol in sugar palm juice extender is able to maintain the spermatozoa motility of South Kalimantan buffalo during the cryopreservation process. Egg yolk contains lecithin and cholesterol which functions to protect the cell plasma membrane due to cold shock during freezing (Hermansson and Linde-Forsberg, 2006). While, glycerol is functions to protect the cell by minimizing the formation of ice crystal in the cell

(Mazur, 1984), preventing cell dehydration, maintain the balance of electrolytes osmotic pressure inside and outside of the cell (Watson, 2000), and maintain the stability of protein and glycoprotein of the cell

plasma membrane (Park and Graham, 1992). Almadaly *et al.* (2019) reported the percentage of buffalo spermatozoa motility after thawing cryopreserved with 7% glycerol is 41.3%.



Fig. 1. Pedets results from AI chilled-semen.

El-Sisy *et al.* (2016) reported that the glycerol concentration was 7% better compared to 3 and 5% in the cryopreservation of buffalo semen. According to Abbas and Andrabi (2002), the cryopreservation process of Nili-Ravi buffalo semen that glycerol concentration was 7% better compared to 5, 6, 8, and 10%. The quality of frozen semen of Saanen goats containing 7% glycerol is better compared to 5 and 9% (Kulaksiz *et al.*, 2013). The opposite occurs at a concentration of 8%, which is thought to be too much so that it increases the osmotic pressure of the extender, and is bad for spermatozoa. Willoughby *et al.* (1996) reported that cells will experience death if they are in an osmotic pressure of medium that exceeds the tolerance limit. According to Mughal *et al.* (2018), the osmotic pressure of the solution used in extender preparation plays an important role in the post-thaw quality of cryopreserved buffalo bull semen.

Estrous response and percentages of pregnancy and birth from AI

The result of this study was showed that all (100%)

acceptors showing estrous symptoms 2–3 days after the second injection of PGF 2α . This shows that all buffaloes have good reproductive conditions because the occurs cyclically and regularly, typically in buffaloes around 20–22 days. According to Brito *et al.* (2002), responses to the administration of PGF 2α to cattle will be effective if the estrous cycle is regular and there is a corpus luteum (CL) in the luteal phase (about 17 days from the estrous cycle for 20–22 days). The results also confirm that the estrous synchronization method applied can initiate various physiological mechanisms such as folliculogenesis, ovulation, spermatozoa and oocyte transport, fertilization, and the synthesis and secretion of hormones directly involved in regulating various reproductive activities. Sianturi *et al.* (2012) reported that female swamp buffaloes injected with the PGF 2α or a combination of the PGF 2α and gonadotropin-releasing hormone (GnRH) as well as a combination of the PGF 2α and human chorionic gonadotropin (hCG) produce a percentage of estrous is 100%, and estrous symptoms began to appear on two days after the second injection of the PGF 2α .

The percentage of pregnancy for chilled-semen was 69.23%, higher than for frozen-thawed semen (66.67%). The percentage of birth for chilled-semen was 61.54%, while for frozen-thawed semen was 55.55%. There was no significant difference between

treatment on the percentage of pregnancy and birth (Table 2). There was an embryo death in each treatment. According to (Hunter, 1981), the mortality of embryo in the early stage of pregnancy in livestock can reach 30%.



Fig. 2. Pedets results from AI using frozen-thawed semen.

The pregnancy rate of chilled-semen was higher than frozen-thawed semen. This is presumably because the quality of spermatozoa in chilled-semen was better than that of spermatozoa in frozen-thawed semen, so it also affects the fertility of spermatozoa. Generally, 40–50% of sperm cells die after cryopreservation even with an optimized methodology (Su *et al.*, 2007). According to Singh *et al.* (2012), freezing-thawing semen procedure is results in more damage to the spermatozoa of bull semen than chilling semen, so that leads to reduced conception rate. Watson (2000) also reported that one of the factors that can reduce spermatozoa fertility is the process of semen cryopreservation. According to Ansari *et al.* (2012), buffalo spermatozoa are more vulnerable to freezing-thawing damage compared with other mammalian species.

The results of this research explain that sugar palm juice was feasible as an alternative extender for semen of the South Kalimantan buffalo because it was able to maintain the fertility of spermatozoa after the cold

preservation (Fig. 1) and cryopreservation (Fig. 2). This is confirmed that the combination of sugar palm juice with egg yolk and glycerol can minimize spermatozoa damage during cryopreservation. According to Barile (2005), the success of artificial insemination is influenced by several factors, i.e., the main thing is the reproductive condition of females, the quality of inseminated semen, the accuracy of estrous detection, and the correct skills and handling of semen.

The percentage of pregnancy obtained in this study was confirmed by the results of several previous researchs. Sianturi *et al.* (2012) reported that the percentage of pregnancy of swamp buffalo in Banten Indonesia was 66.7–75%, which inseminated with frozen-thawed semen. The pregnancy rate of buffaloes which inseminated with frozen-thawed semen was 56.5% (Berber *et al.*, 2002), 40–60% Barile (2005), 34.6–46.2% (Swelum *et al.*, 2011) and 52% (Almadaly *et al.*, 2019) of buffalo in Egypt, 44.45% of buffalo in Iran (Mahmoud *et al.*, 2013),

40% in Bangladeshi buffaloes (Hamid, 2018), and 68.5% of swamp buffalo in Ngawi Indonesia (Budiarto *et al.*, 2019).

Yulnawati *et al.* (2013) reported the pregnancy rate of swamp buffalo inseminated with frozen-thawed epididymal spermatozoa of spotted buffalo was 37.5–40%. In water buffalo, the pregnancy rate is reported as 36% in Italian Mediterranean buffaloes (Neglia *et al.*, 2003), 33% in Murrah buffaloes (Paul and Prakash, 2005), 36% (Warriach *et al.*, 2008) and 37% (Naseer *et al.*, 2011) in Nili-Ravi buffaloes, and 45–62.7% (Gutiérrez-Añez *et al.*, 2018). Based on the results of the study it can be concluded that fresh semen of South Kalimantan buffalo has a good quantity and quality, and is eligible to be processed into chilled-semen or frozen semen for AI purposes. Sugar palm juice is suitable as an alternative extender for South Kalimantan buffalo semen because it is able to maintain the quality and fertility of spermatozoa after preservation and cryopreservation.

Acknowledgements

The authors thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, the Head and Staff of the Artificial Insemination Center of South Kalimantan Province, the Head of Animal Production Laboratory, Department of Animal Science, Lambung Mangkurat University, and the Maju Jaya 1 Farmers Group in Benua Raya Village, Tanah Laut District for funding support, animal, and laboratory facilities.

References

Abbas A, Andrabhi SMH. 2002. Effect of different glycerol concentrations on motility before and after freezing, recovery rate, longevity and plasma membrane integrity of Nili-Ravi buffalo bull spermatozoa. *Pakistan Veterinary Journal* **22**, 1-4. http://pvj.com.pk/abstract/22_1/1.htm

Akhter S, Rakha BA, Ansari MS, Andrabhi SMH, Ullah N. 2011. Storage of Nili-Ravi buffalo (*Bubalus bubalis*) semen in skim milk extender supplemented with ascorbic acid and α -tocopherol.

Pakistan Journal of Zoology **43**, 273-277.

<https://docplayer.net/137400288-Storage-of-nili-ravi-buffalo-bubalus-bubalis-semen-in-skim-milk-extender-supplemented-with-ascorbic-acid-and-alpha-tocopherol.html>

Almadaly EA, Tawfik FS, El-Kon II, Heleil BA, Fattouh EM. 2019. Effect of different cryoprotectants on the post-thaw sperm characteristics and in vivo fertility of buffalo (*Bubalus bubalis*) bull semen. *Slovak Veterinary Research* **56** (22), 541-551.

<https://www.cabdirect.org/cabdirect/abstract/20193447175>

Amin MR, Toelihere MR, Yusuf TL, Situmorang P. 1999. Pengaruh plasma semen sapi terhadap kualitas semen beku kerbau lumpur (*Bubalus bubalis*). *Jurnal Ilmu Ternak dan Veteriner* **4**, 143-147.

<http://dx.doi.org/10.14334/jitv.v4i3.151>

Andrabhi SMH, Ansari MS, Ullah N, Anwar M, Mehmood A, Akhter S. 2008. Duck egg yolk in extender improves the freezability of buffalo bull spermatozoa. *Animal Reproduction Science* **104**, 427-433.

<https://doi.org/10.1016/j.anireprosci.2007.07.003>

Ansari MS, Rakha BA, Andrabhi SMH, Ullah N, Iqbal R, Holt WV, Akhter S. 2012. Glutathione supplemented tris-citric acid extender improves the postthaw quality and *in vivo* fertility of buffalo (*Bubalus bubalis*) bull spermatozoa. *Reproduction Biology* **12**, 271-6.

<https://doi.org/10.1016/j.repbio.2010.10.001>

Aviram M, Dornfield L, Rosenblatt M, Volkova N, Kaplan M, Coleman R. 2000. Pomegranate juice consumption reduces oxidative stress, atherogenic modifications to LDL, and platelet aggregation: studies in humans and in atherosclerotic apolipoprotein E-deficient mice. *American Journal of Clinical Nutrition* **71**, 1062-1076.

<http://dx.doi.org/10.1093/ajcn/71.5.1062>

- Aviram M, Rosenblatt M, Gaitani D, Nitecki S, Hoffman A, Dornfield L.** 2004. Pomegranate juice consumption for 3 years by patients with carotid artery stenosis (CAS) reduces common carotid intima-media thickness (IMT), blood pressure and LDL oxidation. *Clinical Nutrition* **23**, 423-433.
<http://dx.doi.org/10.1016/j.clnu.2003.10.002>
- Barati F, Mahabady KM, Mohammadi GH.** 2009. Cryopreservation of in situ cool stored buffalo (*Bubalus bubalis*) epididymal sperm. *Iranian Journal of Veterinary Research* **10**, 339-345.
http://ijvr.shirazu.ac.ir/article_1718_354f16050fe0fb2226c57858cb3b6c76.pdf
- Barile VL.** 2005. Improving reproductive efficiency in female buffaloes. *Livestock Production Science* **92**, 183-194.
<https://doi.org/10.1016/j.livprodsci.2004.06.014>
- Berber RCA, Madureira EH, Baruselli PS.** 2002. Comparison of two ovsynch protocols (GnRH vs. LH) for fixed timed insemination in buffalo (*Bubalus bubalis*). *Theriogenology* **57**, 1421-1430.
[https://doi.org/10.1016/S0093-691X\(02\)00639-8](https://doi.org/10.1016/S0093-691X(02)00639-8)
- Brito LFC, Satrapa R, Marson EP, Kastelic JP.** 2002. Efficacy of PGF₂α to synchronize estrus in water buffalo cows (*Bubalus bubalis*) is dependent upon plasma progesterone concentration, corpus luteum size and ovarian follicular status before treatment. *Animal Reproduction Science* **73**, 23-35.
[https://doi.org/10.1016/S0378-4320\(02\)00124-0](https://doi.org/10.1016/S0378-4320(02)00124-0)
- Budiarto A, Ciptadi G, Hakim L, Putri AI.** 2019. Reproductive performance and fertility index of swamp buffalo (*Bubalus bubalis*) in ngawi regency, east java. *IOP Conf. Series: Journal of Physics: Conf. Series* 1146 (2019) 012024. Hlm 1-4.
<https://doi.org/10.1088/1742-6596/1146/1/012024>
- Daramola JO, Adekunle EO, Oke OE, Onagbesan OM, Oyewusi IK, Oyewusi JA.** 2016. Effects of coconut (*Cocos nucifera*) water with or without egg-yolk on viability of cryopreserved buck spermatozoa. *Animal Reproduction* **13**, 57-62.
<http://dx.doi.org/10.21451/1984-3143-AR724>
- El-Sheshtawy RI, El-Nattat WS, Ali GAD.** 2017. Cryopreservation of cattle semen using coconut water extender with different glycerol concentrations. *Asian Pacific Journal of Reproduction* **6**, 279-282.
<http://dx.doi.org/10.4103/2305-0500.217343>
- El-Sisy GA, Shahba MI, El-Sheshtawy RI.** 2016. Freezability of buffalo semen with TRIS extender enriched with disaccharides (trehalose or sucrose) and different glycerol concentrations. *Asian Pacific Journal of Reproduction* **5**, 416-418.
<https://doi.org/10.1016/j.apjr.2016.07.007>
- Gutiérrez-Añez JC, Palomares RA, Jiménez-Pineda JR, Camacho AR, Portillo-Martínez GE.** 2018. Pregnancy rate in water buffalo following fixed-time artificial insemination using new or used intravaginal devices with two progesterone concentrations. *Tropical Animal Health Production* **50**, 629-634.
<https://doi.org/10.1007/s11250-017-1479-1>
- Hafez ESE, Hafez B.** 2000. *Reproduction in Farm Animals* 7th Edition. Lippincott Williams & Wilkins, Baltimore.
- Hamid MA.** 2018. Study on the effect of insemination time on pregnancy rate of Bangladeshi buffalo in intensive farming. *SAARC Journal of Agriculture* **16**, 143-152.
<https://doi.org/10.3329/sja.v16i2.40266>
- Hermansson U, Linde-Forsberg C.** 2006. Freezing of stored, chilled dog spermatozoa. *Theriogenology* **65**, 5845593.
<https://doi.org/10.1016/j.theriogenology.2005.06.004>
- Hunter RHF.** 1981. *Physiology and Technology of Reproduction in Female Domestic Animals*. Academic Press Limited, London.

- Kulaksiz R, Ari UC, Daskin A, Uner AG.** 2013. The effect of different glycerol concentrations on freezability of semen from agora, kilis, and saanen goats. *Slovak Journal of Animal Science* **46**, 39-44
http://www.cvzv.sk/slju/13_2/1Kulaksiz.pdf
- Kumar S, Sahni KL, Mohan G.** 1992. Effects of different levels of glycerol and yolk on freezing and storage of buffalo semen in milk, Tris and sodium citrate buffers. *Buffalo Journal* **2**, 151-156.
- Leboeuf B, Restall B, Salamon S.** 2000. Production and storage of goat semen for artificial insemination. *Animal Reproduction Science* **62**, 113-141.
[https://doi.org/10.1016/S0378-4320\(00\)00156-1](https://doi.org/10.1016/S0378-4320(00)00156-1)
- Lemma A.** 2011. Effect of cryopreservation on sperm quality and fertility. In: M. Manafi (editor). 2011. *Artificial Insemination in Farm Animals*. Accessed from
<https://www.intechopen.com/books/artificial-insemination-in-farm-animals/effect-of-cryopreservation-on-sperm-quality-and-fertility>
- Mahmoud KGM, El-Sokary AAE, El-Roos MEAA, Ghaffar ADA, Nawito M.** 2013. Sperm characteristics in cryopreserved buffalo bull semen and field fertility. *Iranian Journal of Applied Animal Science* **3**, 777-783.
http://ijas.iaurasht.ac.ir/article_513585.html
- Martinez RRD, Hernandez J, Reyna L, Santos R.** 2007. Intrauterine insemination with frozen-thawed semen in Creole goats, synchronized in oestrous during the non-breeding season. *Research Journal of Animal Science* **1**, 76-80.
<http://medwelljournals.com/abstract/?doi=rjnasci.2007.76.80>
- Mazur P.** 1984. Freezing of living cells: Mechanisms and implications. *American Journal of Physiology* **247**, 125-142.
<https://doi.org/10.1152/ajpcell.1984.247.3.C125>
- McKinnon AO.** 1999. Breeding and its technology - now and the future. Accessed from
www.harness.org.au/g9wldcon/CONFEREN.HTM.
- Morrell JM.** 2006. Update on semen technologies for animal breeding. *Reproduction Domestic Animal* **41**, 63-67.
<https://doi.org/10.1111/j.1439-0531.2006.00621.x>
- Mughal DH, Ijaz A, Yousaf MS, Wadood F, Farooq U, Mahmood SA, Riaz A.** 2018. Effect of osmotic pressure on spermatozoa characteristics of cryopreserved buffalo bull (*Bubalus bubalis*) semen. *Journal of Applied Animal Research* **46**, 274-277.
<https://doi.org/10.1080/09712119.2017.1295971>
- Naseer Z, Ahmad E, Singh J, Ahmad N.** 2011. Fertility following CIDR based synchronization regimens in anoestrous Nili-Ravi buffaloes. *Reproduction Domestic Animal* **46**, 814-817.
<https://doi.org/10.1111/j.1439-0531.2010.01746.x>
- Neglia G, Gasparrini B, Palo RD, Rosa CD, Zicarelli L, Campanile G.** 2003. Comparison of pregnancy rates with two estrus synchronization protocols in Italian Mediterranean Buffalo cows. *Theriogenology* **60**, 125-133.
[https://doi.org/10.1016/S0093-691X\(02\)01328-6](https://doi.org/10.1016/S0093-691X(02)01328-6)
- Parks JE, Graham JK.** 1992. Effects of cryopreservation procedures on sperm membranes. *Theriogenology* **38**, 209-222.
[https://doi.org/10.1016/0093-691X\(92\)90231-F](https://doi.org/10.1016/0093-691X(92)90231-F)
- Paul V, Prakash BS.** 2005. Efficacy of the ovsynch protocol for synchronization of ovulation and fixed-time artificial insemination in Murrah buffaloes (*Bubalus bubalis*). *Theriogenology* **64**, 1049-1060.
<https://doi.org/10.1016/j.theriogenology.2001.02.004>
- Rizal M, Riyadhhi M.** 2016. Fertilitas semen kerbau rawa (*Bubalus bubalis carabanensis*) yang diencerkan dengan pengencer nira aren. *Jurnal Veteriner* **17**, 457-467.

<https://doi.org/10.19087/jveteriner.2016.17.3.457>

Sansone G, Nastri MFJ, Fabbrocini A. 2000. Storage of buffalo (*Bubalus bubalis*) semen. *Animal Reproduction Science* **62**, 55-76.

[https://doi.org/10.1016/S0378-4320\(00\)00154-8](https://doi.org/10.1016/S0378-4320(00)00154-8)

Seeram NP, Adams LS, Henning SM, Niu Y, Zhang Y, Nair MG, Heber D. 2005. In vitro antiproliferative, apoptotic and antioxidant activities of punicalagin, Ellagic acid and a total pomegranate tannin extract are enhanced in combination with other polyphenols as found in pomegranate juice. *Journal of Nutritional Biochemistry* **16**, 360-367.

<http://dx.doi.org/10.1016/j.jnutbio.2005.01.006>

Sianturi RG, Purwantara B, Supriatna I, Amrozi, Situmorang P. 2012. Effect of glutathione and bovine seminal plasma in lactose extender on viability of swamp buffalo frozen semen. *Jurnal Ilmu Ternak dan Veteriner* **17**, 169-178.

<http://dx.doi.org/10.14334/jitv.v17i3.697>

Singh A, Singh V, Narwade B, Mohanty T, Atreja S. 2012. Comparative quality assessment of buffalo (*Bubalus bubalis*) semen chilled (5°C) in egg yolk and soya milk-based extenders. *Reproduction Domestic Animal* **47**, 596-600.

<https://doi.org/10.1111/j.1439-0531.2011.01928.x>

Singh P, Kumar D, Kumar P, Singh I, Yadav PS. 2013. Cryopreservation and quality assessment of buffalo bull semen collected from farmer's doorstep. *Agricultural Research* **2**, 148-52.

<https://link.springer.com/article/10.1007/s40003-013-0056-8>

Su L, Li X, Quan J, Yang S, Li Y, He X, Tang X. 2007. A comparison of the protective action of added egg yolks from five avian species to the cryopreservation of bull sperm. *Animal Reproduction Science* **104**, 212-9.

<https://doi.org/10.1016/j.anireprosci.2007.06.019>

Stryer L. 1995. *Biochemistry*. WH Freeman and

Company, New York.

Swelum AA, Mansour HA, Elsayed AA, Amer HA. 2011. Comparing ethylene glycol with glycerol for cryopreservation of buffalo bull semen in egg-yolk containing extender. *Theriogenology* **76**, 833-842.

<https://doi.org/10.1016/j.theriogenology.2011.04.015>

Tarig AA, Wahid H, Rosnina Y, Yimer N, Goh YM, Baiee FH, Khumran AM, Salman H, Assi MA, Ebrahimi M. 2017. Effect of different concentrations of soybean lecithin and virgin coconut oil in Tris-based extender on the quality of chilled and frozen-thawed bull semen. *Veterinary World* **10**, 672-678.

<http://dx.doi.org/10.14202/vetworld.2017.672-678>

Tezcan F, Ozguven MG, Diken T, Ozcelik B, Erim FB. 2009. Antioxidant activity and total phenolic, organic acid and sugar content in commercial pomegranate juices. *Food Chemistry* **115**, 873-877.

<https://doi.org/10.1016/j.foodchem.2008.12.103>

Wadood F, Aleem M, Ijaz A, Ahmad M, Yousaf M, Mughal H. 2016. Effect of extender osmolality and butylated hydroxy toluene supplementation on post-thaw quality and fertility of Nili Ravi buffalo bull (*Bubalus bubalis*) semen. *The Journal of Animal and Plant Science* **26**, 605-611.

<https://pdfs.semanticscholar.org/9785/09633a22a4f46fc0cd32201d60fb953b7319.pdf>

Warriach HM, Channa AA, Ahmad N. 2008. Effect of oestrus synchronization methods on oestrus behaviour, timing of ovulation and pregnancy rate during the breeding and low breeding seasons in Nili-Ravi buffaloes. *Animal Reproduction Science* **107**, 62-67.

<https://doi.org/10.1016/j.anireprosci.2007.06.007>

Watson PF. 2000. The causes of reduced fertility with cryopreserved semen. *Animal Reproduction Science* **60-61**, 4815492.

[https://doi.org/10.1016/S0378-4320\(00\)00099-3](https://doi.org/10.1016/S0378-4320(00)00099-3)

Willoughby CE, Mazur P, Peter AT, Kritser JK. 1996. Osmotic tolerance limits and properties of murine spermatozoa. *Biology of Reproduction* **55**, 715-727.

<https://doi.org/10.1095/biolreprod55.3.715>

Yulnawati, Gunawan M, Maheshwari H, Rizal M, Herdis, Boediono A. 2010. Quality of ejaculated and epididymal sperms spotted buffalo in dextrose supplemented extender. *Hayati Journal of Biosciences* **17**, 27-30.

<https://doi.org/10.4308/hjb.17.1.27>

Yulnawati, Maheshwari H, Rizal M, Boediono A. 2013. Frozen-thawed epididymal sperm quality and the success rate of artificial insemination in spotted buffaloes (*Bubalus bubalis carabanensis*). *Buffalo Bulletin (Special Issue 2)* **32**, 494-497.

<http://ibic.lib.ku.ac.th/e->

[Bulletin/IBBUSI201302039.pdf](http://ibic.lib.ku.ac.th/e-Bulletin/IBBUSI201302039.pdf)