BIODIVERSITAS Volume 22, Number 1, January 2021 Pages: 278-287

Birds in the west coast of South Kalimantan, Indonesia

MAULANA KHALID RIEFANI^{1,}, MOCHAMAD ARIEF SOENDJOTO^{2,3,}

¹Faculty of Teacher Training and Education Sciences, Universitas Lambung Mangkurat. Jl. Hasan Basry, Kayutangi, Banjarmasin 70123, South Kalimantan, Indonesia. Tel.: +62-511-3304914, *email: maulanakriefani@ulm.ac.id

²Faculty of Forestry, Universitas Lambung Mangkurat. Jl. Ahmad Yani Km 36, Banjarbaru 70714, South Kalimantan, Indonesia

Tel.: +62-511-4772290, "email: masoendjoto@ulm.ac.id

³Program of Education Biology, Graduate Program, Universitas Lambung Mangkurat. Jl. Hasan Basry, Kayutangi, Banjarmasin 70123, South

Kalimantan, Indonesia

Manuscript received: 29 November 2020. Revision accepted: 23 December 2020.

Abstract. Riefani MK, Soendjoto MA. 2020. Birds in the west coast of South Kalimantan, Indonesia. Biodiversitas 22: 278-287. There is no comprehensive data on birds in the west coast of South Kalimantan, yet the areas are pressured by human activities, especially the rapidly growing tourism. This study aimed to investigate the diversity of bird species that occurred in the west coast of South Kalimantan and to compare the similarities of bird communities across four villages used as the sampling sites, i.e. Sungai Rasau, Sungai Bakau, Tabanio, and Pagatan Besar. Transect method combined with concentration count method were used to record diurnal bird species and number of individuals in five replications from April to September 2020. The observation area was 100 m from the shoreline toward the sea and 400 m from the shoreline toward the inland along 2 km with observation time was 07.00-11.00 and 16.00-19.00. Shannon Wienner species diversity index (H'), a Dice community similarity index based on the presence or absence of species (ISD), and a community similarity index (modification) based on the number of individuals per species (ISM) were analyzed. In total, there are 101 bird species recorded belonging to 41 families with Sungai Bakau Village has the highest number of species (81 species), while Pagatan Besar Village is the lowest (62 species). Thirty-four species are categorized as waterbirds. Twenty-three species are categorized as migratory birds. In terms of diversity index from highest to lowest are Sungai Bakau (4.04), Sungai Rasau (3.95), Pagatan Besar (3.65), and Tabanio (3.48). The similarity of the ISD community ranges from 0.74-0.84, while the ISM ranges from 0.73-0.90. One case (Sungai Bakau - Tabanio) shows that community similarity based on the number of individuals per species is smaller than based on the presence or absence of species. Based on the protection status following Indonesian government regulation, most (84.16%) birds are categorized as unprotected, while based on the conservation status of IUCN, the majority (91.09%) are Least Concern. The results of this study can be used as baseline information for sustainable management of the area in various aspects, including future research (such as on pest and disease control), tourism (such as bird watching), and education (such as field guide).

Keywords: Diurnal bird, diversity, South Kalimantan, similarity, status

INTRODUCTION

South Kalimantan is one of the five provinces in the main island of Kalimantan (Indonesian Borneo) which has coastlines. The coastlines are located in the southwest, south and east of the province. In more detail, the coastline stretches from the west of the Barito River estuary (administratively included in the Barito Kuala District which borders the Central Kalimantan Province), to the east of the Barito River estuary in Banjar District, to the south to Tanjung Selatan in Tanah Laut District, to the northeast through Tanah Laut District to Pagatan in Tanah Bumbu District, and to the north through Tanah Bumbu District and Kotabaru District to the border with East Kalimantan Province.

Like most coastal ecosystems, the west coast of South Kalimantan also consists of two habitats with different characteristics. The first part is the land/terrestrial habitat, which is located from the shoreline toward the inland. The other part is the beach, which is actually the transition area between land and sea, and is located from the shoreline towards the sea at the lowest tide. The shoreline is a virtual line along the coast which is actually the outermost part of the vegetated land. This line separates the land from the ocean, which is covered by seawater. The beach is alternately or periodically flooded and not inundated by seawater. When high tide or seawater moves inland, beach covered with vegetation or non-vegetated either (sandy/muddy beaches) appears inundated or even submerged. On the other hand, when the sea recedes or the seawater moves away from the land towards the sea, the beach becomes dry and the vegetation that grows or appears to be growing on dry land and non-vegetated areas (sandy/muddy beaches) is evident. According to the Ramsar Convention Secretariat (2016), beach is classified as wetlands. The Ramsar Convention stated that wetlands are areas of marsh, fen, peatland or water, whether natural or artificial, permanent or temporary, with water that is static or flowing, fresh, brackish or salt, including areas of marine water the depth of which at low tide does not exceed six meters.

The west coast of South Kalimantan is an interesting area to study its biological resources, especially the diversity of birds (avifauna). There are two rationales for the importance of studying bird diversity in this region. First, there is no comprehensive information on the species richness of bird in this area although preliminary surveys suggested that there were many bird species occurred on the west coast which can be classified into aquatic and nonaquatic birds. The only research is from Riefani and Arsyad (2019) who reported that among the aquatic birds that occurred in Pagatan Besar Village, there were birds that were categorized as migratory birds. Second, four villages on the west coast of South Kalimantan Province have developed tourism, yet the attracting objects have been merely focused on physical resources. The villages of Tabanio, Takisung, and Batakan, for example, make sandy beaches and seawater as tourism features. One other village, namely Pagatan Besar Village, utilizes mangrove forest as a tourist attraction. However, the growth and development of mangroves, especially api-api or white mangrove (Avicennia marina) planted by the community (Soendjoto 2019) hampered activities to observe migratory birds that used to forage on muddy beaches.

This study aimed to investigate the diversity of bird species that occurred in the west coast of South Kalimantan and to compare the similarities of bird communities across four villages used as the sampling sites. We expected the results of this study can be used as baseline information for sustainable management of the area in various aspects, including future research (such as on pest and disease control), tourism (such as bird watching), and education (such as field guide).

MATERIALS AND METHODS

Study area

Data were collected from four villages in the west coast of South Kalimantan which were directly adjacent to the Java Sea, namely Sungai Rasau, Sungai Bakau, Tabanio, and Pagatan Besar (Figure 1). Detailed information on geographical site and habitat characteristics are presented in Table 1.



Figure 1. Map of four research sites located in the west coast of South Kalimantan Province, Indonesia

Table 1. Site of data collection,	geographical coor	dinates and habitat	characteristics
-----------------------------------	-------------------	---------------------	-----------------

Site code	Site	Geographical coordinates	Habitat characteristics
SR	Sungai Rasau Village, Bumi Makmur	3°33'30"S - 3°34'35"S;	Muddy/sandy flat, mangrove forest, coastal forest,
	Sub-district, Tanah Laut District	114°34'06"E - 114°34'50"E	fishpond, rice field, grass field, settlements
SB	Sungai Bakau Village, Kurau Sub-	3°42'26" S - 3°43'48"S;	Muddy/sandy flat, mangrove forest, coastal forest,
	district, Tanah Laut District	114°37'01"E - 114°37'34"E	fishpond, rice field, grass field, settlements
TB	Tabanio Village, Takisung Sub-	3°45'28"S - 3°46'38"S;	Muddy/sandy flat, mangrove forest, coastal forest, rice
	district, Tanah Laut District	114°36'28"E - 114°36'46"E	field, grass field, settlements
PB	Pagatan Besar Village, Takisung	3°47'56"S - 3°49'08"S;	Muddy flat, mangrove forest, grass field (grass height is
	Sub-district, Tanah Laut District	114°36'14"E - 114°36'32"E	about 5 cm; this meadow is used as a pasture for
			livestock), settlements

Data collection

Data collection was carried out 5 times between April-September 2020 using transect and the concentration count methods. The occurrences of diurnal birds were recorded in an area of 100 m from the shoreline toward the sea and 400 m from the shoreline toward the inland along 2 km twice a day at 07.00-10.00 and 16.00-19.00. Along the transect, we walked slowly while identifying species and counting the number of individuals that were seen directly perching or flying on both sides of the transect within a maximum distance of 100 m if the land was open or 25 m if the vegetation (trees) was dense. In the concentration count method, we stopped at a certain point (in this case at a densely vegetated or forested area that was thought to be a gathering place or a completely open area where the birds gathered together), observed, identified the species, and counted the number of individuals. The tools used were GPS, binocular (8x40), counter and camera with telephoto lens. Birds were identified according to MacKinnon et al. (2010), Soendjoto et al. (2015), or Soendjoto et al. (2019). The taxonomical name was referred to the IUCN (2020).

Data analysis

The name of bird species and the number of individuals in five replications were tabulated and counted. Then, species diversity and community similarity indices were calculated using the following formula.

Species diversity index Shannon-Wienner;

$$H' = -\sum \frac{m}{N} \ln \left(\frac{m}{N}\right)$$

Where, ni = number of individuals of species i; N = number of individuals of all species; ln = natural logarithm.

Community similarity index Dice;

$$IS_D = \frac{2A}{2A+B+C}$$

Modified community similarity index;

$$IS_{M} = \frac{ni(A+B)}{ni(A+B) + njA + nkB}$$

Where, ISD = Dice community similarity index calculated based on the presence or absence of species; ISM = community similarity index whose equation is a modification of the ISD and the calculation is based on the number of individuals of each species from the communities being compared; A = number of species present at the two compared sites; B = number of species present only at the first site; C = number of species present only at the second site; ni (A + B) = the number of individuals of species *i* present at site A and at the same time at site B; nA = number of individuals of species *j* only at site A; and nkB = the number of individuals of the *k* species that exist only at site B.

In addition, the status of the bird species protection was determined according to Regulation of the Minister of Environment and Forestry, Republic of Indonesia No. P.106/MENLHK/SETJEN/KUM.1/12/2018 concerning Amendment to Regulation of the Minister of Environment and Forestry No P.20.MENLHK/SETJEN/KUM.1/6/2018 on The Protected Flora and Fauna, while the conservation status assigned according to the IUCN (2020).

RESULTS AND DISCUSSION

Species diversity

In the west coast of South Kalimantan, a total of 101 bird species belonging to 41 families were recorded (Table 2). The village of Sungai Bakau has the richest species with 81 species, Sungai Rasau Village with 78 species, Tabanio Village with 62 species, while Pagatan Besar Village is the lowest with 62 species. The sites with the highest to lowest diversity of bird species are Sungai Bakau (4.04), Sungai Rasau (3.95), Pagatan Besar (3.69), and Tabanio (3.48).

E	To Jamasia mana	Tertere altere la comp	Number of individuals (ni) Status					
r anniy/species indonesia name internasional name		SR	SB	TB	PB	P.106	IUCN	
Acanthizidae								
Gerygone sulphurea	Remetuk laut	Golden-bellied Gerygone	21	24	8	15	TD	LC-d
Accipitridae								
Elanus caeruleus	Elang tikus	Black-winged Kite	5	2	1	1	D	LC-s
Haliaeetus leucogaster	Elang laut perut putih	White-bellied Sea-eagle	1	2	1	1	D	LC-d
Haliastur indus	Elang bondol	Brahminy Kite	3	1	1	2	D	LC-d
Nisaetus cirrhatus	Elang brontok	Changeable Hawk-eagle	1	-	-	-	D	LC-i
Acrocephalidae	C	0						
Acrocephalus orientalis	Kerakbasi besar	Oriental Reed-warbler	1	-	-	-	TD	LC-d
Aegithinidae								
Aegithina tiphia	Cipoh kacat	Common Iora	25	18	14	20	TD	LC-u
Aegithina viridissima	Cipoh jantung	Green Iora	4	4	8	6	TD	NT-d
Alcedinidae	1 5 0							
Alcedo meninting	Raja-udang meninting	Blue-eared Kingfisher	1	4	1	3	TD	LC-d
Halcyon smyrnensis	Cekakak belukar	White-breasted Kingfisher	4	2	1	-	TD	LC-i
Pelargopsis capensis	Pekaka emas	Stork-bellied Kingfisher	3	3	1	4	TD	LC-d
Todiramphus chloris	Cekakak sungai	Collared Kingfisher	43	18	28	22	TD	LC-d
Todiramphus sanctus	Cekakak suci	Sacred Kingfisher	6	29	12	7	TD	LC-i

Table 2. Species name, number of individuals and conservation status of bird at the four study sites in the west coast of South Kalimantan, Indonesian

Anatidae						
Anas gibberifrons	Itik benjut	Sunda Teal	5	6	-	-
Dendrocygna arcuata	Belibis kembang	Wandering Whistling-duck	12	15	-	-
Apodidae						
Apus nipalensis	Kapinis rumah	House Swift	23	20	9	-
Collocalia linchi	Walet linci	Cave Swiftlet	27	24	21	22
Ardeidae				-		
Ardea purpurea	Cangak merah	Purple Heron	-	2	-	-
Ardeola speciosa	Blekok sawah	Javan Pond-heron	24	45	8	9
Butorides striata	Kokokan laut	Green-backed Heron	7	9	1	5
Egretta alba	Kuntul besar	Green White Egret	12	4	-	-
Egretta garzetta	Kuntul kecil	Little Egret	42	10	2	4
Ixobrychus cinnamomeus	Bambangan merah	Cinnamon Bittern	4	5	1	-
Ixobrychus sinensis	Bambangan kuning	Yellow Bittern	4	8	1	-
Artamidae						
Artamus leucoryn	Kekep babi	White-breasted Woodswallow	16	14	8	9
Campephagidae						
Lalage nigra	Kapasan kemiri	Pied Triller	8	6	5	12
Pericrocotus igneus	Sepah tulin	Fiery Minivet	4	-	-	-
Caprimulgidae						
Caprimulgus affinis	Cabak kota	Savanna Nightjar	8	13	5	6
Charadriidae						
Charadrius dubius	Cerek kalung-kecil	Little Ringed Plover	7	62	-	-
Charadrius leschenaultii	Cerek-pasir besar	Greater Sandplover	9	33	-	85
Charadrius mongolus	Cerek-pasir mongolia	Lesser Sandplover	-	20	-	20
Pluvialis fulva	Cerek krenyut	Pacific Golden Plover	-	34	-	27
Ciconiidae						
Leptoptilos javanicus	Bangau tongtong	Lesser Adjutant	2	1	-	-
Cisticolidae						
Orthotomus ruficeps	Cinenen kelabu	Ashy Tailorbird	23	24	11	16
Orthotomus sericeus	Cinenen merah	Rufous-tailed Tailorbird	2	-	2	-
Prinia flaviventris	Prenjak rawa	Yellow-bellied Prinia	8	18	1	6
Columbidae						
Chalcophaps indica	Delimukan zamrud	Emerald Dove	-	-	4	-
Geopelia striata	Perkutut jawa	Zebra Dove	3	8	1	5
Spilopelia chinensis	Tekukur	Eastern Spotted Dove	4	20	2	6
Treron vernans	Punai gading	Pink-necked Green-Pigeon	4	12	2	6
Cuculidae						
Cacomantis merulinus	Wiwik kelabu	Plaintive Cuckoo	3	2	3	5
Cacomantis sonneratii	Wiwik lurik	Banded Bay Cuckoo	-	1	-	-
Centropus bengalensis	Bubut alang-alang	Lesser Coucal	10	10	3	4
Centropus sinensis	Bubut besar	Greater Coucal	3	4	3	4
Chrysococyx minutillus	Kedasi laut	Little Bronze-cuckoo	1	-	-	-
Dicaeidae						
Dicaeum trigonostigma	Cabai bunga api	Orange-bellied Flowerpecker	-	-	1	-
Dicaeum trochileum	Cabai jawa	Scarlet-headed Flowerpecker	9	7	10	8
Estrildidae						
Lonchura fuscans	Bondol kalimantan	Dusky Munia	10	27	6	5
Lonchura malacca	Bondol rawa	Tricoloured Munia	12	18	8	8
Lonchura punctulata	Bondol peking	Scally-breasted Munia	39	13	27	10
Hirundinidae						
Delichon dasypus	Layang-layang rumah	Asian House-martin	-	-	-	15
Hirundo tahitica	Layang-layang batu	Tahiti Swallow	15	14	11	10
Laniidae						
Lanius schach	Bentet kelabu	Long-tailed Shrike	16	20	8	22
Laridae						
Chlidonias leucopterus	Dara laut sayap putih	White-winged Tern	9	14	-	-
Gelochelidon nilotica	Dara-laut tiram	Common Gull-billed Tern	5	-	-	-
Sterna hirundo	Dara-laut biasa	Common Tern	15	16	15	16
Sterna sumatrana	Dara-laut tengkuk-hitam	Black-naped Tern	7	-	-	-
Sternula albifrons	Dara-laut kecil	Little Tern	-	29	-	-

Thalasseus bergii

Merops philippinus

Merops viridis

Meropidae

Dara-laut jambul

Kirik-kirik laut

Kirik-kirik biru

Great Crested Tern

Blue-tailed Bee-eater

Blue-throated Bee-eater

NT-s

LC-d LC-i

LC-d

LC-d

LC-u

LC-d

LC-u

LC-i

LC-s

LC-u

LC-s

LC-d

NT-d

LC-s

LC-s

LC-d

LC-u

LC-d

VU-d

LC-s LC-s

LC-d

LC-d

LC-s

LC-i

LC-s

LC-s

LC-s

LC-i

LC-s

LC-s

LC-s

LC-s

LC-s

LC-s

LC-s

LC-i

LC-u

LC-u

LC-s

LC-d

LC-u

LC-u

LC-d

LC-s

LC-s

LC-s

TD

D

TD

D

D

D

D

D

D

TD

TD

12

8

5

-

4

31

34

13

11

-

9

14

BIODIVERSITAS 22 (1): 278-287, January 2021

Motacillidae								
Anthus novaeseelandiae	Apung tanah	Australasian Pipit	1	3	2	3	TD	LC-s
Motacilla flava	Kicuit kerbau	Western Yellow Wagtail	2	2	1	-	TD	LC-d
Muscicapidae			-	-	-	-		
Cyornis rufigastra	Sikatan bakau	Mangrove Blue-flycatcher	-	-	1	1	TD	LC-d
Nectariniidae		с .						
Aethopyga siparaja	Burung-madu sepah-raja	Crimson Sunbird	2	3	2	1	D	LC-s
Anthreptes malacensis	Burung-madu kelapa	Brown-throated Sunbird	15	12	11	6	TD	LC-s
Cinnvris iugularis	Burung-madu sriganti	Olive-backed Sunbird	13	19	14	4	TD	LC-s
Leptocoma calcostetha	Burung-madu bakau	Copper-thorated Sunbird	1	-	-	-	TD	LC-s
Pandionidae			-					
Pandion haliaetus	Elang tiram	Osprev	1	-	-	-	D	LC-i
Passeridae	214119 11411	osproj	-				2	201
Passer montanus	Burung gereia erasia	Furasian Tree Sparrow	22	21	41	23	TD	I C-d
Pellorneidae	Burung gereja erasia	Eurasian free Sparlow	22	21	71	25	10	LC u
Malacocincla seniaria	Pelanduk semak	Horsfield's Babbler	_	_	2	_	ТD	I C-d
Dicidaa	I clanduk semak	Horsheld's Dabbler			2		ID	LC-u
Chrysophlagma miniacaum	Pelatuk merah	Banded Woodpecker	_	_	2	_	тD	I C-s
Diagidas moluceansis	Caladi tilik	Sunda Dygmy Woodnocker	- 5	- 5	1	- 2		
Picolaes moluccensis		Sunda Fyginy Woodpecker	5	5	1	5	ID	LC-I
Ditta condida	Deals biim	Western Heeded Ditte			2		D	
	Paok injau	western Hooded Pitta	-	-	3	-	D	LC-û
Ploceidae	M	C 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2	10			TD	IC
Ploceus manyar	Manyar Jambul	Streaked weaver	2	10	-	-	ID	LC-s
Pycnonotidae			50	(1		50	ΠD	
Pycnonotus aurigaster	Cucak kutilang	Sooty-headed Bulbul	53	61	76	50	TD	LC-d
Pycnonotus brunneus	Merbah mata merah	Red-eyed Bulbul	15	4	44	2	TD	LC-d
Pycnonotus goiavier	Merbah cerukcuk	Yellow-vented Bulbul	15	34	26	37	TD	LC-1
Pycnonotus plumosus	Merbah belukar	Olive-winged Bulbul	4	-	4	-	TD	LC-s
Rallidae								
Amaurornis cinerea	Tikusan alis-putih	White-browed Crake	-	1	-	-	TD	LC-u
Amaurornis phoenicurus	Kareo padi	White-brested Waterhen	3	9	3	12	TD	LC-u
Recurvirostridae								
Himantopus leucocephalus	Gagang bayam timur	White-headed Stilt	12	8	-	3	TD	LC-i
Rhipiduridae								
Rhipidura javanica	Kipasan belang	Sunda Pied Fantail	15	11	13	8	TD	LC-s
Scolopacidae								
Actitis hypoleucos	Trinil pantai	Common Sandpiper	11	12	3	6	TD	LC-d
Calidris alba	Kedidi putih	Sanderling	-	40	-	5	TD	LC-u
Calidris ruficollis	Kedidi leher merah	Rufous-necked Stint	-	41	-	18	TD	NT-d
Gallinago megala	Berkik rawa	Swinhoe's Snipe	-	-	4	-	TD	LC-u
Limosa lapponica	Biru laut ekor-blorok	Bar-tailed Godwit	7	13	-	8	TD	NT-d
Numenius madagascariensis	Gaiahan timur	Far Eastern Curlew	_	4	-	7	D	EN-d
Numenius phaeopus	Gajahan penggala	Whimbrel	-	6	-	-	D	LC-d
Tringa glareola	Trinil semak	Wood Sandniper	5	24	-	8	TD	LC-s
Tringa nebularia	Trinil kaki-hijau	Common Greenshank	-	11	-	-	TD	LC s
Tringa stagnatilis	Trinil rawa	Marsh Sandniner	5	8	_	_	TD	LC J
Yanus cinaraus	Trinil badaran	Terek sandpiper	10	23	_	28	TD	
Strigidaa		Telek sandpiper	10	23	-	20	ID	LC-u
Strigiuae	Dalult Itatuna	Buffy Fish and		2			тD	IC.
	Calarrala rahar	Sunda Saama and	-	2	-	-		LC-S
Otus lempiji	Celepuk reban	Sunda Scops-owi	-	-	1	-	ID	LC-s
Sturmdae	77 1 1 1	T M	25		25	7	TD	1 7 1 1
Acriaotheres javanicus	Kerak kerbau	Javan Myna	25	22	25	/	ID	vu-a
Timaliidae	<i>a</i>		1.5	25	_	10	TD	I.C.
Mixornis gularis	Clung-air coreng	Pin-striped Tit-babbler	15	35	1	13	TD	LC-s
Vangidae								
Hemipus hirundinaceus	Jinjing batu	Black-winged Flycatcher-shrike	-	4	-	1	TD	LC-d
Zosteropidae								
Zosterops flavus	Kacamata jawa	Javan White-eye	3	-	-	-	TD	EN-d
Zosterops palpebrosus	Kacamata biasa	Indian White-eye	2	8	-	-	TD	LC-d
		Number of individuals (N)	817	1242	576	705		
		Number of species (S)	78	81	63	62		
		Species diversity index (H')	3 95	4 04	3 4 8	3 69		

Species diversity index (H') 3.95 4.04 3.48 3.69 Note: 1) Site of data collection: SR: Sungai Rasau, SB: Sungai Bakau, TB: Tabanio, PB: Pagatan Besar. 2) Protection status according to Minister of Environment and Forestry, Republic of Indonesia No P.106/MENLHK/SETJEN/KUM.1/12/2018 (2018): D: protected, TD: unprotected. 3) Conservation status according to IUCN (2020): EN: endangered, VU: vulnerable, NT: near threatened, LC: least concern, i: increasing, d: decreasing, s: stable, u: unknown Most (66.34% or 67 species) of the 101 bird species are categorized as non-waterbirds, while the rest (34.66%) are waterbirds. Eight waterbird families in the west coast include of Anatidae, Ardeidae, Charadriidae, Ciconiidae, Laridae, Rallidae, Recurvirostridae, and Scolopacidae, and are among 34 families of waterbirds in the world described by Ducks Unlimited New Zealand (2017) or among 33 families described by Wetlands International (2020).

Twenty-one of 34 waterbird species are categorized as migratory birds, including Greater Sandplover, Lesser Sandplover, Pacific Golden Plover, Little Ringed Plover, Common Gull-billed Tern, Little Tern, White-winged Tern, Great Crested Tern, Common Tern, White-headed Stilt, Far Eastern Curlew, Marsh Sandpiper, Whimbrel, Common Sandpiper, Terek Sandpiper, Wood Sandpiper, Common Greenshank, Swinhoe's Snipe, Sanderling, Rufous-necked Stint, and Bar-tailed Godwit. Based on the colored flag attached to the bird's leg and referring to the meaning of the flag color in EAAFP (2018), R. Yuliansyah (2020, pers. com.) found Sanderling of South Australia on the sandy stretch of the Sungai Bakau, and T.K. Utomo (2020, pers. com.) found Rufous-necked Stint of China-Chongming Island in Pagatan Besar.

Besides the migratory waterbirds, there are also nonwaterbirds that are categorized as migrants. They are Osprey, the raptor and Blue-tailed Bee-eater, the insectivore. Osprey has been observed soaring above waters and ponds and then perching on the dry tree of the Sungai Rasau. The presence of Osprey at that site indicates that the migratory raptors (Germi et al. 2009, Bierregaard et al. 2014, Purwanto et al. 2015) or these wetland raptors (Irham 2012) make the west coast as a stopover site. In the southern hemisphere, these birds that spread around the world become non-breeding visitors in Indonesia (van Balen 1998, Purwanto et al. 2015) as well as breeding visitors in Australia (Clancy 2006; Dennis and Clancy 2014, Detmar and Dennis 2018). Osprey was recorded in February-March 2016 flying around an artificial lake in the Tarjun Cement Factory Area, Kotabaru (Riefani et al. 2019).

Blue-tailed Bee-eater and Blue-throated Bee-eater are migratory birds in Malaysia (Nisbet 2013) and Thailand (DeCandido et al. 2010). There has not been any specific publication on the bee-eaters as migratory birds in Indonesia. According to Soendjoto et al. (2018) and Riefani et al. (2019), Blue-tailed Bee-eater was not always found, while Blue-throated Bee-eater was almost always found in 2013-2017 observations in Tabalong District, South Kalimantan and more than 300 hours of field observation in Kotabaru District, South Kalimantan. Therefore, we only include Blue-tailed Bee-eater as migratory birds and not Blue-throated Bee-eater as one of the migratory birds in South Sumatra and do not mention Blue-throated Bee-eater as such.

The following two bird species were also found on the west coast. Both are highlighted in this paper because their presence and distribution are not stated in field books, such as Smythies (1960) and MacKinnon et al. (2010). Therefore, these findings can be considered as new records

that are important for updating data or revising field books. The first is White-breasted Kingfisher which was found in Tanah Laut District, to be precise at Sungai Rasau, Sungai Bakau, and Tabanio. The presence of the bird was observed 9 times, both solitary and 2 individuals, adding to new information that this bird is indeed distributed in Kalimantan Selatan or even Kalimantan in general. Smythies (1960) did not mention the presence of this bird in Borneo. Avibase (2020b) even said that not a single subspecies of White-breasted Kingfisher existed in Indonesia. According to Avibase (2020b), the birds spreading across the Arabian Peninsula to the Caucasus Mountains and the northwestern part of India are Halcyon smyrnensis smyrnensis; in West India and Sri Lanka H.s. fusca, in the Andaman H.s. saturatior; as well as in Myanmar to Peninsular Malaysia and Indochina H.s. perpulchra. MacKinnon et al. (2010) stated that the distribution of this bird in Sumatra and Java. According to Strange (2012), its distribution was even only in Sumatra. The distribution of White-breasted Kingfisher in Sumatra is supported by the report of Ayat (2011), Arief et al. (2015), Kamal et al. (2016), Santosa et al. (2016), Syamsi (2017), and Iswandaru et al. (2018), while in Java there was the report by Taufiqurrahman et al. (2019). White-breasted Kingfisher was also found in West and South Kalimantan (Myers 2016), industrial plantation companies in West Kalimantan (Ata Marie and Ekologika Consultans 2017), Sebangau National Park, Central Kalimantan (Adi 2018), oil palm plantations in Central Kalimantan (Santosa et al. 2018), and the Liang Anggang Protected Forest Area, Kalimantan Selatan (Wulansari et al. 2020).

The second is Streaked Weaver which was found breeding in Sungai Rasau and Sungai Bakau. In the observations conducted in July 2020, an average of 7.3 nests hung per coconut tree (*Cocos nucifera*) (n = 10) at about 20 m above ground level. More nests (up to 32 nests) were found hanging from branches of the Acacia auriculiformis tree at about 6-10 m above ground level. The tree is on the roadside of Sungai Bakau Village. The distribution of Streaked Weaver in Tanah Laut District adds to new record in primary publications. Early records suggest that the distribution of Streaked Weaver in Indonesia is only in Java and Bali (MacKinnon et al. 2010) and not recorded in Borneo (Smythies 1960, MacKinnon et al. 2010). IUCN (2020) said that this bird spreads in Southeast Asia (Cambodia, Myanmar, Singapore, Thailand, Vietnam), South Asia (India, Bangladesh, Bhutan, Nepal, Pakistan, Sri Lanka), China, and even Egypt. Particularly in Indonesia, the map attached to the IUCN report (2020) shows that the distribution of Streaked Weaver is consistent with MacKinnon et al. (2010) mentioned above. According to Avibase (2020a), the Streaked Weaver spreading across Indonesia (the islands of Java, Bali and Bawean) is a subspecies of Ploceus manyar manyar; in eastern Pakistan to western India and Sri Lanka P.m. flaviceps; in Northeast India (Assam) to Bangladesh and northern Myanmar P.m. peguensis; and in Southwest China (Yunnan) to Thailand and Vietnam P.m. williamsoni.

The presence of Streaked Weaver in Kalimantan was published by Soendjoto et al. (2014) and Iqbal et al. (2016).

The nests hung from the branches of rambai or mangrove apple (*Sonneratia caseolaris*) and rumbia or sago palm (*Metroxylon sago*) trees. From interviews with the community, Soendjoto et al. (2016) stated that this bird had been detected in Banjar District in 2011. It is likely that the birds were brought by bird traders from Java Island and then accidentally released. Birds can also be deliberately released by keepers who are tired of raising animals or are not careful with raising them after buying them from traders.

In a field trip in 2017, we even found 3 occurrences of this bird with its nest hanging individually on bundung or greater club rush (Cyperaceae: *Scirpus grossus*), one of the swamp plants around Sungai Puting, Tapin Distcrit. On each stem of 1-1.5 m high there was only one nest attached. In 2018 the weaver's nest was found hanging on a branch of the mangrove apple at a height of about 20 m above the ground in the swamplands of Lambung Mangkurat University, Banjarmasin City. Streaked Weaver is considered to be pests, along with Dusky Munia, Dusky Munia, Scally-breasted Munia, and Eurasian Tree Sparrow because they eat rice grown by the people living in the settlements west of the university campus complex.

It is important to note that the states of the bird species richness and diversity indices are temporary (subject to change) and only apply to this study due to the following reasons. First, this study focuses on diurnal birds, so that nocturnal birds are not observed and identified. Observation and identification of only certain diurnal bird species, for example, those that are small in size and inhabit mangrove forests, coastal forests, or forests with thick canopy cover or those that are migrants, require a longer duration or frequency of observation. Observation and identification of nocturnal bird species are relatively challenging. Certain strategies or techniques need to be developed because what is faced is not only the dark sight, but also the thickness of the canopy and/or tidal inundation. In this study, two nocturnal bird species that were found by chance were Buffy Fish-owl and Sunda Scops-owl. It is likely that there are other nocturnal bird species at the sites.

Second, one sub-district on the west coast (Panyipatan sub-district) has not been represented in the study. However, the following findings of bird species from a wildlife photographer can be used as a guide that there are actually other birds on the west coast. Lesser Frigatebird (*Fregata andrewsi*), Great Tit (*Parus major*), Black-bellied Malkoha (*Phaenicophaeus diardi*), Buff-banded Rail (*Hypotaenidia philippensis*), and Copper-throated Sunbird (*Nectarinia calcostetha*) were sighted in the area by P. Irawan (2020, pers. com.).

Community similarity

The community similarity index between sites based on the presence or absence of species (0.77-0.84) is lower than that based on the number of individuals per species (0.79-0.90) (Table 3). The exception is the index based on the presence or absence of species between Tabanio and the Sungai Bakau which is slightly higher than that based on the number of individuals of each species. Since the index based on the number of individuals for each species is generally higher or closer to 1.00, this index is better to use to measure community similarity between two sites.

The similar resources with adequate availability are the factors that cause the similarity of bird communities in the two compared sites to be categorized as high. These resources include food, nesting, microclimate and occurrence of predators. In the case of migratory bird species, both intra and inter continents, birds leave certain sites and come to other sites because they have the same resource. Site and season cause the availability of prey and species requirements for energy to vary and in turn the proportion of foraging birds also varies (Burton et al. 2004). The presence and diversity of migratory birds in various habitat types is largely determined by microhabitat, availability of food sources, and safety in resting and foraging areas (Sonobe and Usui 1993, Howes et al. 2003).

In the Sungai Bakau, the feed source area is disturbed by tidal inundation. Some waterbirds, or to be precise in this case are shorebirds, generally forage on mudflats that lie after the shoreline towards the sea. They forage in colonies within the same species and even with other waterbird species. However, when the high tide inundates or submerges the muddy areas, they move to and forage on other muddy areas, such as active ponds, non-operational ponds or former ponds. In this non-vegetated inland muddy field, they can still stand on the surface of the mud and their beaks can still reach the food on the surface or in the mud. Moving and looking for food in the mud inland is a strategy for waterbirds to survive. In general, the waterbird diet consists of plants, such as terrestrial plants, seeds and aquatic macrophytes, and animals, such as zooplankton, snails, shells, worms, crabs, mollusks, crustaceans, arthropods, larvae, and imago of terrestrial and aquatic insects (Coleoptera, Diptera, Heteroptera, Odonata), nectonic macroinvertebrates, fish, amphibians, and water snakes (Ntiamoa-Baidu et al. 1998, Horváth et al. 2012). The feed is obtained easily when the mud bed is not inundated by seawater and facilitates the birds to behave according to their morphology; for example, the soles and toes may step on the mud and the beak reaches for feed in the mud.

Table 3. Bird community similarity index between sites

Site	Sungai Rasau	Sungai Bakau	Tabanio	Pagatan Besar
Sungai Rasau	1.00	0.87	0.90	0.85
Sungai Bakau	0.84	1.00	0.73	0.88
Tabanio	0.78	0.74	1.00	0.79
Pagatan Besar	0.77	0.84	0.78	1.00

Note: The numbers below the diagonal line (top left - bottom right) are the community similarity index based on the presence or absence of species, and while above the diagonal line are that based on the number of individuals of each species.

The conditions that make water birds move are not only the tidal mud area, but also the growth and development of mangrove species. In Pagatan Besar village, for example, the muddy area which was originally an open area has gradually turned into a mangrove forest. The development of mangrove forests makes it difficult for migratory birds who are accustomed to foraging for food together (in colonies) on muddy plains in open areas. This change was observed from four ironwood walkways, which according to Soendjoto (2019) have 1.5 m wide and about 100 m long, jutting from the shoreline towards the Java Sea. The height of the walkway floor is 0 m from the ground level at the starting point of the walkway (at the shoreline) and about 2 m from the ground level (mud) at the end of the walkway (towards the sea). At Titian 1, which is located at the northernmost point, mangrove forests which are dominated by white mangrove are observed to be getting higher. The height in 2018 was around 4-5 m and in 2020 it is about 6 m. However, the sea and the mud can still be observed from the end of the walkway. At Titian 2, which is located 490 m to the south of Titian 1, the sea and mud can still be observed from half the length of the walkway to the end of the path in 2018. In 2020, the sea and mud can no longer be observed. The white mangroves reach 3-4 m in high and are covering the view. At Titian 3, which is located 190 m to the south of Titian 2, the sea and mud can be observed from the base to the end of the bridge in 2018 because the height of the white mangroves growing on the left and right is only about 0.5 m from the surface of the mud. In 2020, the height of the flames is 2 m around the base of the walkway and about 1 m around the end of the walkway. Several individual white mangroves had grown out of the direction of the sea from the end of the bridge. In other words, mangrove forests are growing or expanding. The sea and mudflats can still be observed starting from one-third of the length of the walkway to the end of the path. Titian 4 which is located 180 m to the south from Titian 3, the sea and mud can still be observed freely. The white mangroves are still few in number. In other words, the density is very low compared to the density of the white mangroves around Titian 1, 2, and 3. The height is also about 0.5 m above the ground (mud).

The stretch of mud on the west coast is formed by sediments carried by the Maluka River, Tabanio River, Pagatan Besar River, and Barito River as the widest river in South Kalimantan and then deposited in the Java Sea, where these rivers flow into. The main sediments on the Barito River are sand, clay and silt (Arisanty et al. 2014). Because the Barito River is the main route for coal transportation (Novico and Priohandono 2012), coal shale is also observed in the mud. According to Arisanty et al. (2012), shorelines can change and the changes are caused by human activities, such as logging, forest and land fires, intensive agriculture, and mining.

Bird status

Based on protection status of the regulation of Minister of Environment and Forestry, Republic of Indonesia No P.106/MENLHK/SETJEN/KUM.1/12/2018 (2018), most bird species (84.16%) are categorized as unprotected. Only 16 species (15.84%) are protected. Based on their conservation status of IUCN (2020), most bird species (91.09%) are categorized as Least Concern with a tendency of population to decrease (30 species), increase (13 species), stable (36 species), and unknown (13 species). The rest are categorized as Near Threatened (5 species, namely Green Iora, Sunda Teal, Fiery Minivet, Rufous-necked Stint, and Bar-tailed Godwit), Vulnerable (2 species, namely Lesser Adjutant and Javan Myna), and Endangered (2 species, namely Far Eastern Curlew and Javan White-eye. A large number of unprotected bird species while at the same time the least concern status is of course alarming. Many bird species have not been studied intensively, but their sustainability or survival is threatened.

Currently, people catch birds for at least two purposes. First, birds are used as a pet. Zebra Dove, Eastern Spotted Dove, Sooty-headed Bulbul, Yellow-vented Bulbul, and Javan Myna are kept because of their voice. Unfortunately, birds are kept without partners so they cannot reproduce. Second, bird species are consumed. White-breasted Waterhen, Yellow Bittern, and Wandering Whistling-duck are species that are traded and then enjoyed for their meat. The over catching and consumption of birds has definitely wiped them out.

In addition, land conversion, especially the protected areas or conservation areas into production areas also threatens the bird. Forest-dwelling birds who like lush bushes or under-canopy dwellers such as Western Hooded Pitta, Ioras, Pin-striped Tit-babbler, and Emerald Dove have lost their habitats. In contrast, birds that are able to adapt to open areas or grass vegetation find new habitats. They include Eurasian Tree Sparrow, Australasian Pipit, Western Yellow Wagtail, and Javan Myna.

Each bird species has its own ecological roles and functions. As carnivores, eagles and Brahminy Kite help maintain the balance of nature. Sooty-headed Bulbul, Yellow-vented Bulbul, and flowerpeckers are frugivores that help spread plant seeds from certain sites to other sites, so do Zebra Dove, Eastern Spotted Dove, and munias as granivores. Long-tailed Shrike, bee-eaters, and Sunda Pied Fantail as insectivores play a role in controlling pests and diseases in plants. Sunbirds as nectarivores not only act as nectar eaters, but also helps pollinate.

In conclusion, 101 bird species found in the west coast of South Kalimantan are classified as minimal. The species diversity index of the four sample sites ranges from 3.48 to 4.04. In general, the similarity index based on the number of individuals per species is greater than that based on the presence or absence of the species. Migratory birds can be developed as unique attractions or ecotourism objects on this west coast. White-breasted Kingfisher and Streaked Weaver are worth noting in field books when these birds have been found spreading in Kalimantan.

ACKNOWLEDGEMENTS

This research was funded by PNBP of Lambung Mangkurat University, Banjarmasin, Indonesia in the Fiscal

Year of 2020. We express our gratitude to the Rector of Lambung Mangkurat University who approved the funding for this research. Thanks also go to the Head of Tanah Laut District, Indonesia who allowed the research to be carried out in his administrative area and the enumeration team (Nur Abdi Suga) who helped collect data in the field.

REFERENCES

- Adi. 2018. Bird Species Identification in Sungai Kaki and Sebangau Kecil. https://www.tnsebangau.com/kegiatan-identifikasi-jenisburung-di-sungai-kaki-dan-sebangau-kecil. [Indonesian]
- Arief H, Mijiarto J, Rahman A. 2015. Diversity and protection status of wildlife in PT Riau Sawitindo Abadi. Media Konservasi 20 (1): 159-165. DOI: 10.29244/medkon.20.2.%25p [Indonesian].
- Arisanty D, Sartohadi J, Marfai MA, Hadmoko DS. 2012. The Long-term morphodynamic of Barito Delta, Southern Kalimantan, Indonesia. J Environ Sci Eng B 1: 1196-1202.
- Arisanty D, Sartohadi J, Marfai MA, Hadmoko DS. 2014. Sediment dynamic in Barito Delta, Southern Kalimantan, Indonesia. J Environ 1 (1): 30-37.
- Ata Marie and Ekologika Consultans. 2017. Summary Report High Conservation Value Assessment PT Hutan Ketapang Industri-Kabupaten Ketapang, West Kalimantan Province. Ata Marie & Ekologika Consultans, Jakarta.
- Avibase. 2020a. Streaked Weaver *Ploceus manyar* (Horsfield, 1821). https://avibase.bsc-eoc.org.
- Avibase. 2020b. White-throated Kingfisher Halcyon smyrnensis (Linnaeus, 1758). https://avibase.bsc-eoc.org.
- Ayat A. 2011. Agroforest Birds in Sumatra. World Agroforestry Centre-ICRAF, SEA Regional Office, Bogor, Indonesia. [Indonesian]
- Bierregaard RO, Poole AF, Washburn BE. 2014. Ospreys (Pandion haliaetus) in the 21st century: Populations, migration, management, and research priorities. J Raptor Res 48 (4): 301-308. DOI: 10.3356/0892-1016-48.4.301
- Burton NHK, Musgrove AJ, Rehfisch MM. 2004. Tidal variation in numbers of waterbirds: how frequently should birds be counted to detect change and do low tide counts provide a realistic average? Bird Stud 51: 48-57.
- Clancy GP. 2006. The breeding biology of the Osprey Pandion haliaetus on the North Coast of New South Wales. Corella. J Austr Bird Stud Assoc 30 (1): 1-8.
- Decandido R, Nualsri C, Allen D. 2010. Mass northbound migration of Blue-tailed *Merops philippinus* and Blue-throated *M. viridis* Beeeaters in southern Thailand, spring 2007-2008. Forktail 26: 42-48.
- Dennis TE, Clancy GP. 2014. The status of the Osprey (*Pandion haliaetus cristatus*) in Australia. J Raptor Res 48 (4): 1-7. DOI: 10.3356/JRR-OSPR-14-01.1
- Detmar SA, Dennis TE. 2018. A review of Osprey distribution and population stability in South Australia. South Australian Ornithologist (43): 1-2.
- Ducks Unlimited New Zealand. 2017. What are waterbirds? https://ducks.org.nz.
- EAAFP. 2018. Shorebird Color Flagging Protocol on the East Asian-Australasian Flyway. https: //eaaflyway.net/wpcontent/uploads/2017/12/Protocol_birds-marking.pdf
- Germi F, Young GS, Salim A, Pangimangen W, Schellekens M. 2009. Over-ocean raptor migration in a monsoon regime: spring and autumn 2007 on Sangihe, North Sulawesi, Indonesia. Forktail 25: 104-116.
- Horváth Z, Ferenczi M, Móra A, Vad CF, Ambrus A, Forró L, Szövényi G, Andrikovics S. 2012. Invertebrate food sources for waterbirds provided by the reconstructed wetland of Nyirkai-Hanyy, northewestern Hungary. Hydrobiologia 697: 59-72. DOI: 10.1007/s10750-012-1170-5.
- Howes J, Bakewell D, Noor YR. 2003. Panduan Studi Burung Pantai. Wetlands International Indonesia Program, Bogor. [Indonesian]
- Iqbal M, Irawan P, Fauzi A, Rahman I. 2016. Introduced Species: Streaked Weaver *Ploceus manyar*, a new species for Kalimantan, Borneo. Birding ASIA 25: 71-72.
- Irham M. 2012. Avifauna diversity at Central Halmahera North Maluku, Indonesia. Zoo Indonesia 21 (1): 17-31.

- Iswandaru D, Khalil ARA, Kuniawan B, Pramana R, Febryano IG, Winarno GD. 2018. Abundance and diversity of bird species in Mangrove Forest KPHL Gunung Balak. Indonesian J Conserv 7 (1): 57-62. [Indonesian]
- IUCN [International Union for Conservation of Nature]. 2020. The IUCN Red List of Threatened Species. Version 2020-2. https: //www.iucnredlist.org
- Kamal S, Agustina E, Rahmi Z. 2016. Bird species in some habitat types in Kecamatan Lhoknga, Kabupaten Aceh Besar. Jurnal Biotik 4 (1): 15-32. [Indonesian]
- MacKinnon J, Phillipps K, van Balen B. 2010. LIPI- Field Guide of Birds in Sumatra, Kalimantan, Java and Bali. Puslitbang Biologi- LIPI, Bogor. [Indonesian]
- Myers S. 2016. Helm Field Guides, Birds of Borneo, 2nd ed. NHBS, UK.
- Nash SV, Nash AD. 1985. A checklist of the forest and forest edge birds of the Padang-Sugihan Wildlife Reserve, South Sumatra. Kukila 2 (3): 51-59.
- Nisbet ICT. 2013. Seasonal occurrence, migrations and habitat relationships of Blue-tailed and Blue-throated Bee-eaters *Merops philippinus* and *M. viridis* in Peninsular Malaysia. Raffles Bull Zool Suppl. 29: 219-223.
- Novico F, Priohandono YA. 2012. Analysis of erosion and sedimentation patterns using software of Mike 21 HDFM-MT in The Kapuas Murung River Mouth Central Kalimantan Province. Bull Mar Geol 27 (1): 35-53.
- Ntiamoa-Baidu Y, Piersma T, Wiersma P, Poot M, Battley P, Gordon C. 1998. Water depth selection, daily feeding routines and diets of waterbirds in coastal lagoons in Ghana. Ibis 140 (1): 89-103. DOI: 10.1111/j.1474-919X.1998.tb04545.x
- Purwanto AA, Rakhman Z, Supriatna AA, Sutito ASB, Srirejeki I. 2015. Current information on migratory raptors and its conservation efforts in Indonesia. Asian Raptors 1: 54-62.
- Rachman R, Sudiarty N, Ismawati, Sumbawati NK, Hun S. 2020. Analysis of household income in swiftlet cultivation in Karang Dima Village, Labuhan Badas District. Prosiding Seminar Nasional IPPeMas 2020 Inovasi Hasil Penelitian dan Pengabdian kepada Masyarakat dalam Menunjang Era Industri 4.0. [Indonesian]
- Ramsar Convention Secretariat. 2016. An Introduction to the Convention on Wetlands (previously The Ramsar Convention Manual). Ramsar Convention Secretariat, Gland, Switzerland.
- Riefani MK, Arsyad M. 2019. Bird species in Mangrove Ecotourism Area of Pagatan Besar, Tanah Laut Regency, Indonesia. Prosiding Seminar Nasional Lingkungan Lahan Basah 4 (1): 192-196. [Indonesian]
- Riefani MK, Soendjoto MA, Munir AM. 2019. Bird species in the cement factory complex of Tarjun, South Kalimantan, Indonesia. Biodiversitas 20 (1): 218-225. DOI: 10.13057/biodiv/d200125
- Santosa RA, Harianto SP, Nircahyani N. 2016. Comparison of the population of Kingfisher bird (Halcyonidae) in the wetlands of Sungai Luar Village and the wetlands of Kibang Pacing Village, Menggala Timur District, Tulang Bawang Regency. Jurnal Sylva Lestari 4 (2): 79-88. DOI: 10.23960/jsl2479-88. [Indonesian]
- Santosa Y, Sugiharti W, Erniwati. 2018. The variation of bird diversity in different land cover at oil palm plantation: Case study at Asm Oil Palm Estate in Central Kalimantan, Indonesia. E3S Web of Conferences 52, 00051. DOI: 10.1051/e3sconf/20185200051.
- Smythies BE. 1960. The Birds of Borneo. The Sabah Society & The Malaya Nature Society, Kuala Lumpur, Malaysia.
- Soendjoto MA, Nugroho Y, Suyanto, Riefani MK, Supandi, Yudha HES. 2019. Avifauna in the Area of PT Borneo Indobara Kalimantan Selatan. Banyubening, Banjarbaru. [Indonesian]
- Soendjoto MA, Riefani MK, Ready A. 2014. New record: Streaked Weaver (*Ploceus manyar*) in Kalimantan Selatan. Warta Konservasi Lahan Basah 22 (3): 16-17. [Indonesian]
- Soendjoto MA, Riefani MK, Triwibowo D, Metasari D. 2018. Birds observed during the monitoring period of 2013-2017 in the revegetation area of ex-coal mining sites in South Kalimantan, Indonesia. Biodiversitas 19 (1): 323-329. DOI: 10.13057/biodiv/d190144
- Soendjoto MA, Riefani MK, Triwibowo D, Wahyudi F. 2015. Avifauna in Reclamation Area of PT Adaro Indonesia. Universitas Lambung Mangkurat Press, Banjarbaru, Indonesia. [Indonesian]
- Soendjoto MA. 2019. Potential and problems in development of the ecotourism area (Case in the Pagatan Besar Mangrove Forest, Tanah Laut Regency, Indonesia). Prosiding Seminar Nasional Lingkungan Lahan Basah 4 (3): 635-642.

- Sonobe K, Usui S. 1993. A Field Guide to the Waterbirds of Asia. Wild Bird Society of Japan, Tokyo.
- Strange M. 2012. A Photographic Guide to the Birds of Indonesia. 2nd ed. Tuttle Publishing, Tokyo.
- Sudiyanto T. 2018. The impact of the swiftlet business on changes in the economy of the Betung Banyuasin community. Jurnal Media Wahana Ekonomika 12 (3): 101-109. [Indonesian]
- Syamsi F. 2017. Bird diversity in wetland area of Kota Batam. Simbiosa 6 (1): 17-30. [Indonesian]
- Taufiqurrahman I, Purwanto AA, Tirtaningtyas FN, Ade S, Rachman M, Daryana A, Hermawan AS, Yordan K, Emmanuel B, Kristanto A, Kiryono, Purnama S, Sujatmiko DA. 2019. Recent records of White-

breasted Kingfisher Halcyon smyrnensis on Java, Indonesia, with notes on its status and distribution. BirdingASIA 31: 74-78.

- van Balen SB. 1998. Tropical forest raptors in Indonesia: Recent information on distribution, status, and conservation. J Raptor Res 32 (1): 56-63.
- Wetlands International. 2020. Waterbird Population Estimates. https://wpe.wetlands.org.
- Wulansari A, Soendjoto MA, Suyanto. 2020. Bird species diversity in natural vegetation of Liang Anggang Protected Forest Area- Block I, Banjarbaru, Indonesia. Jurnal Sylva Scienteae 3 (5): 818-824. [Indonesian]