

# Bacillus thuringiensis As Local Biological Agent

*by* Akhmad Rizali

---

**Submission date:** 09-Oct-2021 09:51AM (UTC+0700)

**Submission ID:** 1669252306

**File name:** sis\_As\_Local\_Biological\_Agent\_9-ICGT-2018\_paper\_3-dikonversi.pdf (122.07K)

**Word count:** 2202

**Character count:** 12178

## *Bacillus thuringiensis* As Local Biological Agent

Akhmad Rizali<sup>1</sup>

<sup>1</sup>Department of Agrotechnology, Lambung Mangkurat University. Indonesia

### Abstract

The chemical insecticides are still contributing to human life enormously, but they have been distributed in ecological system of organisms including human beings because of their low specific toxicity to any organism and their low specific toxicity to any organism and their slight decomposition in nature. Therefore, many biological control of insects have been investigated. Currently, researches on the use pathogenic microorganisms to control insect pests are increasing. Microbial pest control is practiced in different parts of the world though utilization of pathogen like fungi, bacteria, viruses and nematodes. Bacterial research causing disease in insects began in the late nineteenth century. It was a study of flacherie of the silkworm, *bombx mori* in this report on the discovery of *sotto bacillus*, referred briefly to occurrence of sotto bacillus-like organism, which causes the disease to silkworm larvae. Ten-fold serial dilutions of the heated suspension in sterile distilled water were placed on nutrient agar (NA-pH 7.5). After two days of incubation at 28°C, *Bacillus* colonies were recorded. After 2 to 3 days incubation, crystalliferous sporeforming bacteria were determined in phase contrast microscop. Isolation from five soil samples yielded about 35 isolates, only one was identified as *B. thuringiensis*.

Key Word: *B. thuringiensis*. Biological Agent

### Introduction

Cabbage moth, *C. binotalis* Zell. (Lepidoptera: Pyraustidae) is considered the most important limiting factor for a successful production of cruciferous vegetable not only in the Indonesia but

in other country in the world. The larva feeds on folage from seedling to harvest causing 100% yield loss if not control (Rejesus and Sayaboc, 1990).

Numerous chemical insecticides have been used in order to control pests, which

damage for agriculture. While chemical insecticides have knock down effect to the insect pests, they are too expensive in the developing countries and harmful to both human and the environment. In addition, target insect pests rapidly develop biological resistance especially at higher rates of application. The chemical insecticides are still contributing to human life enormously, but they have been distributed in ecological system of organisms including human beings because of their low specific toxicity to any organism and their low specific toxicity to any organism and their slight decomposition in nature (Shorey and Hall, 1962). Therefore, many biological control of insects have been investigated. Currently, researches on the use pathogenic microorganisms to control insect pests are increasing. Microbial pest control is practiced in different parts of the world though utilization of pathogen like fungi, bacteria, viruses and nematodes. Bacterial research causing disease in insects began in the late nineteenth

century. It was a study of flacherie of the silkworm, *bombx mori* (Burgess and Hussey, 1971; Burgess, 1981). Ishiwata (1901) in this report on the discovery of *sotto bacillus*, referred briefly to occurrence of sotto bacillus-like organism, which causes the disease to silkworm larvae .

Berliner (1911) proposed the name of *B. thuringiensis* for a species of bacillus which was isolated from the diseased larvae of the Mediterranean flour moth *Anagasta (Ephestia) kuhniella* Zell. Later, Berliner (1915) noted infection of the larvae after the ingestion of the bacillus or its spore, described and named it *Bacillus thuringiensis*. Mattes (1927) isolated the same bacillus from the same insect host, which Berliner had found earlier.

*B. thuringiensis* is a gram-positive soil bacterium, and produce a crystalline inclusion body during sporulation (Bulla et al., 1980). This parasporal body is composed of proteins termed “delta-endotoxin”, and specifically toxic to

insects. In addition, *B. thuringiensis* produce another toxins namely: alpha-toxin, beta-exotoxin, and gamma-exotoxin. All of the toxic substance may not present in the bacterium (Heimpel, 1967). In another hand, Krieg (1961) has defined various toxic substance produced *B. thuringiensis* as follow: (a) thermolabile endotoxic; (b) thermostable exotoxin; (c) bacillogenic antibiotic; (d) lecithinase; (e) proteinase.

Most strains of *B. thuringiensis* produce delta-endotoxin crystals toxic to lepidopteran insects such as moth (Dulmage et al., 1970). Recently, however several researches have shown that *B. thuringiensis* is also widely distributed in natural soils of various area. Delucca et al., (1982) reported that *B. thuringiensis* made up less than 0.5% of more than 46,000 bacterial isolates recovered from various soils in the United States.

The objective of the studies to survey, collect and determine the distribution of

*B. thuringiensis* in selected diverse crop-growing area.

### **Materials and Methods**

#### *Isolation of B. thuringiensis*

Soil samples were collected in areas planted to vegetables, rice, citrus, peanut and corn in South Kalimantan (Indonesia) following multistage random sampling. Soil samples were collected at random in a 1- hectare area for a total of 5 kg. The soil sample were taken from the top 1 cm of the soil layer. The 5-kg soil samples were mixed thoroughly and composite sample of 1 kg was taken from which isolation were made for as long as one month. The samples were labeled denoting date, place of collection and crops planted. Five 1-g soil samples were separately suspended to 9 ml of distilled water. After allowing the suspension to stand for 5 minute, 3-4 ml of the the suspension were taken. One half of the suspension was transferred to a test tube and heated in a waterbath of 80°C for 15

minutes, so that all microorganisms were killed except *Bacillus* and other sporeforming bacteria, then allowed to cool at room temperature. Ten-fold serial dilutions of the heated suspension in sterile distilled water were placed on nutrient agar (NA-pH 7.5). After two days of incubation at 28°C, *Bacillus* colonies were recorded. After 2 to 3 days incubation, crystalliferous sporeforming bacteria were determined in phase contrast microscop.

## RESULTS AND DISCUSSION

Isolation and distribution of *B. thuringiensis* in different plants growing areas.

Fifty *Bacillus sp.* were isolated from 5 soil samples collected from diverse crop growing areas in the South Kalimantan (Indonesia) (Table 1). The different isolates were obtained from the same area planted with diverse crop. Two soil samples great were found in this area namely: typic calciborrols with pH 6.5

was planted to rice and typic tropudults pH ranging from 4.8 – 5.7 planted to vegetable, corn, citrus, peanut and cabbage. Out of the 35 *Bacillus sp.* isolates, only one (2%) was identified as *B. thuringiensis* based on phase contrast microscope examination for the presence of parasporal inclusion bodies. Only the soil sample from citrus yielded *B. thuringiensis*.. The possible reasons for the low incidence of *B. thuringiensis* isolated from the samples taken in the areas surveyed are the small number of samples size from which the isolation were made, the area where sampling was done and also the difference in the physic-chemical characteristic of the soil where samples were taken. The low incidence of *B. thuringiensis* was also reported in Japan by Ohba and Aizawa (1966) in soil sample from non-agricultural areas. Out of 6910 isolates only 189 (2.7%) isolates were identified as *B. thuringiensis*

Table 1. South Kalimantan Isolates with different plants areas source

ISOLATES	SOIL SOURCE		
	Crop Planted	Great Soil	pH
AA	Watermelon	Typic calciborolls	6.5
AA 1.1			
AA 1.2			
AA 2.1 (1)			
AA 2.2			
AA 2.3			
AA 4.1			
AA 2.5			
AA 2.5 (1)			
AA 2.5 (2)			
BB	Cucumber	Typic tropudults	4.8
BB 1.1			
BB 1.2			
BB 2.3			
BB 1.4			
BB 3.1			
CC	Corn	„	5.2
CC 1.2			
CC 2.1			
CC 2.3			

CC 2.4

CC 3.2

CC 5.1

CC 5.1 (1)

DD            Citrus            ,,            5.7

DD 1.2 (1)

DD 1.3

EE            Peanut            ,,            5.1

EE 1

EE 2

EE 1.3

EE 1.4

EE 3.1

EE 4.3

---

### Conclusion

Six soil samples were collected from different crop growing areas in South Kalimantan (Indonesia). Isolation from five soil samples yielded about 35 isolates, only one was identified as *B. thuringiensis*. This

result suggest that *B. thuringiensis* is rare and not widespread in the places where sampling was done and isolates of *B. thuringiensis* will screen to the cabbage worm (*Crocidolomia binotalis*).

## Reference

Abdel-Hameed, A., Carlberg, G. and El-Tayeb, O. M (1990). Studies on *Bacillus thuringiensis* H-14 strains isolated in Egypt. Screening for active strains. World Journal of Microbiology and Biotechnology. 6: 299-304.

Asano, S., Bando, H. and Iizuka T. (1993). Amplification and identification of *cryII* genes from *Bacillus thuringiensis* by PCR procedures. J. Seric. Sci. Jpn. 62: 223-227.

Baba, F., Asano, S. and Iizuka T. (1990). Purification of crystals from *Bacillus thuringiensis* by using Percoll. J. Sci. Jpn 59: 487-489.

Balarman, K., Hoti, S. L. and Manonmani, L. M (1981). An Idigenous virulent strain of *Bacillus thuringiensis*, highly pathogenic and specific to mosquitoes. Current Science 50: 199-200.

Berliner, E. (1915). Ober die schalaffsucht der mehmottenraupe (*Ephestia kuhniella* Zeller) und ihren erreger *Bacillus thuringiensis* n. sp. Zangue. Entomol. 2: 29-56.

Bulla, L. A., faust, R. M., Andrews, R. and Goodman, N. (1985). Insecticidal bacilli, pp. 185-209. In: the Moleculer Biology of the Bacilli, Volume II. D. A. Dubnau (ed.). Academic Press, New York.

Burges, H.D. 1981. Microbial control of pest and plant disease. 1970-1980. Academic Press. 949 pp.

Burges, H.D. and N.W. Hussey, 1971. Microbial control of insects and mites. Academic Press. 876 pp.

Berliner, E. (1915). Ober die schalaffsuch der Mehmottenraupe (*Ephestia kuhniella* Zeller) und ihren erreger *Bacillus thuringiensis* n. Sp. Z angue. Entomol. 2: 29-56.

Bulla, L. A., Jr., Kramer, K. J. and Davidson, L. I. (1977). Characterization of the enmocidal parasporal crystal of *Bacillus thuringiensis*. J. Bacteriol. 130: 375-383.



Delucca, A. J. II, Simonson, J. G and Larson, A. D. (1981). *Bacillus thuringiensis* distribution in soils of the United States. *Canadian J. Microbiol.* 27: 865-870.

Dulmage, H. T. ( 1992). Insecticidal activity of *Bacillus thuringiensis* and their potential for pest control in *Microbial control for pests and plant diseases and plant diseases 1970-1980* (ed.H.D Burges). Acad. Press. N.Y. PP.

Golberg, L. J. and Margalit, J. (1977). A bacterial spore demonstrating rapid larvicidal activity against *Anopheles sergentii*, *Uranotaenia unguiculata*, *Culex univittatus*, *Aedes aegypti* and *Culex pipiens*, *Mosq. New* 37: 355-358.

Heimpel, A. M. (1967). A critical review of *Bacillus thuringiensis* Berl. And other crystalliferous bacteria. *Ann. Rev. Entomol.* 12: 287-322.

Held, G. A., Kawanishi, C. Y. and Huang, Y. -S. (1990). Characterization of the parasporal inclusion of *Bacillus thuringiensis* subsp. *Kyushuensis*. *J. Bacteriol.* 481-483.

Iizuka, T., Ishino, M. and Nakajima, T (1982). Comparative morphology of Parasporal crystal and characterization of plasmid DNA from various subspecies of entomopathogenic bacteria, *Bacillus thuringiensis*. *J. Fac. Agric. Hokkaido Univ.* 13: 423-431.

Iizuka, T. and Yamamoto, T. (1984). Serological properties of the mosquitoicidal protein of *Bacillus thuringiensis* and the morphology of its parasporal crystal. *J. fac. Hokkaido Univ.* 62: 98-114.

Iizuka, T., Sasaki, J., Asano, S. and Bando, H. (1995). Comparative studies on isolation and identification of *Bacillus thuringiensis*. *Biotechnology and Environ. Benefits*, Vol. I, 143-153.

Ishii, T. and Ohba, M. (1997). Investigation of mosquito-specific larvicidal activity of a soil isolate of *Bacillus thuringiensis* serovar *Canadensis*. *Curr. Microbiol.* 35: 40-43.

Ishiwata. S. (1901). On a kind of severe flacherie (sotto disease). *Dainihon Sanshi kaiho* 114: 1-5.

Kalman, S., Kiehne, K. K., Libs, J. L. and Yamamoto, T. (1993). Cloning of novel cryIC-type gene from a strain *Bacillus thuringiensis* subs. *Galleriae*. *Appl. Environ. Microbiol.* 59: 1131-1137.

Kawalek, M. D., Benjamin, S., Lee, H. L. and Gill, S. S. (1995). Isolation and identification of novel toxin from a new mosquitocidal isolate from Malaysia, *Bacillus thuringiensis* subsp. *Jegathesan*. *Apl. Environ. Microbiol.* 2965-2969.

Kim, K. H., Ohba, M. and Aizawa, K. (1984). Purification of the toxic protein from *Bacillus thuringiensis* serotype 10 isolate demonstrating a preferential larvicidal activity to mosquito. *J. Invertebr. Pathol.* 44: 214-219.

Krieg, A. (1961). *Bacillus thuringiensis* Berliner. In disease caused by certain sporeforming bacteria. Heimpel and Angus (eds.). 21-67.

Mattes, O. (1927). Parasitare krankheiten der mehnottenlarvaen und versuche ober ihre verwundbarkeitals biologische becampfungsmittel. (Zuheich lire beitrags zur zytologie de bacgerien). *Gesell f. befond, gedam, naturw. Sitzber (Marbnog)* 62: 381-417.

Ohba, M. and Aizawa, K. (1986). Insect toxicity of *Bacillus thuringiensis* isolated from soils of Japan. *J. Invertebr. Pathol.* 47: 12-20.

Padua, L. E., Ohba, M. and Aizawa, K. (1984). Isolation of a *Bacillus thuringiensis* strain (serotype 8a:8b) highly and selectively toxic against mosquito larvae. *J. Invertebr. Pathol.* 44: 12-17.

Rejesus, B. M. and A. Sayaboc, (1990). Management of DBM with *Apanteles plutellae* prospects in the Philippines. Paper presented in the second International Workshop on Diamondback Moth Management. Dec. 10-15, 1990. Tainna, Taiwan. 17 pp.

Rizali, A., Shin-ichiro Asano, Ken Sahara, Hisanori Bando, Bibiana W, Lay, Sugyo Hastowo and Toshihiko Iizuka. (1998). Novel *Bacillus thuringiensis* serovar aizawai strains isolated from mulberry leaves in Indonesia. *Appl. Entomol. Zool.* 33 (1): 111-114.

Shorey, H. H. and I. M. Hall. (1962). Effect of chemical and microbial insecticides on several insect pests of lettuce in southern California. J. Econ. Entomol. 56: 169-174.

Yu, Y. -M., Ohba, M. and Gill, S. S (1991). Characterization of mosquitocidal activity of *Bacillus thuringiensis* subsp. *Fukuokaensis* crystal proteins. Appl. Environ. Microbiol. 1075-1081.

Zhang, Y., Ku, Z., Chan, Z., Xu, B., Yuan, F., Chen, G., Zhong, T. and Ming, G. (1984). A new isolate of *Bacillus thuringiensis* possessing high toxicity against the mosquitoes. Acta Microbiologica Sinica 24: 320-325

# Bacillus thuringiensis As Local Biological Agent

---

## ORIGINALITY REPORT

---

15%

SIMILARITY INDEX

6%

INTERNET SOURCES

11%

PUBLICATIONS

5%

STUDENT PAPERS

---

## MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

1%

★ Yoshisue, H.. "Identification of a second transcriptional start site for the insecticidal protein gene cryIVA of *Bacillus thuringiensis* subsp. *israelensis*", *Gene*, 19970207

Publication

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off