

# Comparison Study Of Toxicity *Bacillus thuringensis* from Soil Isolate and Mulberry Leaves in Indonesia

*by* Akhmad Rizali

---

**Submission date:** 07-Oct-2021 07:51AM (UTC+0700)

**Submission ID:** 1667298757

**File name:** ringensis\_from\_Soil\_Isolate\_and\_Mulberry\_Leaves\_in\_Indonesia.pdf (515.82K)

**Word count:** 3119

**Character count:** 18176



## Comparison Study Of Toxicity *Bacillus thuringiensis* from Soil Isolate and Mulberry Leaves in Indonesia

Akhmad Rizali

Department of Agroechotechnology, Lambung Mangkurat University, Banjarbaru, Indonesia

Email address:  
arizali25@yahoo.com

### To cite this article:

Akhmad Rizali. Comparison Study Of Toxicity *Bacillus thuringiensis* from Soil Isolate and Mulberry Leaves in Indonesia. *American Journal of Entomology*. Vol. 2, No. 1, 2018, pp. 1-5. doi: 10.11648/j.aje.20180201.11

33

Received: March 30, 2018; Accepted: April 19, 2018; Published: May 9, 2018

**Abstract:** Background and Objective: Most strains of *B. thuringiensis* produce delta-endotoxin crystals toxic to lepidopteran insects such moth. But some strains of *B. thuringiensis* produce delta-endotoxin crystals toxic to dipteran insects such as mosquitoes and blackflies. To compare *B. thuringiensis* isolate from soil and mulberry leaves. Material and Methods: One g of soil samples was suspended in 9 ml of sterile distilled water and shaken for 5 min. the upper layer of the soil suspension was transferred to a test tube and heated at 80°C for 5 min in water bath to kill non-spore-forming organism and vegetative cells to prepare the sporulated culture, bacteria were grown on nutrient agar pH 7.2, at 30°C for 4 days. The isolates were obtained from mulberry leaves collected in West Java, Indonesia, using the leaf-lift technique. Leaves were trimmed to fit inside a 100 mm petri dish. Abaxial leaf surfaces were placed in contact with nutrient agar, and a sterile, per forated stainless steel disk was placed on the leaf sections to ensure maximum contact with the agar. After the sample was coated with carbon and gold, it was observed and photographed with (SEM). However they had differences form crystal protein, but they had unique as insecticidal to control the same orders (mosquitocidal). Results: The study found that *B. thuringiensis* serovar *entomocidus* INA288 produced large cuboidal-form crystals and *B. thuringiensis* serovar *aizawai* BUN 1-14 a little homology with serovar *entomocidus* INA288 but, *B. thuringiensis* *israelensis* ONR60A has irregular shaped crystal protein. Toxicity results that serovar *entomocidus* INA288, serovar *aizawai* BUN 1-14 and serovar *israelensis* ONR60A showed toxic to mosquito insects. which was composed of major protein of 130 kDa peptides. *B. thuringiensis* serovar *entomocidus* INA288 has 70 kDa and *aizawai* BUN 1-14 had 69 kDa. Conclusion: *B. thuringiensis* strain had different crystal protein form but had same toxicity.

**Keywords:** *B. Thuringiensis*, Soil, Leaves

## 1. Introduction

Numerous chemical insecticides have been used to control some insects. While chemical insecticides have knock down effect, they are too expensive harmfull to both humans and the environment. In addition, target insect pests develop biological resistance rapidly especially at higher rates of application. Thus, the increase in pesticidal application to control this pest has urged to researcher to search for biological control alternatives that would be a good component of Integrated Pest Management [1, 32, 33].

*Bacillus thuringiensis* is a gram-positive, spore-forming bacterium that produce parasporal crystal during the sporulation stage. The crystal is made of one or more proteins toxic to some insect species. Most strains of *B. thuringiensis* produce delta-endotoxin crystal toxic to lepidopteran insect such as moth [2, 28, 29].

The name proposed of *Bacillus thuringiensis* for a species of *bacillus* which was isolated from diseased larvae of Mediterranean flour moth Angasta (Ephestia) kuhniella Zell [3]. Later, noted infection of the larvae after the ingestion of the *bacillus* or its spores, described and named it *Bacillus thuringiensis* [3, 4]. It isolated the same *bacillus* from the same insect host [4], which had found earlier. This strain is now maintained as *B. thuringiensis* serovar *thuringiensis* (serotype H-1). They noticed that the vegetative remains of sporulating cells assumed a rhomboid shape [3, 4]. He described this crystalline inclusion in the sporangium of the organism and made further interpretations of the data being accumulated on this bacillus at that time. Neither Berliner nor Mattes attributed those parasporal bodies any role in the disease process caused by the ingestion of sporulating *B. thuringiensis* [5].

*B. thuringiensis* is a gram-positive soil bacterium, and

produce a crystalline inclusion body during sporulation [6, 27, 28]. This parasporal body is composed of proteins termed "delta-endotoxin", and specifically toxic to insects. In addition, *B. thuringiensis* produce another toxins namely: alpha-exotoxin, beta-exotoxin, and gamma-exotoxin. All of the toxic substances may not be present in the bacterium [7, 20, 21]. In another hand, Krieg [8] has defined various toxic substance produced by *B. thuringiensis* as follow: (a) thermolabile endotoxin; (b) thermostable exotoxin; (c) bacillogenic antibiotic; (d) lecithinase; (e) proteinase.

*B. thuringiensis* has been studied world wide over the past decades, mainly because this gram-positive bacterium produce significant amount of crystal proteins with toxic activity against economically important insect larvae [9, 18, 28]. The most attractive characteristics of the *B. thuringiensis* proteins for insect control are their specificity and high unit activity. Members of non-target insect orders are not susceptible to the potent effects of the lepidopteran-specific and dipteran-specific insecticidal proteins [10, 27, 29]. Most strains of *B. thuringiensis* produce delta-endotoxin crystals toxic to lepidopteran insects such moth [11, 23, 25]. But some strains of *B. thuringiensis* produce delta-endotoxin crystals toxic to dipteran insects such as mosquitoes [12, 22-24] and blackflies.. Lepidopteran-specific delta-endotoxin are composed 130 kDa proteins [13, 15-17] while dipteran-specific delta endotoxin are composed of several protein [13, 18-20].

*Bacillus* strains possessing a high larvical activity, specific for mosquitoes, from the soil of mosquitoes-breeding site in Israel [11, 25-27]. The objective of the studies to compare *B. thuringiensis* strains serovar *entomocidus* INA288 (soil isolated), serovar *aizawai* Bun 1-14 (leaves isolated) and *israelensis* ONR60A (standard isolate).

## 2. Materials and Methods

Bacterial stains. The strain of *B. thuringiensis* used in the present study were *B. thuringiensis* serovar *entomocidus* INA 288, serovar *aizawai* BUN 1-14 and *israelensis* ONR60A (standard isolate).

### 2.1. Isolation and Identification

*B. thuringiensis* serovar *entomocidus* INA288 which had been isolated from Indonesia [3], was prepared according to the method [14]. One gram of soil samples was suspended in 9 ml of sterile distilled water [1] and shaken for 5 min. the upper layer of the soil suspension was transferred to a test tube and heated at 80°C for 5 min in water bath to kill non-spore-forming organism and vegetative cells to prepare the sporulated culture, bacteria were grown on nutrient agar pH 7.2, at 30°C for 4 days. Formation of spores and parasporal inclusion were monitored with a phase-contrast microscope. The culture was scratched on the agar slant as a stock. *B. thuringiensis* serovar *aizawai* BUN 1-14 which had been isolated from mulberry leaves. Isolates were obtained from mulberry leaves collected in West Java, Indonesia, using the leaf-lift technique [15, 16]. Leaves were trimmed to fit inside

a 100 mm petri dish. Abaxial leaf surfaces were placed in contact with nutrient agar, and a sterile, per forated stainless steel disk was placed on the leaf sections to ensure maximum contact with the agar. The lid was replaced, and samples were incubated at 30°C overnight. To prepare the sporulated culture, bacteria were grown on nutrient agar, pH 7.0, at 30°C for 4 days. Formation of spores and parasporal inclusion were monitored with a phase-contrast microscope.

### 2.2. Morphology of Parasporal Inclusion

Isolates were examined with a HITACHI S-800 Scanning Electron Microscope (SEM) at a magnification of 10,000x, according to the method presented by [23, 24]. *B. turingiensis* serovar *entomocidus* INA288 were cultured on N-broth agar at 30°C until almost all cells [28] died (overnight). The crystal and spores (about 100 mg wet weight) were washed in 10 [8] of 50 mM Tris-HCl (pH 8.0). The final precipitate was resuspended in 1 ml of distilled water, and 20 µl of the suspension was air-dried on a glass disk (O 10 mm). After the sample was coated with carbon and gold, it was observed and photographed with SEM.


### 2.3. Biological Activity

The strain were examined for their larvicidal against the larvae of the silkworm, *Plutella xylostella* and *Spodoptera litura*. The insect cultures were maintained in this laboratory. Toxicity test with the Lepidopteran insect, *B. mori*, *P. xylostella* and *S. litura*, were done by introducing ten 3<sup>rd</sup>-instar larvae were fed on an artificial diet dropped with 0.3 ml of the bacterial suspension and rear at 25°C for 48 hr to determine mortality. The *B. thuringiensis* isolates were examined for oral insecticidal activity against the insects were prepared by the following procedures. Overnight culture of serovar *entomocidus* INA288, BUN 1-14 and *israelensis* ONR60A were grown on 2 ml of nutrient broth at 30°C using tube glass. Then, 200 µl of the [13] overnight culture was plating on nutrient agar, reincubated for 4 days at 30°C. Sporulated cultures were harvested by centrifugation at 10,000 g for min at 4°C. The pellet was washed 5 [6] times by centrifugation in mMTris-HCl and 1 M NaCL at 4°C, the bacterial suspensions were finally [24] pended in 500 µl of sterile distilled water. The bacteria were also tested against larvae of the mosquitoes, *Aedes aegypti*, *Aedes japonicus* and *Culex quinquefasciatus*. Ten 2<sup>nd</sup>-instar larvae were placed in a test tube containing 10 ml of the spore-parasporal inclusion suspension, respectively, under levels 1 µl/ml. The tubes were kept at 22°C for 24 hr without feeding.

## 3. Result and Discussion

In order to identify serovar *entomocidus* INA288 strains by H-serotype cell with broth, they were dropped to glass slide and the motility of cells was observed under phase-contrast microscope. Since isolate of serovar *entomocidus* INA288 gave positive reaction in the H 6 serum agglutination test, it was identified as *B. thuringeisis* serovar *entomocidus*

INA288. interestingly, *entomocidus* INA288 produced large cuboidal-form crystals (Figure 1). On the other hand, *B. thuringiensis* serovar *aizawai* BUN 1-14 a little homology with serovar *entomocidus* INA288 (Figure 2). Interestingly, *B. thuringiensis* serovar *israelensis* ONR60A [23, 24] has irregular shaped crystal protein.



**Figure 1.** Scanning electron microscopy (SEM) showing spores and parasporal crystal of *Bacillus thuringiensis* serovar *entomocidus* INA288. Bar indicates 3 um.

*B. thuringiensis* serovar *israelensis* ONR60A had mosquitocidal activity, which was composed of major protein of 130 kDa peptides. On the other hand, *B. thuringiensis* serovar *entomocidus* INA288 and *aizawai* BUN 1-14 had mosquitocidal activity, while *aizawai* Bun 2-1 both did not have. In addition, neither serovar *entomocidus* original strain nor serovar *aizawai* IPL had lepidopteracidal activity. SDS-PAGE profiles of *entomocidus* INA288 crystals indicated that the polypeptide of 70 kDa was dominant, while *entomocidus* original strain was constituted of 130-65 kDa. However, *B. thuringiensis* serovar *aizawai* Bun 1-14 was composed of polypeptides of 69 kDa, while *entomocidus* Bun 2-1 was constituted of 130-65 kDa. Interestingly, using antibody of *B. thuringiensis* serovar *entomocidus* INA288 crystal protein demonstrated that there is cross-reactivity among the parasporal inclusion proteins of *B. thuringiensis* serovar

*aizawai* Bun 1-14, *israelensis* ONR60A, and *fukoukaensis*. However, [27] demonstrated that using antibodies of *B. thuringiensis* serovar *israelensis* ONR60A showed weakly cross-reactivity to serovar *jegathesan*. Indeed, *B. thuringiensis* serovar *jegathesan* included *cryIVD* and showed immunological similarity with antibodies of serovar *israelensis*. While, *entomocidus* INA288 contained a novel crystal protein gene *cryINA288*, on plasmid. In addition, the similarity of amino acid sequence between *cryINA288* and *cry4Aa* was 38%. On the other hand, serovar *aizawai* Bun 1-14 [27] a little homology with serovar *entomocidus* INA288 (Figure 2).



**Figure 2.** Scanning electron microscopy (SEM) showing spores and parasporal crystal of *Bacillus thuringiensis* serovar *entomocidus* INA288. Bar indicates 3 um.

#### Quantitative toxicity test isolate

The serovar entomocidus INA288, serovar *aizawai* BUN 1-14 and *israelensis*ONR60A were bioassayed against 4<sup>th</sup>-instar larvae of *B. mori*, *P. xylostella*, *S. litura*, *A. japonicas*, *A. aegypti* and *C. quinquefasciatus*. Respectively, serovar entomocidus INA288, *aizawai* BUN 1-14 showed not toxic activity against latter 3 former lepidopteran species, while serovar entomocidus INA288, *aizawai*BUN1-14 and *israelensis* ONR 60A had the toxicity against latter 3 dipteran species (*A. aegypti*, *A. japonicas* and *C. quinquefasciatus*) (Table 1). However they had differences form crystal protein, but they had unique as insecticidal to control the same orders (Mosquitoes).

**Table 1.** Toxic activity of three *B.thuringiensis* strains against some insect species.

Strain	Lepidopteracidal			Dipteracidal		
	<i>B. m.</i>	<i>P.x.</i>	<i>S.l.</i>	<i>A.j.</i>	<i>A.a.</i>	<i>C.q.</i>
Serovar <i>entomocidus</i> INA288	-	-	-	+	+	+
Serovar <i>aizawai</i> BUN 1-14	-	-	-	+	+	+
Serovar <i>israelensis</i> ONR60A	-	-	-	+	+	+

31

*B. m:* *Bombyx mori*, *P.x:* *Plutella xylostella*, *S.l:* *Spodoptera litura*  
*A.j:* *Aedes japonicas*, *A.a:* *Aedes aegypti*, *C.q:* *culex quinquefasciatus*

## 4. Conclusion

In the search for potential alternatives to the application of *B. thuringiensis* serovar *israelensis* ONR60, *entomocidus* INA288 and *aizawai* BUN 1-14 isolation of novel mosquitocidal strains. The crystal proteins form of *B. thuringiensis* serovar *entomocidus* INA288 has cuboidal shaped, serovar *aizawai* BUN 1-14 has homology shaped ones and *B. thuringiensis* serovar *israelensis* ONR60A has irregular, which was composed of major protein of 130 kDa peptides. *B. thuringiensis* serovar *entomocidus* INA288 has 70 kDa and *aizawai* BUN 1-14 had 69 kDa.

## Acknowledgements

The author wishes to express his sincerely gratitude to his adviser, Dr. TOSHIHIKO Iizuka, Professor of Bioscience at the Faculty of Agriculture, Hokkaido University. Sapporo, Japan for his excellent guidance, suggesting, constant encouragement, and critical review in the preparation of this article.

## References

- [1] Brar, S. K., Verma, M., Tyagi, R. D. and Valero, J. R. 2006. Recent advances in downstream processing and formulations of *Bacillus thuringiensis* based biopesticide. *Process Biochem.* 41(2): 323-342.
- [2] Sanahuja, G., Banakar, R., Twyman, R. M., Capell, T. and Christou, P. 2011. *Bacillus thuringiensis*: a century of research, development and commercial application, *Plant Biotechnology J.* 9: 283-300.
- [3] Berliner, E. (1915). Oberdieschaffauch der mehlmottenraupe (*Ephestiakuhniella* Zeller) und ihrererreger*Bacillus thuringiensis* n. sp. *Zangue. Entomol.* 2: 29-56.
- [4] Mattes, O. (1927). Parasitarekrankheiten der mehnottenlarvaen und versuecheoberreverwndbarkeitalsbilogischebecampfungismitt el. (Zuheich lire beitragzurzytologie de bacgerien). *Gesell f. befond, gedam, naturw. Sitzber (Marbnog)* 62: 381-417.
- [5] Rizali, A., Shin-ichiro Asano, Ken Sahara, Hisanori 6ndo, Bibiana W, Lay, SugyoHastowo and Toshihiko Iizuka.(1988). Novel *Bacillus thuringiensis* serovar *aizawai* strains isolated from mulberry leaves in Indonesia. *Appl. Entomol. Zool.* 33 (1): 111-114.
- [6] Bulla, L. A., faust, R. M., Andrews, R. and Goodman, N. (1985). Insecticidal bacilli, pp. 185-209. In: the Molecular Biology of the Bacilli, Volume II. D. A. Dubnau (ed.). Academic Press, New York.
- [7] Heimpel, A. M. 26(67). A critical review of *Bacillus thuringiensis*Berl. And other crystalliferous bacteria. *Ann. Rev. Entomol.* 12: 287-322.
- [8] Krieg, A. (1961). *Bacillus thuringiensis* Berliner. In disease caused by certain sporeforming bacteria. Heimpel and Angus (eds.). 21-67.
- [9] Delucca, A. J. II, Simonson, J. G and Larson, A. D. (1981). 19  
*Bacillus thuringiensis* distribution in soils of the United States. *Canadian J. Microbiol.* 27: 865-870.
- [10] Dulmage, H. T. (1968). Insecticidal activity of *Bacillus thuringiensis* and their potential for pest control in Microbial control for pests and plant diseases and plant diseases 1970-1980 (ed. H.D Burges). Acad. Press. N. Y. PP.
- [11] Golberg, L. J. and Margalit, J. (1977). A bacterial spore demonstrating rapid larvicidal activity against *Anopheles* *sergentii*, *Uranotaeniaunguiculata*, *Culexunivittatus*, *Aedesaegypti* and *Culexpipiens*. *Mosq. New* 37: 355-358.
- [12] Pfannenstiel, M. A., Ross, E. J., Kramer, V. C. and Nickerson, K. W. (1984). Toxicity and composition of protease-inhibited *Bacillus thuringiensis* var. *israelensis* crystal. *FEMS Microbio. Lett.* 21: 39-42.
- [13] Hastowo, S., Lay, B. W. and Ohba, M. (1982). Naturally occurring *Bacillus thuringiensis* in Indonesia. *J. Apply. Bacteriol.* 73: 108-113.
- [14] Iizuka, T., Ishino, M. and Nakajima, T (1982). Comparative morphology of Parasporal crystal and characterization of plasmid DNA from various subspecies of entomopathogenic bacteria, *Bacillus thuringiensis*. *J. Fac. Agric. Hokkaido Univ.* 13: 423-431.
- [15] Asano, S., Bando, H. and Iizuka T. (1993). Amplification and identification of *cryII* genes from *Bacillus thuringiensis* by PCR procedures. *J. Seric. Sci. Jpn.* 62: 223-227.
- [16] Asano, S. (1996) Identification of *cry* gene from *Bacillus thuringiensis* by PCR and isolation of unique insecticidal bacteria. *Mem. Fac. Agric. Hokkaido Univ.* 19: 529-563.
- [17] Baba, F., Asano, S. and Iizuka T. (1990). Purification of crystals from *Bacillus thuringiensis* by using Percoll. *J. Sci. Jpn* 59: 487-489.
- [18] Balarman, K., Hoti, S. L. and Manonmani, L. M (1981). An Idigenous virulent strain of *Bacillus thuringiensis*, highly pathogenic and specific to mosquitoes. *Current Science* 50: 199-200.
- [19] Burges, H. D. 1981. Microbial control of pest and plant disease. 1970-1980. Academic Press. 949 pp.
- [20] Burges, H. D. and N. W. Hussey, 1971. Microbial control of insects and mites. Academic Press. 876 pp.
- [21] Bulla, L. A., Jr., Kramer, K. J. and Davidson, L. I. (1977). Characterization of the entomocidal parasporal crystal of *Bacillus thuringiensis*. *J. Bacteriol.* 130: 375-383.
- [22] Old, G. A., Kawanishi, C. Y. and Huang, Y. -S. (1990). Characterization of the parasporal inclusion of *Bacillus thuringiensis* subsp. *Kyushuensis*. *J. Bacteriol.* 481-483.
- [23] Saka, T. and Yamamoto, T. (1984). Serological properties of the mosquitocidal protein of *Bacillus thuringiensis* and the morphology of its parasporal crystal. *J. fac. Hokkaido Univ.* 62: 98-114.
- [24] Iizuka, T., Sasaki, J., Asano, S. and Bando, H. (1995). Comparative studies on isolation and identification of *Bacillus thuringiensis*. *Biotechnology and Enviro. Benefits*, Vol. 1, 143-153.
- [25] Ishiwata, S. (1901). On a kind of severe flacherie (sotto disease). *Dainihon Sanshikaiho* 114: 1-5.

- 11
- [26] Kalman, S., Kiehne, K. K., Libs, J. L. and Yamamoto, T. (1993). Cloning of novel cryIC-type gene from a strain *Bacillus thuringiensis* subs. *Galleriae*. *Appl. Enviro. Microbiol.* 59: 1131-1137.
- [27] Kawalek, M. D., Benjamin, S., Lee, H. L. and Gill, S. S. (1995). Isolation and identification of novel toxin from a new mosquitocidal isolate from Malaysia, *Bacillus thuringiensis* subsp. *Jegathesan*. *Apl. Enviro. Microbiol.* 2965-2969.
- [28] Kim, K. H., Ohba, And aizawa, K. (1984). Purification of the toxic protein from *Bacillus thuringiensis* serotype 10 isolate demonstrating a preferential larvicidal activity to mosquito. *J. Invertebr. Pathol.* 44: 214-219.
- 9
- [29] Ohba, M. and Aizawa, K. (1986). Insect toxicity of *Bacillus thuringiensis* isolated from soils of Japan. *J. Invertebr. Pathol.* 47: 12-20.
- 17
- [30] Padua, L. E., Ohba, M. and Aizawa, K. (1984). Isolation **of** a *Bacillus thuringiensis* strain (serotype 8a:8b) highly and selectively toxic against mosquito larvae. *J. Invertebr. Pathol.* 44: 12-17.
- 29
- [31] Rejesus, B. M. and A. Sayaboc, (1990). Management of DBM with Apanteles plutellae prospects in the Philippines. Paper presented in the second International Workshop on Diamondback Moth Management. Dec. 10-15, 1990. Tainan, Taiwan. 17 pp.
- [32] Shorey, H. H. and I. M. Hall.(1962). Effect of chemical and microbial insecticides on several insect pests of lettuce in southern California. *J. Econ. Entomol.* 56: 169-174.
- 18
- [33] Yu, Y. -M., Ohba, M. and Gill, S. S (1991). Characterization of mosquitocidal activity of *Bacillus thuringiensis* subsp. *Fukuokaensis* crystal proteins. *Appl. Enviro. Microbiol.* 1075-1081.
- [34] Zhang, Y., Ku, Z., Chan, Z., Xu, B., Yuan, F., Chen, G., Zhong, T. and Ming, G. (1984). A new isolate of *Bacillus thuringiensis* possessing high toxicity against the mosquitoes. *Acta Microbiologica Sinica* 24: 320-325.

# Comparison Study Of Toxicity Bacillus thuringensis from Soil Isolate and Mulberry Leaves in Indonesia

ORIGINALITY REPORT



PRIMARY SOURCES

- |   |   |     |
|---|---|-----|
| 1 | Guang - Mao Shen, Shi - Yuan Ou, Chuan - Zhen Li, Kai - Yang Feng, Jin - Zhi Niu, Michael J. Adang, Lin He. "Transcription factors CncC and Maf connect the molecular network between pesticide resistance and resurgence of pest mites", Insect Science, 2021<br>Publication | 1 % |
| 2 | Submitted to Kaplan International Colleges<br>Student Paper   | 1 % |
| 3 | f1000research.com<br>Internet Source  | 1 % |
| 4 | Sainsbury, Frank, Meriem Benchabane, Marie-Claire Goulet, and Dominique Michaud. "Multimodal Protein Constructs for Herbivore Insect Control", Toxins, 2012.<br>Publication   | 1 % |
| 5 | HIMENO, Michio, Masao IKEDA, Kikuo SEN, Naoto KOYAMA, Tohru KOMANO, Haruyasu YAMAMOTO, and Isamu NAKAYAMA. "Plasmids and insecticidal activity of delta-  | 1 % |

endotoxin crystals from *Bacillus thuringiensis* var. *israelensis.*", Agricultural and Biological Chemistry, 1985.

Publication

6 Hernandez F.D., Castillo F., Gallegos G., Rodriguez R., C. Aguilar. "Chapter 4 Biological Control of Root Pathogens by Plant- Growth Promoting *Bacillus* spp.", IntechOpen, 2013

Publication

1 %

7 digitalcommons.unl.edu

Internet Source

1 %

8 Rachel E. Down. "Dissemination of the biocontrol agent *Vairimorpha necatrix* by the spined soldier bug, *Podisus maculiventris*", Entomologia Experimentalis et Applicata, 2/2004

Publication

1 %

9 T. A. Angus. "EXTRACTION, PURIFICATION, AND PROPERTIES OF *BACILLUS SOTTO* TOXIN", Canadian Journal of Microbiology, 1956

Publication

1 %

10 pesquisa.bvsalud.org

Internet Source

1 %

11 Chenguang Zhu. "Restraining *Erwinia* virulence by expression of N-acyl homoserine lactonase gene *pro3A-aiiA* in *Bacillus thuringiensis*

1 %

subspleesis", Biotechnology and  
Bioengineering, 10/20/2006

Publication

- 
- 12 Nikolai Galitsky, Vivian Cody, Andrzej Wojtczak, Debashis Ghosh, Joseph R. Luft, Walter Pangborn, Leigh English. " Structure of the insecticidal bacterial δ-endotoxin Cry3Bb1 of ", Acta Crystallographica Section D Biological Crystallography, 2001 1 %  
Publication
- 
- 13 S. Babashpour, S. Aminzadeh, N. Farrokhi, A. Karkhane, K. Haghbeen. "Characterization of a Chitinase (Chit62) from *Serratia marcescens* B4A and Its Efficacy as a Bioshield Against Plant Fungal Pathogens", Biochemical Genetics, 2012 1 %  
Publication
- 
- 14 H. Salama, M. Saker, M. Salama, A. El-Banna, M. Ragaie, N. Abd El-Ghany. " isolates from Egyptian soils and their potential activity against lepidopterous insects ", Archives Of Phytopathology And Plant Protection, 2012 1 %  
Publication
- 
- 15 Regis, L.. "Bacteriological larvicides of dipteran disease vectors", Trends in Parasitology, 20010801 <1 %  
Publication
- 
- 16 hec.gov.pk

- 
- 17 A. Abdel-Hameed, R. Landén. "Studies on *Bacillus thuringiensis* strains isolated from Swedish soils: insect toxicity and production of *B. cereus*-diarrhoeal-type enterotoxin", *World Journal of Microbiology & Biotechnology*, 1994 <1 %  
Publication
- 
- 18 A. Guerchicoff, A. Delecluse, C. P. Rubinstein. "The *Bacillus thuringiensis* cyt Genes for Hemolytic Endotoxins Constitute a Gene Family", *Applied and Environmental Microbiology*, 2001 <1 %  
Publication
- 
- 19 Ben Raymond, Kelly L. Wyres, Samuel K. Sheppard, Richard J. Ellis, Michael B. Bonsall. "Environmental Factors Determining the Epidemiology and Population Genetic Structure of the *Bacillus cereus* Group in the Field", *PLoS Pathogens*, 2010 <1 %  
Publication
- 
- 20 core.ac.uk <1 %  
Internet Source
- 
- 21 sciencepublishinggroup.com <1 %  
Internet Source
- 
- www.ijeat.org

22

Internet Source

<1 %

23

A. M.E. Abd El-Salam, Awad M. Nemat, Attia Magdy. "Potency of and against the cotton leafworm, (Bosid.) Larvae", Archives Of Phytopathology And Plant Protection, 2011

Publication

<1 %

24

D. Diallo, A. Marston, C. Terreaux, Y. Touré, B. Smestad Paulsen, K. Hostettmann. "Screening of Malian medicinal plants for antifungal, larvicidal, molluscicidal, antioxidant and radical scavenging activities", Phytotherapy Research, 2001

Publication

<1 %

25

Philipp Gerhardt, H. Stuart Pankratz, René Scherrer. "Fine Structure of the Spore", Applied and Environmental Microbiology, 1976

Publication

<1 %

26

Somerville, H.J.. "Lethal effect of purified spore and crystalline endotoxin preparations of Bacillus thuringiensis on several lepidopterous insects", Journal of Invertebrate Pathology, 197009

Publication

<1 %

27

[Www.intechopen.com](http://www.intechopen.com)

Internet Source

<1 %

- 28 H Hofstra, J D Van Tol, J Dankert. "Cross-reactivity of major outer membrane proteins of Enterobacteriaceae, studied by crossed immunoelectrophoresis", Journal of Bacteriology, 1980 <1 %  
Publication
- 
- 29 H. Katayama. "Parasporin-1, a Novel Cytotoxic Protein to Human Cells from Non-Insecticidal Parasporal Inclusions of *Bacillus thuringiensis*", Journal of Biochemistry, 01/01/2005 <1 %  
Publication
- 
- 30 Karim Ennouri, Saoussen Ben Khedher, Samir Jaoua, Nabil Zouari. " Correlation between delta-endotoxin and proteolytic activities produced by var. growing in an economic production medium ", Biocontrol Science and Technology, 2013 <1 %  
Publication
- 
- 31 L. Xu, H.-J. Huang, X. Zhou, C.-W. Liu, Y.-Y. Bao. " Pancreatic lipase-related protein 2 is essential for egg hatching in the brown planthopper, ", Insect Molecular Biology, 2017 <1 %  
Publication
- 
- 32 S. Edward Stevens, Randy C. Murphy, William J. Lamoreaux, Lewis B. Coons. "A genetically engineered mosquitocidal cyanobacterium", Journal of Applied Phycology, 1994 <1 %

33

[www.journal.csj.jp](http://www.journal.csj.jp)

Internet Source

---

<1 %

Exclude quotes Off

Exclude bibliography Off

Exclude matches Off