

Effect of Poultry Excreta on Water Quality and Daphnia Magna Production in Chlorella Powder Medium

by Fpk Ulm

Submission date: 08-Nov-2021 07:46PM (UTC+0700)

Submission ID: 1696622029

File name: try_Excreta_on_Water_Quality_and_Daphnia_Magna_Production_in.pdf (447.83K)

Word count: 4788

Character count: 19798

Open Access Article

Effect of Poultry Excreta on Water Quality and *Daphnia Magna* Production in Chlorella Powder Medium

H. Herliwati¹, M. Rahman¹, A.S. Hidayat¹, I. Sumantri^{2*}

¹ Faculty of Fisheries and Marine, University of Lambung Mangkurat, Banjarbaru, Indonesia

² Faculty of Agriculture, University of Lambung Mangkurat, Banjarbaru, Indonesia

Abstract: *Daphnia magna* Straus, 1820, plays a significant role in food supply for fish farming; thus, it is important to formulate a medium that can increase the *D. magna* population. This can be achieved by supplying a direct nutrient for *D. magna* while supporting the growth of the phytoplankton and zooplankton population. However, at certain levels, the nutrient source addition adversely affects the culture medium. This study aims to investigate the effects of different concentrations of poultry excreta included in a culture medium enriched with chlorella powder on the biomass production of *D. magna*. An experiment was conducted by adding poultry excreta at concentrations (0, 2, 4, and 6 g·L⁻¹) to the chlorella powder culture medium. The culture was maintained for 15 days, and samples were collected on days 0, 5, 10, and 15 to analyze the *D. magna* population, water pH, and concentrations of ammonia, nitrate, and dissolved oxygen (DO). Furthermore, a statistical evaluation was conducted using one-way analysis of variance in a completely randomized research design. The results showed that an increase in the poultry excreta concentration reduced the water quality ($P < 0.01$), as indicated by the water pH and ammonia, nitrate, and DO concentrations. In addition, a longer duration of the experiment substantially improved the qualitative parameters evaluated. Similarly, the population of *D. magna* was significantly affected ($P < 0.01$) by both factors. In conclusion, the addition of poultry excreta decreased the water quality of the chlorella powder medium. The water quality improved on prolonged days; thus, the highest population of *D. magna* was achieved on day 15 using 2 g·L⁻¹ of poultry excreta.

Keywords: *Daphnia Magna*, *Daphnia* production, Chlorella powder medium, poultry excreta, water quality.

家禽排泄物对小球藻粉培养基水质和大溞产量的影响

摘要: 大型水蚤, 1820年, 在鱼类养殖的食物供应中发挥着重要作用; 因此, 重要的是配制可以增加大麦格纳菌种群的培养基。这可以通过为D.麦格纳提供直接养分同时支持浮游植物和浮游动物种群的生长来实现。然而, 在某些水平上, 添加营养源会对培养基产生不利影响。本研究旨在研究富含小球藻粉的培养基中不同浓度的家禽排泄物对大麦格纳生物量产生的影响。通过向小球藻粉培养基中添加四种浓度(0、2、4和6g·升⁻¹)

的家禽排泄物进行了一项实验。培养物维持15天, 并在第0、5、10和15天收集样品以分析D.麦格纳种群、水的酸碱度值以及氨、硝酸盐和溶解氧(做)的浓度。此外, 在完全随机的研究设计中使用方差的单向分析进行了统计评估。结果表明, 家禽排泄物浓度的增加降低了水质(磷<0.01), 如水的酸碱度值和氨、硝酸盐和溶解氧浓度所示。此外, 更长的实验持续时间大大改善了评估的定性参数。同样, 这两个因素都显著影响了D.麦格纳的种群(磷<0.01)。总之, 家禽排泄物的添加降低了小球藻粉培养基的水质。水质在延长的日子里有所改善; 因此, 使用2g·升⁻¹的家禽排泄物在第15天实现了最大数量的D.麦格纳。

关键词: 大型水蚤, 水蚤产量, 小球藻粉培养基, 家禽排泄物, 水质。

Received: May 15, 2021 / Revised: June 6, 2021 / Accepted: July 9, 2021 / Published: August 30, 2021

Fund Project: PNPB University of Lambung Mangkurat

About the authors: Dr. H. Herliwati, M. Rahman, A.S. Hidayat, Faculty of Fisheries and Marine, University of Lambung Mangkurat, Banjarbaru, Indonesia; I. Sumantri, Faculty of Agriculture, University of Lambung Mangkurat, Banjarbaru, Indonesia

Corresponding author I. Sumantri, isumantri@ulm.ac.id

1. Introduction

Fish farming requires specific culture techniques as well as live food organisms to serve as feed, including zooplankton, which is naturally used in aquaculture. *Daphnia magna* Straus is considered an important zooplankton species in freshwater ecosystems, owing to its ability to control microbial populations and serve as a food supply for fish [1]. These cladocerans exhibit superior nutritional and physical properties in terms of particle size, digestibility, and essential nutrient content, which meet the requirements for the growth and reproduction of larvae and fish [2]. In addition, *Daphnia sp.* contains several enzymes, such as proteinases, peptidases, amylases, lipases, and cellulase, which can serve as digestive enzymes for fish larvae [3]. Therefore, the production of *Daphnia sp.* biomass has attracted considerable research interest over the decades, particularly in terms of their feeding techniques and nutrient requirements.

Some previous studies have indicated the effect of adding organic substrates, such as wheat bran, soybean powder [4], tofu waste [5], and animal wastes [6], to the culture medium. Animal wastes can supply nutrients for the proliferation of phytoplankton and bacteria, which serve as feed for *Daphnia sp.* However, certain levels of animal wastes in the culture medium tend to produce toxic effects on the reproduction of *Daphnia magna* Straus, 1820 [6].

This toxicity can be avoided by developing numerous culture media (e.g., chlorella powder) with the aim of directly supplying the relevant nutrients [4] [7], [8], especially essential amino acids and unsaturated fatty acids [9]. However, it is necessary to develop a medium that can adequately support the growth of both phytoplankton and zooplankton [10] [11]. Therefore, this study aims to determine the optimum concentration of poultry excreta that needs to be added to chlorella powder-enriched cultures to support the production of *D. magna*.

2. Materials and Methods

2.1. Materials

Daphnia magna Straus, 1820, was originally isolated from the river area of South Kalimantan, followed by its reservation and maintenance in the laboratory. Conversely, the poultry excreta were obtained from a local broiler farm, and the *Chlorella Sp.* powder was purchased from Ugoplanktonshop (Central Java, Indonesia).

2.2. Procedure for *D. Magna* Culture

An experiment was conducted by adding four concentrations (0, 2, 4, and 6 g·L⁻¹) of poultry excreta,

which was dried and ground through a 1-mm mill size. These samples were individually incorporated into a 60 × 40 × 30 cm³ chamber containing 20 L of tap water, with each level replicated three times. The experimental medium was aerated for eight days, followed by inoculation with 30 heads·L⁻¹ of *D. magna* or 600 heads per chamber; then, the chlorella powder was added at 1000 mg per day. Next, 500 mL of the water sample was collected on days 0, 5, 10, and 15 for determining the microorganism population and water quality, including pH, as well as the concentrations of ammonium, nitrate, and dissolved oxygen (DO).

2.3. Data Analysis

The data were statistically evaluated using the analysis of variance (one-way ANOVA) with a completely randomized research design. All statistical analyses were conducted using SPSS 22.0 software (IBM, USA).

3. Results

3.1. Water Quality

The concentration of poultry excreta added to the chlorella powder culture medium significantly ($P < 0.01$) affected the medium pH according to the *D. magna* culture period (Table 1). On day 0, the poultry excreta levels resulted in a relatively similar pH of culture media, ranging within a neutral pH (6.95–7.32). Then, higher amounts of poultry excreta increased the water pH water on day 5. However, on days 10 and 15, the medium without poultry excreta resulted in the highest pH value (8.39). Meanwhile, the media with poultry excreta addition had slightly neutral pH values, except for the poultry excreta at 6 g·L⁻¹ (pH 8.01).

Table 1 Effects of poultry excreta on the pH of chlorella powder culture medium

| Days | Excreta levels (g·L ⁻¹) | | | |
|------|-------------------------------------|------------------------|------------------------|------------------------|
| | 0 | 2 | 4 | 6 |
| 0 | 7.32±0.61 | 7.25±0.01 | 7.13±0.04 | 6.95±0.35 |
| 5 | 6.98±0.01 ^c | 6.08±0.01 ^a | 6.95±0.02 ^b | 7.21±0.01 ^d |
| 10 | 8.39±0.01 ^d | 7.20±0.01 ^a | 7.23±0.01 ^b | 8.01±0.01 ^c |
| 15 | 8.39±0.01 ^d | 7.20±0.01 ^a | 7.23±0.01 ^b | 8.01±0.01 ^c |

^{a, b, c, d} Means with different superscripts in the same row are significantly different ($P < 0.01$)

The amount of poultry excreta had a significant ($P < 0.01$) positive effect on ammonium concentration in the experimental medium, as shown in Table 2. Subsequently, a decline in the ammonia concentration was observed when extending the culture period.

Table 2 Effects of poultry excreta on ammonium concentration (mg.L^{-1}) in chlorella powder culture medium

| Days | Excreta levels (g.L^{-1}) | | | |
|------|--------------------------------------|------------------------|------------------------|------------------------|
| | 0 | 2 | 4 | 6 |
| 0 | 0.23±0.19 | 1.14±0.01 | 1.68±0.01 | 2.64±0.02 |
| 5 | 0.28±0.01 ^a | 0.64±0.10 ^b | 2.24±0.10 ^c | 0.52±0.01 ^b |
| 10 | 0.26±0.01 ^b | 0.19±0.01 ^a | 0.32±0.01 ^c | 0.37±0.01 ^d |
| 15 | 0.26±0.01 ^b | 0.19±0.01 ^a | 0.32±0.01 ^c | 0.37±0.01 ^d |

^{a, b, c, d} Means with different superscripts in the same row are significantly different ($P < 0.01$)

The concentration of poultry excreta conferred a significant effect on the concentration of nitrate present in the experimental medium ($P < 0.01$), as shown in Table 3. However, a decline in the concentration was observed during a prolonged period of exposure.

Table 3 Effects of poultry excreta on the nitrate concentration (mg.L^{-1}) of chlorella powder culture medium

| Days | Excreta levels (g.L^{-1}) | | | |
|------|--------------------------------------|------------------------|------------------------|------------------------|
| | 0 | 2 | 4 | 6 |
| 0 | 3.00±0.00 | 39.60±0.00 | 54.60±1.00 | 30.40±0.20 |
| 5 | 0.28±0.01 ^a | 0.64±0.10 ^b | 2.24±0.10 ^c | 0.52±0.10 ^b |
| 10 | 3.35±0.28 ^b | 4.80±0.10 ^c | 4.80±0.10 ^c | 0.10±0.00 ^a |
| 15 | 3.50±0.10 ^b | 4.80±0.10 ^c | 4.80±0.10 ^c | 0.10±0.00 ^a |

^{a, b, c} Means with different superscripts in the same row are significantly different ($P < 0.01$)

The DO content on day 5 was significantly affected by the concentration of poultry excreta ($P < 0.01$), although a reduction in the concentration was subsequently observed on days 10 and 15 ($P > 0.05$). This study showed increased DO concentrations after prolonged days of culture which were not significantly affected by the concentration of poultry excreta on day 10 and 15 (Table 4).

Table 4 Effects of poultry excreta on DO (mg.L^{-1}) of chlorella powder culture medium

| Days | Excreta levels (g.L^{-1}) | | | |
|------|--------------------------------------|------------------------|------------------------|------------------------|
| | 0 | 2 | 4 | 6 |
| 0 | 7.80±0.10 | 7.20±0.10 | 7.30±0.10 | 7.20±0.10 |
| 5 | 5.50±0.10 ^b | 7.70±0.10 ^a | 7.60±0.10 ^a | 4.20±0.10 ^c |
| 10 | 7.03±0.87 | 7.03±0.97 | 7.83±0.38 | 7.03±0.87 |
| 15 | 7.03±0.87 | 7.03±0.97 | 7.83±0.38 | 7.03±0.87 |

^{a, b, c} Means with different superscripts in the same row are significantly different ($P < 0.01$)

3.2. *D. Magna* Population

Table 5 shows the significant effect of poultry excreta concentration on the *D. magna* population on

day five ($P < 0.05$). This was very significant at day 10 ($P < 0.01$), and it was the reverse on day 15 ($P > 0.05$). Despite the relatively higher population in medium, without the treatment, a subsequent decline was observed after prolonged period of exposure. However, the presence of poultry excreta increased the *D. magna* population in the following 5 days, with the highest obtained on day 15, using a concentration of 2 g.L^{-1} .

Table 5 Effects of poultry excreta on *D. Magna* population (head.L⁻¹)

| Days | Excreta levels (g.L^{-1}) | | | |
|------|--------------------------------------|-----------------------|---------------------|----------------------|
| | 0 | 2 | 4 | 6 |
| 0 | 300±0 | 300±0 | 300±0 | 300±0 |
| 5 | 1,044±315 ^b | 255±93 ^a | 99±61 ^a | 865±245 ^b |
| 10 | 888±305 ^l | 611±224 ^{kl} | 362±30 ^k | 835±159 ^l |
| 15 | 876±208 | 1,488±917 | 836±378 | 782±34 |

^{a, b} Means with different superscripts in the same row are significantly different ($P < 0.01$)

^{k, l} Means with different superscripts in the same row are significantly different ($P < 0.05$)

4. Discussion

Daphnia is capable of tolerating a wide range of pH (6.5–9.5), although 7.2–8.5 was identified as the optimum value for the population growth of most species [1]. A study by [12] demonstrated 7.9–8.3 as an optimum pH range for the growth and reproduction of *D. magna*, as values lesser than 4.55 and more than 10.13 were assumed to decrease survival and reproduction.

This study showed a decline in the medium pH at the initiation of experiment, which was simultaneous with the increase of concentration of poultry excreta and an extended period of exposure. The responses were possibly linear or quadratic. However, the water pH was within a range that was suitable for *D. magna*, which is optimal for the commencement of growth and reproduction in experimental medium using poultry excreta concentration of 2 g.L^{-1} .

The ammonium content in water was produced from the decomposition of nitrogen compounds. Hence a higher supply results in the generation of more ammonium. This is known to confer detrimental and lethal effects on *Daphnia* at high amounts, according to [13], in a study that showed a decline in growth and reproduction after increasing the concentration from 0.366 to 0.581 mg.L^{-1} . Furthermore, the organic nitrogen compound used in this study was supplied by the chlorella powder and poultry excreta.

This study showed an increase in ammonium content of the experimental medium, which occurred as a response to the elevated poultry excreta concentration. This value decreased after a prolonged day of culture, reaching 0.37 mg.L^{-1} on day 15, with 6 g.L^{-1} of poultry excreta. Furthermore, the maximum

concentration required for *D. magna* to grow and reproduce was surpassed, based on the limit of $<0.2 \text{ mg.L}^{-1}$ [14]. Therefore, the addition at 2 and 4 g.L^{-1} treatment is expected to fulfil the nitrogen requirement for the growth and reproduction of *D. magna*.

Table 3 shows the presence of high nitrate content in the medium at the initiation of the experiment using poultry excreta, which subsequently decreased over a prolonged exposure period. One experiment reported 462 mg.L^{-1} as the lethal concentration for *D. magna* production; hence the values recorded in this study were below the toxic level [15]. However, the lowest quantity was identified in treatments with a 6 g.L^{-1} concentration.

The addition of poultry excreta was perceived to cause a decline in the DO content of the experimental medium. Low oxygen concentration (below 3.5 mg.L^{-1}) reduced growth rate, body size, and egg size of *Daphnia sp.* [16]. Smaller adult body sizes in hypoxia could be an adaptive response of *Daphnia sp.* to improve hypoxia tolerance.

The results showed higher concentrations after prolonged exposure, which were not significantly affected by poultry excreta on days 10 and 15 (Table 4). The decrease in DO on day five is possibly affiliated with the high microbial activity observed during the degradation of organic matter. As shown by [17], the decomposition of organic materials mediated by bacteria results in high respiration in the water. However, the DO content in this study was still in the optimum range required for the growth of *D. magna*.

Based on the initial water quality parameters, the experimental mediums with poultry excreta demonstrated worse outcomes than the control, especially with the ammonium (Table 2), nitrate (Table 3), and DO concentration (Table 4). These tend to decrease in content after five days, except for the DO, which increased. Low quality of food and environmental conditions contributed to reduced lifespan and fertility of *D. magna* [18]. *Daphnia sp.* is a sensitive and responsive species to water quality and environmental health [1]. Therefore, sources and levels of food inclusion in culture mediums should create a better environment for the growth and reproduction of *D. magna*.

5. Conclusion

Nutrient enrichment could change the water environment. This study indicated poultry excreta levels had a significant impact on pH, oxygen conditions, and other water quality parameters that determine *D. magna* growth and reproduction. Higher levels of poultry excreta caused a decline in the water quality of mediums enriched with chlorella powder. This was, however, in the tolerable range for the growth and reproduction of *D. magna*. In addition, the quality parameters were observed to have improved after prolonged days of culture, which was followed by

an increasing *D. magna* population, with the highest amount reported on day 15, using a poultry excreta concentration of 2 g.L^{-1} .

Acknowledgments

Authors would like to thank the University of Lambung Mangkurat for supporting this research through the PNPB ULM Fundamental Research grant.

References

- [1] DEMERTZIOGLOU M., ANTONOPOULOU E., VOUTSA D., KOZARI A., MOUSTAKA-GOUNI M., and MICHALOUDI E. MAPKs and HSPs' activation of a natural *Daphnia magna* population in a man-perturbed lake: implications of ecological significance. *Water*, 2021, 13(3): 283. <https://doi.org/10.3390/w13030283>
- [2] LAVENS P., & SORGLOOS P. Introduction. In: LAVENS P., & SORGLOOS P. (eds.) *Manual on the production and use of live food for aquaculture*. Food and Agriculture Organization, Rome, 1996.
- [3] RADHAKRISHNAN D. K., AKBARALI I., SCHMIDT B. V., JOHN E. M., SIVANPILLAI S., and VASUNAMBESAN S. T. Improvement of nutritional quality of live feed for aquaculture: an overview. *Aquaculture Research*, 2020, 51(1): 1-17. <https://doi.org/10.1111/are.14357>
- [4] EL-FEKY M. M., & ABO-TALEB H. Effect of feeding with different types of nutrients on intensive culture of the water flea, *Daphnia magna* Straus, 1820. *Egyptian Journal of Aquatic Biology and Fisheries*, 2020, 24(1): 655-666. <https://doi.org/10.21608/EJABF.2020.76554>
- [5] HERAWATI V. E., NUGROHO R. A., PINANDOYO, and HUTABARAT J. Nutritional value content, biomass production and growth performance of *Daphnia magna* cultured with different animal wastes resulted from probiotic bacteria fermentation. *IOP Conference Series: Earth and Environmental Science*, 2017, 55: 012004. <https://doi.org/10.1088/1755-1315/55/1/012004>
- [6] DAMLE D. K., & CHARI M. S. Performance evaluation of different animal residue on culture of *Daphnia sp.* *Journal of Fisheries and Aquatic Science*, 2011, 6(1): 57-61. <http://dx.doi.org/10.3923/jfas.2011.57.61>
- [7] ABDEL-TAWWAB M., ABDULRAHMAN N. M., BAIZ A. I., NADER P. J., and AL-REFAIEE I. H. A. The using of *Chlorella pyrenoidosa* and *Daphnia magna* as feed supplements for common carp, *Cyprinus carpio*: growth performance, somatic indices, and hemato-biochemical biomarkers. *Journal of Applied Aquaculture*, 2020. <https://doi.org/10.1080/10454438.2020.1787291>
- [8] MEHDIPOUR N., FALLAHI M. A., TAKAMI G., VOSSOUGH G., and MASHINCHIAN A. Freshwater green algae *Chlorella sp.* and *Scenedesmus obliquus* enriched with B group of vitamins can enhance fecundity of *Daphnia magna*. *Iranian Journal of Science and Technology*, 2011, 35(2): 157-163. <https://doi.org/10.22099/IJSTS.2011.2139>
- [9] MUNIRASU S., RAMASUBRAMANIAN V., UTHAYAKUMAR V., and MUTHUKUMAR S. Bioenrichment of live feed *Daphnia magna* for the survival and growth of freshwater fish *Catla catla*. *International Journal of Current Research and Review*, 2013, 5(8): 20-31. http://ijcrr.com/article_html.php?did=1371

[10] KILLHAM S. S., DANIEL A. K., LYNN S. G., GOULDEN C. E., and HERRERA L. COMBO: a defined freshwater culture medium for algae and zooplankton. *Hydrobiologia*, 1998, 377: 147-159. <https://doi.org/10.1023/A:1003231628456>

[11] CHEBAN L., GRYNKO O., and DOROSH I. Co-cultivation of *Daphnia magna* (Straus) and *Desmodesmus armatus* (chod.) Hegew. in recirculating aquaculture system wastewater. *Fisheries and Aquatic Life*, 2018, 26: 57-64. <https://doi.org/10.2478/aopf-2018-0007>

[12] GHAZY M. M. E., HABASHY M. M., and MOHAMMADY E. Y. Effects of pH on survival, growth and reproduction rates of the crustacean, *Daphnia magna*. *Australian Journal of Basic and Applied Sciences*, 2011, 5(11): 1-10. <http://ajbasweb.com/old/ajbas/2011/November-2011/1-10.pdf>

[13] YANG Z., LÜ K., CHEN Y., and MONTAGNES D. J. S. The Interactive Effects of Ammonia and Microcystin on Life-History Traits of the Cladoceran *Daphnia Magna*: Synergistic or Antagonistic? *PLoS ONE*, 2012, 7(3): e32285. <https://doi.org/10.1371/journal.pone.0032285>

[14] DELBARE D., DHERT P., and LAVENS P. Zooplankton. In: LAVENS P., & SORGLOOS P. (eds.) *Manual on the production and use of live food for aquaculture*. Food and Agriculture Organization, Rome, 1996.

[15] SCOTT G., & CRUNKILTON R. L. Acute and chronic toxicity of nitrate to fathead minnows (*Pimephales promelas*), *Ceriodaphnia dubia*, and *Daphnia magna*. *Environmental Toxicology and Chemistry*, 2000, 19(12): 2918-2922. <https://doi.org/10.1002/etc.5620191211>

[16] WILCZYNSKI W., DYNAK P., BABKIEWICZ E., BERNATOWICZ P., LENIOWSKI K., and MASZCZYK P. The combined effects of hypoxia and fish kairomones on several physiological and life history traits of *Daphnia*. *Fresh Water Biology*, 2019, 64(12): 2204-2220. <https://doi.org/10.1111/fwb.13407>

[17] GOCKE K., PINEDA J. E. M., and VALLEJO A. Heterotrophic microbial activity and organic matter degradation in coastal lagoons of Colombia. *Revista de Biología Tropical*, 2003, 51(1): 85-98. <https://revistas.ucr.ac.cr/index.php/rbt/article/view/15638>

[18] BOUCHNAK R., & STEINBERG C. E. W. Modulation of longevity in *Daphnia magna* by food quality and simultaneous exposure to dissolved humic substances. *Limnologia*, 2010, 40: 86-91. <https://doi.org/10.1016/j.limno.2009.11.010>

参考文献:

[1] DEMERTZIOGLOU M., ANTONOPOULOU E., VOUTSA D., KOZARI A., MOUSTAKA-GOUNI M. 和 MICHALOUDI E. 地图和热电偶在人为干扰的湖泊中激活天然水蚤种群：对生态的影响意义。水，2021，13(3)：283. <https://doi.org/10.3390/w13030283>

[2] LAVENS P., & SORGLOOS P. 介绍。载于：LAVENS P., & SORGLOOS P. (编辑。) 水产养殖活饵料生产和使用手册。粮食及农业组织，罗马，1996。

[3] RADHAKRISHNAN D. K., AKBARALI I., SCHMIDT B. V., JOHN E. M., SIVANPILLAI S., 和 VASUNAMBESAN S. T.

提高水产养殖活饲料的营养质量：概述。水产养殖研究，2020，51(1)：1-17. <https://doi.org/10.1111/are.14357>

[4] EL-FEKY MM, & ABO-TALEB H. 喂食不同类型营养物质对水蚤集约化养殖的影响，大型水蚤，1820. 埃及水生生物学和渔业杂志，2020，24(1): 655-666. <https://doi.org/10.21608/EJABF.2020.76554>

[5] HERAWATI V. E., NUGROHO R. A., PINANDOYO 和 HUTABARAT J. 用益生菌发酵产生的不同动物粪便培养的大蚤的营养价值含量、生物量生产和生长性能。眼压会议系列：地球与环境科学，2017，55：012004. <https://doi.org/10.1088/1755-1315/55/1/012004>

[6] DAMLE D. K., & CHARI M. S. 不同动物残留物对水蚤培养的性能评价。渔业与水产学杂志，2011，6(1)：57-61. <http://dx.doi.org/10.3923/jfas.2011.57.61>

[7] ABDEL-TAWWAB M., ABDULRAHMAN N. M., BAIZ A. I., NADER P. J., 和 AL-REFAIEE I. H. A. 使用蛋白核小球藻和大水蚤作为鲤鱼、鲤鱼的饲料添加剂：生长性能、体细胞指数和血生生物标志物。应用水产养殖杂志，2020. <https://doi.org/10.1080/10454438.2020.1787291>

[8] MEHDIPOUR N., FALLAHI M. A., TAKAMI G., VOSSOUGH G. 和 MASHINCHIAN A. 淡水绿藻小球藻富含维生素B群的斜生栅藻能增强大蚤的繁殖力。伊朗科学技术杂志，2011，35(2)：157-163. <https://doi.org/10.22099/IJSTS.2011.2139>

[9] MUNIRASU S., RAMASUBRAMANIAN V., UTHAYAKUMAR V. 和 MUTHUKUMAR S. 活饲料水蚤的生物富集，用于淡水鱼卡特拉卡特拉的生存和生长。国际当代研究与评论杂志，2013，5(8)：20-31. http://ijcr.com/article_html.php?did=1371

[10] KILLHAM S. S., DANIEL A. K., LYNN S. G., GOULDEN C. E. 和 HERRERA L. 组合：用于藻类和浮游动物的特定淡水培养基。水生物学，1998，377：147-159. <https://doi.org/10.1023/A:1003231628456>

[11] CHEBAN L., GRYNKO O. 和 DOROSH I. 大型水蚤(施特劳斯)和铁线莲(乔德)黑格共同培养。在循环水产养殖系统废水中。渔业和水生生物，2018，26：57-64. <https://doi.org/10.2478/aopf-2018-0007>

[12] GHAZY M. M. E., HABASHY M. M. 和 MOHAMMADY E. Y. 酸碱度值对甲壳类动物大型水蚤的存活、生长和繁殖率的影响。澳大利亚基础与应用科学杂志，2011，5(11)：1-10. <http://ajbasweb.com/old/ajbas/2011/November-2011/1-10.pdf>

[13] YANG Z., LÜ K., CHEN Y., 和 MONTAGNES D. J. S. 氨和微囊藻毒素对枝角水蚤生活史特征的相互作用：协同还是拮抗？公共图书馆一号，2012，7(3)：e32285. <https://doi.org/10.1371/journal.pone.0032285>

[14] DELBARE D., DHERT P. 和 LAVENS P. 浮游动物。载于：LAVENS P., & SORGLOOS P. (编辑。) 水产养殖活饵料生产和使用手册。粮食及农业组织，罗马，1996。

-
- [15] SCOTT G., & CRUNKILTON R. L. 硝酸盐对黑头鲈鱼(斑马目普罗梅拉斯)、银耳木虱和大型水蚤的急性和慢性毒性。环境毒理学与化学, 2000, 19(12): 2918-2922。 <https://doi.org/10.1002/etc.5620191211>
- [16] WILCZYNSKI W., DYNAK P., BABKIEWICZ E., BERNATOWICZ P., LENIOWSKI K., 和 MASZCZYK P. 缺氧和鱼类开罗酮对水蚤的几个生理和生活史特征的综合影响。淡水生物学, 2019, 64(12): 2204-2220。 <https://doi.org/10.1111/fwb.13407>
- [17] GOCKE K., PINEDA J. E. M. 和 VALLEJO A. 哥伦比亚沿海泻湖的异养微生物活动和有机物降解。热带生物学杂志, 2003, 51(1) : 85-98。 <https://revistas.ucr.ac.cr/index.php/rbt/article/view/15638>
- [18] BOUCHNAK R., & STEINBERG C. E. W. 通过食品质量和同时暴露于溶解的腐殖质物质对大蚤长寿的调节。湖沼学, 2010, 40 : 86-91。 <https://doi.org/10.1016/j.limno.2009.11.010>

Effect of Poultry Excreta on Water Quality and Daphnia Magna Production in Chlorella Powder Medium

ORIGINALITY REPORT

14%

SIMILARITY INDEX

11%

INTERNET SOURCES

8%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|---|---|----|
| 1 | Submitted to Universitas Negeri Manado Student Paper | 2% |
| 2 | www.mdpi.com Internet Source | 1% |
| 3 | e-sciencecentral.org Internet Source | 1% |
| 4 | www.science.gov Internet Source | 1% |
| 5 | www.tandfonline.com Internet Source | 1% |
| 6 | Bu-Min Kim, Yu-Na Jang, Sun Moon Kang, Yunseok Kim, Eun-Seon Lee, Jong-Hui Kim. "Analysis of quality changes and generation of sulfur volatiles according to the storage period of chicken", Korean Journal of Food Preservation, 2021 Publication | 1% |
| 7 | Divya Kandathil Radhakrishnan, Isamma AkbarAli, Bjorn V. Schmidt, Elizabeth Mary | 1% |

John et al. "Improvement of nutritional quality of live feed for aquaculture: An overview", Aquaculture Research, 2019

Publication

8

Bogdan Drugă, Patrick Turko, Piet Spaak, Francesco Pomati. "Cyanobacteria Affect Fitness and Genetic Structure of Experimental Populations", Environmental Science & Technology, 2016

Publication

<1 %

9

pure.rug.nl

Internet Source

<1 %

10

unctad.org

Internet Source

<1 %

11

H Heryani, N R Yanti. "Potentials of biomass waste sources for heterogeneous catalyst production", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

12

Paula Andrea Conforti, Mariela Patrignani. "Increase in the antioxidant content in biscuits by infusions or pod flour", Open Agriculture, 2021

Publication

<1 %

13

journal.yykxjz.cn

Internet Source

<1 %

14 Dewi Erika Adriani, Michael Dingkuhn, Audrey Dardou, H el ene Adam, Delphine Luquet, Tanguy Lafarge. "Rice panicle plasticity in Near Isogenic Lines carrying a QTL for larger panicle is genotype and environment dependent", Rice, 2016
Publication <1 %

15 Qiao-Guo Tan, Wen-Xiong Wang. " Acute Toxicity of Cadmium in under Different Calcium and pH Conditions: Importance of Influx Rate ", Environmental Science & Technology, 2011
Publication <1 %

16 Sarah K. Wise, Sandra Y. Lin, Elina Toskala, Richard R. Orlandi et al. "国际过敏与鼻科学共识声明：变应性鼻炎", International Forum of Allergy & Rhinology, 2019
Publication <1 %

17 www.ncbi.nlm.nih.gov
Internet Source <1 %

18 www.pesca.sp.gov.br
Internet Source <1 %

19 xbbjb.swu.cn
Internet Source <1 %

20 en.bio-protocol.org
Internet Source <1 %

21 hdl.handle.net Internet Source <1 %

22 www.ecologica.cn Internet Source <1 %

23 www.esap-ethiopia.org Internet Source <1 %

24 www.jlakes.org Internet Source <1 %

25 Smith Jean. "Ethanol Decreases Rat Hepatic Arylsulfatase A Activity Levels", Alcoholism Clinical and Experimental Research, 11/2006
Publication <1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On