

Search all conversations ☰ Active ? ⚙️ ☰  



← 📅 🕒 🗑️ ✉️ 🕒 🔄 📁 🗨️ ⋮ 369 of 415 < >

Fwd: Saya sedang berbagi 'Artikel Jurnal Bambang P\_130220.docx' dengan Anda Inbox x ⌵ 🖨️ 📄

📧 Feb 17, 2020, 9:51 AM ★ ↶ ⋮

🌐 Indonesian > English Translate message Turn off for: Indonesian x

----- Forwarded message -----  
Dari: **Benny Pratasik** <spjong07@yahoo.com>  
Date: Sen, 17 Feb 2020 08:36  
Subject: Re: Saya sedang berbagi 'Artikel Jurnal Bambang P\_130220.docx' dengan Anda


Search all conversations ☰ Active ? ⚙️ ☰  


← 📅 🕒 🗑️ ✉️ 🕒 🔄 📁 🗨️ ⋮ 256 of 415 < >

📧 Feb 29, 2020, 12:36 PM ★ ↶ ⋮

Dear Dr. Gavriolaie,  
I have rechecked the whole document of Poerwanto et al's final form and made some minor corrections (yellow). Now, it is ready for uploading it. Thank you

Sincerely,





## The analysis of factors influencing carp (*Cyprinus carpio*) seed production in Bungur District, Tapin Regency, South Kalimantan

<sup>1</sup>Bambang Poerwanto, <sup>2</sup>Idianoor Mahyudin, <sup>2</sup>Leila A. Sofia

<sup>1</sup> Student of Magister Program of Fisheries Science, Lambung Mangkurat University, South Kalimantan, Indonesia; <sup>2</sup> Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, South Kalimantan, Indonesia.  
Corresponding author: L. A. Sofia, leila.ariyani@ulm.ac.id

**Abstract.** Seed production has become one of the factors supporting the continuity of fish farming activity. Carp, *Cyprinus carpio* is one of the largest freshwater fish productions that support the human food needs in the world. This study aims to know factors affecting carp seed production. The study locality was purposively selected in Bungur District, Tapin Regency, South Kalimantan, and samples were collected through census from 42 carp farmers. The primary data were descriptively quantitatively analyzed using multiple regression. Results showed that there were 4 factors affecting the seed production, number of male spawners, feed amount for fry, spawning frequency, and the use frequency of the same female spawners. Carp seed production development could be done by considering supply of enough number and good quality spawners, good protein feeding and efficient feeding technique, and spawning recovery time of the female spawners.

**Key Words:** fish farmer, pond, stocking density, spawners, feeding.

**Introduction.** World human population in 2016 reached 7.4 billion people. This number has increased as many as 400 million people from that in 2011 (United Nations 2019). Increase in human population causes increase in world food needs including fisheries products. In 2016, the world fisheries production also reached 170.9 million tons, from which 151.2 million tons were used for human consumption and 19.7 million tons for non consumption need. This production comprises fishing fisheries and aquaculture. Nevertheless, the world fishing fisheries production has declined from 2011 to 2016 from 92.2 to 90.9 million tons, while world aquacultural production has significantly risen from 61.8 to 80.0 million tons in the same period (FAO 2018). Systematic fishing in tremendously high numbers, weak reporting, and other complex pressures added with ecosystem change consequences have resulted in decline in fishing fisheries production (Allan et al 2005). Hence, fish farming becomes a very important sector that plays important roles in world food security due to increasing human population and food needs (Wang et al 2017).

Indonesia as the fourth highest human population country, 267.6 million people (United Nations 2019) required fish consumption up to 43.94 kg capita<sup>-1</sup> in 2016. This need was supported by 16.68 million tons of aquaculture and 6.83 million tons of fishing fisheries. Total fisheries production of Indonesia in 2016 reached 23.51 million tons and has contributed to 13.75% of world fisheries production (Ministry of Marine Affairs and Fisheries (MMAF) of RI 2018a). The need of Indonesian fisheries products can be achieved by 70% from aquaculture. This condition makes aquaculture be very crucial. There are several fish commodities developed in Indonesia, one of which is carp, *Cyprinus carpio*. This species becomes the fourth largest freshwater aquacultural commodity in Indonesia with a volume of 316,646 tons in 2017 (MMAF RI 2018b). This species also became the third largest fish culture commodity in the world (Xu et al 2012; FAO 2018) and one of the fish species that can reach high biomass and its distribution

**Commented [indra1]:** provide the affiliation only, remove the position

**Commented [indra2]:** add the city as well

**Commented [indra3]:** city to be added

**Commented [indra4]:** all the factors or only some of them?

**Commented [indra5]:** can you update the info for 2020? Or, at least for 2019?

has been developed by human due to its important economic value (Arteni & Rosca 2010; Saikia & Das 2009).

The magnitude of carp culture commodity needs fish seed availability as important prerequisite of carp cultivation (Bhuyan et al 2012). The supply of good quality fish seeds is necessary for faster development of aquacultural industry. Hatchery efforts are also needed to reduce the dependence of aquaculture upon the use of wild seeds (Ghosh et al 2019). The development of carp seed quality has been conducted through various techniques, such as application of certain treatment to carp seeding production factors. The treatment could be implemented by previously knowing which production factor yields significant effect on the seeding quantity and quality. Studies on the effect of various factors on carp condition have been done a lot, such as how the intestinal microbes affect the fish growth (Li et al 2013) up to how the toxin effect, such as cobalt chloride, on the hematological condition of carp (Saravi et al 2009). Nevertheless, there are very few studies on carp seeding quality and quantity development. Nowadays, the development of good fish seed quality production is needed that it can help the cultured seed requirement and be practiced as a profitable business (Ismi et al 2014).

Tapin Regency, South Kalimantan province, is one of the regions that run carp culture activities with production level up to 16,235,000 individuals. Bungur District is one of the largest carp seeding localities in Tapin Regency. The seed production must be raised due to increasing human population and food demand, fisheries product in particular. Increased fish seed production is expected to be able to maintain the sustainability of the carp culture industry. Developing the carp seeding production requires a right strategy. This study aims to know factors affecting the carp seed production in Bungur District, Tapin Regency, the province of South Kalimantan.

## Material and Method

**Study site description.** The work was carried out from January to May 2019 in Bungur District, Tapin Regency, South Kalimantan province with an area of 148.98 km<sup>2</sup>. Bungur District is the largest aquacultural area in Tapin Regency, where 75% is located in Bungur District, and most of it is used for fish hatchery business, including carp seeding (Statistic Center of Tapin Regency 2019). This study site selection was done since Bungur District is the fish seeding center of Tapin Regency with the biggest aquacultural area and number of fish farmers.

**Population and samples.** Fish hatchery farmers in Shabah village and Kalumpang village, Bungur District are 42 people. All population members were taken as research samples through census method.

**Data analyses.** Factors influencing carp seed production were analyzed using descriptively quantitative method and OLS multiple regression with Cobb-Douglas-typed production function approach (Soekartawi 1994) as follows:

$$Y = a X_1^{b_1} X_2^{b_2} X_3^{b_3} X_4^{b_4} X_5^{b_5} X_6^{b_6} X_7^{b_7} X_8^{b_8} X_9^{b_9} X_{10}^{b_{10}} D_1 e$$

$$\ln Y = \ln a + b_1 \ln X_1 + b_2 \ln X_2 + b_3 \ln X_3 + b_4 \ln X_4 + b_5 \ln X_5 + b_6 \ln X_6 + b_7 \ln X_7 + b_8 \ln X_8 + b_9 \ln X_9 + b_{10} \ln X_{10} + D_1 + e$$

where a is constant, b<sub>1</sub>-b<sub>10</sub> are variable coefficient, Y is the fish seeding production (ind year<sup>-1</sup>); X<sub>1</sub> is pond size (m<sup>2</sup>), X<sub>2</sub> is number of female spawners (ind), X<sub>3</sub> is number of males (ind), X<sub>4</sub> is feed volume for spawners (kg), X<sub>5</sub> is total feed volume for fry (kg), X<sub>6</sub> is amount of manure fertilizer (pack), X<sub>7</sub> is spawning frequency (times), X<sub>8</sub> is use frequency of the same female spawners, X<sub>9</sub> is number of labor used (HOK), X<sub>10</sub> is fish farmer's experience (year), D<sub>1</sub> (dummy variable) is trained fish farmer = 1, untrained fish farmer = 0, and e is error term.

Commented [indra6]: to be supported by references

Commented [indra7]: make it more explicit

Commented [indra8]: for what period of time? Each year?

Commented [indra9]: also add some info about the statistical analysis since you refer to it in one of the tables

## Results

**Seeding activity description.** The Public Hatchery Unit has been done by the farming group in Bungur District for 8 years in average. The hatchery business run for 1-5 years was 48%, 6-10 years was 24%, and more than 10 years was 29%. The construction of carp seeding pond used by fish farmers in Bungur District is soil ponds with area around 1,490-18,212 m<sup>2</sup> and number of ponds managed were 4-29 units hatchery<sup>-1</sup> (Table 1) with total labors of 12-306 working days. Number of labors in each hatchery was 2-3 people depending upon number of spawning ponds operated. The pond localities are generally near the irrigation route so that the ponds are close enough to the water source. Before use, the fish farmers have fertilized the pond with manure. The organic fertilizer is effective to grow natural food that the productivity and fish growth can be increased (Kour et al 2016). The fertilizer applied in the study site was about 160-2,000 bags yr<sup>-1</sup> hatchery<sup>-1</sup> or about 14 bags m<sup>-2</sup>.

Fish spawner is major production factor in seed production, in either quality or quantity. Both females and males were prepared by the fish farmers. The female spawner candidates were 1.5-3 years old with individual weight of about 2-4 kg, the males were older than 6 months old with an individual weight of 0.2-1 kg. In general, number of spawners prepared by the fish farmers were 4-53 females and 14-178 males per business unit. The female and male spawners ratio was 2:8 each spawning.

Table 1  
Carp hatchery production and input production description in Public Hatchery Unit in Bungur district

Description	Unit	Value		
		Maximum	Minimum	Average
Fish seeding production	ind yr <sup>-1</sup>	2,080,000	100,000	340,520
Total pond size	m <sup>2</sup>	18,212	1,490	6,874
Number of ponds	unit	29	4	9
Pond size	m <sup>2</sup> unit <sup>-1</sup>	1,197	298	782
Number of female spawners	ind	53	4	16
Weight of female spawners	kg ind <sup>-1</sup>	4	2	3
Number of male spawners	ind	178	14	61
Weight of male spawners	kg ind <sup>-1</sup>	1	0.2	0.34
Feed volume of spawners	kg yr <sup>-1</sup>	840	40	318
Feed volume for fish fry	kg yr <sup>-1</sup>	1,560	200	647
Volume of manure fertilizer	bags yr <sup>-1</sup>	2,000	160	651
Spawning frequency	no yr <sup>-1</sup>	32	4	7
Use frequency of the same female spawners	no yr <sup>-1</sup>	6	2	4
Number of labors used	working days	306	12	83
Fish farmer's experience	year	18	1	8

Total time utilized for spawning in one-year is about 8 months. One production cycle requires more or less 60 days, and it could be done 1-2 time mo<sup>-1</sup> or 4-32 times yr<sup>-1</sup>. To maintain the quality of the produced seed, the frequency use of the same female spawners was 2-6 times yr<sup>-1</sup>, and every year the new spawners are prepared to replace the old ones. Total number of seeds produced by the hatchery were 100,000-2,080,000 ind yr<sup>-1</sup>.

The seeds were then reared in the nursery ponds from larval stage to selling size, 2-3 cm, 3-5 cm, and 5-8 cm long that needed about 2 months. In this stage, the seeds were given fine pellet with total feed weight of 200-1560 kg y<sup>-1</sup> per business unit. Similarly, the spawners were fed with artificial feed as much as 40-840 kg yr<sup>-1</sup> per business unit.

**Financial analysis.** Understanding the business profit level is important for the fish farmers to make decision in relation with the production planning. Carp hatchery

Commented [indra10]: why three classes of lengths?

Commented [indra11]: which length: standard, total or fork?

Commented [C12]: These are the fry size ranges that are commonly prepared by the hatchery to the fish farmers under different selling price

Commented [indra13]: to be rephrased, it has no meaning, because the verb is missing

business needs an investment of IDR 2.9 million – IDR 254.23 million (Table 2) depending upon land size, number of ponds used, and other supporting facilities, such as keeping house and storage, net/happa, kakaban/egg substrate, bucket, sorting box, and scoop net. Each business unit required the operational costs of IDR 29.69 million – IDR 159.34 million yr<sup>-1</sup> with the highest in feed preparation for the seeds, IDR 1.05 million – IDR 94 million yr<sup>-1</sup> depending upon the seed numbers produced. The additional feed was administered at the life stage of about 7-day old larvae (with yolk sac) up to be ready for sale.

The second highest operational cost is labor’s wages, IDR 16.4 million – IDR 27 million. It was allocated for pond preparation or processing, dyke improvement, and feeding. The lowest cost was recorded in male spawner supply, IDR 135,000 – IDR 1,750,000 yr<sup>-1</sup>. Previous studies found no enough difference, with total costs of IDR. 66,262,510.89, approximately 31% is use for feed costs (Shivakumar et al 2014). This finding is in agreement with Kholifah et al (2012) that carp hatchery business has good prospectus with a profit of IDR. 7,329,301.90 cycle<sup>-1</sup>. Different from that of Yulinda (2012) concerning the business financial analysis of catfish *Clarias gariepinus* with much lower profit, averagely IDR. 1,745,194 cycle<sup>-1</sup>. It could result from higher market price of carp fry than that of catfish.

Table 2  
Investment, operational costs, and profit of carp hatchery unit in Bungur District

Description	Unit	Value		
		Maximum	Minimum	Mean
Investment (total):	IDR	<b>254,225,000</b>	<b>2,935,000</b>	<b>46,256,475</b>
Pond	IDR	200,000,000	1,800,000	32,222,500
Other supporting facilities	IDR	54,225,000	1,135,000	14,033,975
Operational costs (total):	IDR yr <sup>-1</sup>	<b>159,340,000</b>	<b>29,692,500</b>	<b>61,318,504</b>
Female spawner preparation	IDR yr <sup>-1</sup>	7,500,000	480,000	2,139,500
Male spawner preparation	IDR yr <sup>-1</sup>	1,750,000	135,000	621,038
Spawner’s feed preparation	IDR yr <sup>-1</sup>	22,800,000	900,000	6,181,167
Seed feed preparation	IDR yr <sup>-1</sup>	94,000,000	1,050,000	19,699,750
Manure fertilizer	IDR yr <sup>-1</sup>	33,500,000	2,680,000	10,445,300
Other input preparation	IDR yr <sup>-1</sup>	14,750,000	350,000	3,081,750
Labor’s wages	IDR yr <sup>-1</sup>	27,300,000	16,400,000	19,150,000
Revenue and profit:				
Revenue	IDR yr <sup>-1</sup>	268,000,000	57,600,000	120,215,750
Business profit	IDR yr <sup>-1</sup>	177,795,000	15,370,000	58,897,246

Commented [indra14]: check again the calculations, because these totals here are not correct. For example, for the first column the total exceeds 200 million IDR

The revenue (from seed sales) obtained by a hatchery business unit ranged from IDR 57.6 million to IDR 268 million. The amount of revenue is dependent upon the number of production, seed size, and selling price. The larger the seed size is, the more expensive the seed will be. Under an agreement among the fish farmers, the selling price was IDR 250 ind<sup>-1</sup> for 2-3 cm long seed, IDR 300 ind<sup>-1</sup> for 3-5 cm length seed, and IDR 350 ind<sup>-1</sup> for 5-8 cm length seed, respectively. Therefore, the business profit varied as well from IDR 15.37 million to IDR 547.34 million.yr-1 or IDR 1.28 million to IDR 45.61 million mo<sup>-1</sup>.

Commented [indra15]: we cannot see this value in the table...

**Production and production factors.** The effect of production factors on the carp seed production showed that several production factors (input) involved in production function model gave the consistent values with the theoretical estimate. The production factors of positive value and significant effect on production were number of male spawners, feed volume for seeds, and spawning frequency done by the fish farmers. However, the use of the same female spawners for several spawning activities during the reproductive cycle will cause seed production decline as well (Table 3).

Table 3

Factors influencing the production of common carp hatchery

Variable	Coefficient	SE	t-Statistic	P value
Intercept	7.543	1.264	5.945	0.000
Total pond size (m <sup>2</sup> )	0.059	0.123	0.477	0.637
No. female spawners (ind.)	0.065	0.106	0.620	0.540
No. male spawners (ind.)	0.343	0.116	2.951**	0.006
Feed volume for spawners (kg)	-0.071	0.088	-0.813	0.423
Feed volume for seeds (kg)	0.381	0.138	2.760**	0.010
No. manure fertilizer (bag)	0.094	0.103	0.920	0.365
Spawning frequency (times)	0.910	0.110	8.241**	0.000
Use frequency of the same female spawners (times)	-0.324	0.139	-2.338**	0.026
Labor's working days	-0.204	0.113	-1.802	0.082
Fish farmer's experience (year)	-0.048	0.079	-0.608	0.548
Trained fish farmers (dummy)	0.112	0.139	0.808	0.425
R-squared		0.828		
Adjusted R-squared		0.765		
Standard error		0.321		
Observations		42		
F-statistic		13.154		
Prob (F-statistic)		0.000		
Durbin-Watson		2.190		

Commented [indra16]: you need to say few words about these indices in the Material and Method section - Data analyses

Hence, to increase the seed production, the fish farmers could increase several inputs and reduce the use frequency of the same female spawners to give them some time to recover and to have perfect maturation.

**Discussion.** Fish gamete quality, male or female, is determined by a number of factors, such as age, management, food, chemical and physical factors, water quality, and so on, that will impact the survivorship of the embryos, larvae, and/or fry in short or long term (Valdebenito et al 2013). Fish spawners are a major production factor in the hatchery business. The use of higher number of spawners in spawning activity will positively influence the number of eggs produced. The more the spawners are spawned, the higher the spawning frequencies could be done and the higher the number of eggs will be produced (Syaifudin et al 2007). The female carp spawner appropriate for spawning should be 1.5-3 years old with minimum weight of 1.5 kg ind<sup>-1</sup>, while the male should be at least 6 months old with minimum weight of 0.5 kg ind<sup>-1</sup> (Mantau et al 2004; Masoumeh et al 2013). The weight ratio of female and male is 1:1 (kg m<sup>-2</sup>) meaning that a 2 kg ind<sup>-1</sup> female could be balanced by 3 males of 600-700 g ind<sup>-1</sup>.

Some fish hatchery businesses showed that female-male ratio became factor determining the quality and quantity of seeds produced. For instance, in rainbow fish spawning, it is known that the use of sex ratio with 1:3 of total weight is more efficient than 1:1 and 1:2 ratios. It is possible that more male spawners will give higher opportunity to fertilize the eggs. Also, the seed quality produced at the spawner sex ratios of 1:2 and 1:3 gave 5-day old seeds with good survival rate (Herjayanto et al 2016). The climbing perch *Anabas testudineus* at 1:4 ratio spawners yields good fecundity (Burmansyah et al 2013). However, the female-male ratio practiced by the fish farmers in the study site indicated that total weight of male spawner was still below the total weight of female spawner. Thus, number and weight of male spawners need to be added in order to increase the seed production.

*C. carpio* naturally showed 2 clear reproductive cycle per year. Each reproductive cycle is divided into 4 phases. Males have growth phase, maturation, spawning, and resting, while females have preparatory phase, pre-spawning, spawning, and post-spawning (Guha & Mukherjee 1991). At 2 years of age, carp has fecundity of 523,500

Commented [indra17]: provide the scientific name too

eggs and at 3 years of age, the fecundity reaches 657,701 eggs (Aliniya et al 2013). Thus, the more the number of spawners spawned are, the more the number of eggs and eventually more larvae are produced. The availability of spawners spawned will also interchangeably give the opportunity to the spawners to have perfect reproductive cycle. The spawning frequency using the same female spawners gave impact on the seed growth. This study found that the seeds obtained from the spawners often used had lower body mass and shorter size. Also, the larvae produced from the often used spawners looked paler than those from the less used spawners.

Commercial feasibility of intensive fish culture depends on market demand and production costs. The largest part of the production costs is feed, so that it is important to know the optimal feeding and the right feed management strategy implementation (Jia et al 2016). The determination of different feeding strategy will yield different cultural conditions. For instance, the right feeding frequency can reduce the production costs and prevent water quality degradation (Daudpota et al 2016). Higher feeding frequency gives positive impact on the fish growth and makes the feed conversion ratio value be more efficient (Ganzon-Naret 2013a). Besides that, the percent of feeding becomes important factor for fish growth and the percentage of optimal feeding determines the aquaculture success (Deyab & Hussein 2015; Aryani et al 2017). The percent feed amount given will influence the fish growth rate as weight increment (Ganzon-Naret 2013b; Sofia et al 2013).

Feed is an important component for the spawners in order to produce good quality eggs and seeds in high number. The fish spawners spawned at the high frequency need good quality feed. Specific additional feeding also affects the aquacultural production. For example, addition of tauge extract in gold fish *Carrasius auratus* feed gives faster spawning cycle (Fajrin et al 2012). Addition of probiotic to the feed can increase body weight and fish endurance, so that the production rises and feed costs decline (Aitkaliyeva et al 2019). Besides feeding technique, the fish farmers should understand how to store the feed before use. According to Solomon et al (2016), storage in closed condition can prevent the nutritional damage of the feed.

Carp seeds produced need different feed treatments with life stages divided into 3 nursing phases. **First**, this stage was done for 7-day old larvae (0.6-0.7 cm long) at a stocking density of 100 ind m<sup>-2</sup> and in a pond of about 60 cm deep. At this stage, they were fed with the aquatic biota living in the pond and fine pellet. In about 15 days of rearing, the fish grew to 1-3 cm long. **Second**, this stage was conducted at the stocking density was about 50 ind m<sup>-2</sup> in about 20 days of rearing up to producing 3-5 cm long fish. **Third**, the stocking density of this stage was about 25 ind m<sup>-2</sup> at rearing period of about 30 days until 5-8 cm long-fish were obtained. After these nursing phases, the seeds are ready to sell, and they are sold when reaching the individual size of 3-5 cm and 5-8 cm long.

So far, feeding in each nursing phase still relied upon the artificial feed so that the operational cost for the fry feed was very high. On the other hand, the economic return of the larvae produced in the hatchery is more important than the fry growth. Therefore, alternative additional feed and feed cost efficiency need to be applied in order to increase the survivorship. One of these is tubifex supplement. According to Rawat et al (2019), 5% tubifex supplement can be used to increase the fish survivorship, especially in indoor seed rearing.

**Conclusions.** There were 4 factors affecting the production of *C. carpio* hatchery, three of which had positive effect on seed production, namely the number of male spawners, the amount of feed for the fry, and the spawning frequency. While the use frequency of female spawners during the production cycle negatively impact the seed production. This finding suggested that the fish farmers need to pay attention on the availability of sufficient number of good fish spawners, good protein feeding, efficient feeding technique, and longer post-spawning recovery time for female spawners.

**Acknowledgements.** We would greatly appreciate the Animal Husbandry and Fisheries Services of Tapin Regency, the Fisheries Consultant for field information and guidance in

Commented [indra18]: make it more clear

Commented [C19]: Erased

Commented [indra20]: add the Latin name as well

this study. We would also thank the fish farmers in the Public Hatchery Unit in Bungur District, Tapin Regency for their collaboration.

## References

- Aitkaliyeva A.A., Alpeisov S.A., Isbekov K.B., Assylbekova S.Z., Badryzlova N.S., 2019 Influence of different feeds and feed additives on fish-breeding and biological indicators at rearing rainbow trout. *EurAsian Journal of BioSciences* 13(1): 437-442.
- Aliniya, M., Khara, H., Noveiri, S.B., & Dadras, H., 2013 Influence of age of common carp (*Cyprinus carpio*) broodstock on reproductive traits and fertilization. *Turkish Journal of Fisheries and Aquatic Sciences* 13: 19-25.
- Allan J. D., Abeli R., Hogan Z., Revenga C., Taylor B. W., Welcomme R.L., Winemiller K., 2005 Overfishing of inland waters. *BioScience* 55 (12): 1041-1051.
- Arteni M. O., Rosca I., 2010 Some morphological feature of carp (*Cyprinus carpio* Linnaeus, 1758) in different stages of development. *Animal Biology and Animal Husbandry Bioflux* 2 (1): 47 -52.
- Aryani N., Azrita, Mardiah A., Syandri H., 2017 Influence of feeding rate on the growth, feed efficiency and carcass composition of the giant gourami (*Osphronemus goramy*). *Pakistan Journal Zoology* 49(5): 1775-1781.
- Bhuyan A., Akther S., Aktar N., 2012 Present status and fish seed production of the hatcheries of six upazillas of Rajshahi District. *University Journal of Zoology, Rajshahi University* 30: 29-32.
- Burmansyah, Muslim, Fitrani M., 2013 [Semi-natural spawning of the climbing perch (*Anabas testudineus*) under different sex ratio.] *Jurnal Akuakultur Rawa Indonesia* 1(1): 23-33. [in Indonesian]
- Daudpota A. M., Abbas G., Kalhoro I. B., Shah S. S. A. Shah, Kalhoro H., Hafeez-ur-Rehman M., Ghaffar A., 2016 Effect of feeding frequency on growth performance, feed utilization and body composition of juvenile Nile tilapia, *Oreochromis niloticus* (L) reared in low salinity water. *Pakistan Journal Zoologi* 48(1): 171-177.
- Deyab E.S.D.M.S., Hussein E.E.M., 2015 Effects of different feeding rates on growth performance and body composition of red tilapia, *Oreochromis mossambiquense* x *O. niloticus*, Fingerlings, *International Journal of Aquaculture* 5 (12) 1-7.
- Fajrin Ch. N., Buwono I. D., Sriati., 2012 [Addition of tauge extract in feed for spawning success development of goldfish (*Carassius auratus*).] *Jurnal Perikanan & Kelautan*. 3(3): 51-60. [in Indonesian]
- FAO, 2018 The state of world fisheries and aquaculture, meeting the sustainable development goals. Food and Agriculture Organization of the United Nations, Rome. 28 pp.
- Ganzon-Naret, E.S., 2013a Effects of feeding frequency on growth, survival rate and body composition in sea bass (*Lates calcarifer*) juveniles fed a commercial diet under laboratory condition. *Animal Biology & Animal Husbandry Bioflux* 5 (2): 175-176.
- Ganzon-Naret, E.S., 2013b Influence of different feeding rate using commercial dry pellets on growth, feed efficiency and survival for hatchery produced sea bass *Lates calcarifer*. *Animal Biology & Animal Husbandry Bioflux* 5 (2): 241-248.
- Ghosh A., Mohapatra B. C., Chakrabarti P. P., Hussan A., Das A., 2019 Induced breeding of *Catla catla* carried out at low temperature in FRP carp hatchery of Arunachal Pradesh, India. *Journal of Environmental Biology* 40 (3): 328-334.
- Guha D., Mukherjee D., 1991 Seasonal cyclical change in the gonadal activity of Common carp, *Cyprinus carpio* Linn. *Indian Journal of Fisheries* 38 (4): 218-223.
- Herjayanto M., Carman O., Soelistyowati D. T., 2016 [Spawning behavior, reproductive potential of females, and spawning technique optimization of rainbow fish *Iriatherina wernerii* Meinken, 1974.] *Jurnal Ikhtologi Indonesia* 16 (2): 171-183. [in Indonesian]
- Ismi S., Asih Y. N., Kusumawati D., 2014 Improvement of seed production and quality of grouper by hybridization program. *Jurnal Oseanologi Indonesia*. 1 (1): 1-5. [in Indonesian]



- Jia B., Hilaire St. S., Singh K., Gardner I. A., 2016 Farm level return and costs of yellow catfish (*Pelteobagrus fulvidraco*) aquaculture in Guangdong and Zhejiang provinces, China. *Aquaculture Reports* 4: 48-56.
- Kholifah S., Antara I. M., Dewi R. K., 2012 [Carp (*Cyprinus carpio*) hatchery business feasibility in "Sari Nadi" Group, Tua village, Marga district, Tabanan regency.] *E-Jurnal Agribisnis & Agrowisata*. 1 (2): 88-95. [in Indonesian]
- Kour S., Masud S., Khan A., 2016 Effect of organic manure and inorganic fertilizer on the growth and proximate composition of common carp, *Cyprinus carpio*. *Journal of Environmental Biology* 37: 149-153
- Li X. M., Yan Q. Y., Xie S., Hu W., Yu Y., 2013 Gut microbiota contributes to the growth of fast-growing transgenic Common carp (*Cyprinus carpio* L.). *PLoS One* 8 (5): e64577.
- Mantau Z., Rawung J.B.M., Sudarty, 2004 [Effective and efficient carp spawning.] *Jurnal Litbang Pertanian* 23 (2): 68-73. [in Indonesian]
- Masoumeh A., Khara H., Noveiri S. B., Dadras H. 2013 Influence of age of common carp (*Cyprinus carpio*) broodstock on reproductive traits and fertilization. *Turkish Journal of Fisheries and aquatic sciences*. 13: 19-25.
- Ministry of Marine Affairs and Fisheries (MMAF) of Indonesian Republic, 2018a [Fisheries productivity of Indonesia in the forum of Merdeka Barat 9, the ministry of communication and informatics] 49 pp [in Indonesian].
- Ministry of Marine Affairs and Fisheries (MMAF) of Indonesian Republic, 2018b [Volume and aquaculture fisheries production value with major commodities and province of 2017.] *Kelautan dan Perikanan dalam Angka 2018*. 590 pp [in Indonesian]
- Rawat P., Biswas P., Jena A. K., Patel A. B., Pandey P.K., 2019 Effect of dietary incorporation of natural attractants on growth and survival during seed rearing of Indian butter catfish, *Ompok bimaculatus*. *Journal of Environmental Biology* 40: 661-667.
- Saikia S. K., Das D. N., 2009 Feeding ecology of common carp (*Cyprinus carpio* L.) in a rice-fish culture system of the apatani plateau (Arunachal Pradesh, India). *Aquatic Ecology* 43 (2): 559-568.
- Saravi Saeedi S. S., Karami S., Karami B., Shokrzadeh M., 2009 Toxic effects of cobalt chloride on hematological factors of common carp (*Cyprinus carpio*). *Biological Trace Element Research* 1 (3):144-152.
- Shivakumar M., Bala S., Rajanna C., Naveenkumar B. T., 2014 Economics of seed rearing and farming of carps. *International Journal of Fisheries and Aquatic Studies* 2(1): 42-45.
- Soekartawi, 1994 [Cobb-Douglass function-based production economic theory]. PT RajaGrafindo Persada, Jakarta, 258 pp. [in Indonesian]
- Sofia L. A., Muhammad S., Hanani N., Fatah L., 2013 Simulation model of household economy on production and welfare of catfish (*Pangasius hypophthalmus*) farmer in Banjar minapolitan, South Kalimantan. *IOSR Journal* 7(6): 1-8.
- Solomon S.G., Tiarniyu L.O., Okomoda V.T., and Adaga K., 2016 Nutrient profile of commercial aqua-feeds under different storage conditions. *International Journal of Aquaculture*. 6(12): 1-11.
- Statistic Center of Tapin Regency, 2019 [Tapin regency in figures 2019. Statistic Center of Tapin regency, South Kalimantan.] 451 pp [in Indonesian]
- Syaifudin M., Aliah R. S., Muslim, Sumantadinata K., 2007 [Relation between broodstock number and spawning frequency and egg production of humpback grouper (*Cromileptes altivelis*).] *Jurnal Akuakultur Indonesia* 6 (2): 191-196. [in Indonesian]
- United Nations, 2019 World Population Prospects 2019. File POP/1-1: Total Population (both sexes combined) by region, sub region, and country, annually for 1950-2100 (thousand) estimates,1950-2020. Population Division. Department of Economic and Social Affairs. United Nations.
- Valdebenito I. I., Gallegos P. C., Effer B. R., 2013 Gamete quality in fish: evaluation paramaters and determining factors. *Zygote* 23: 177-197.

- Wang Q. D., Zhongjie L., Jian F., Liu J., Shao W. Y., Jing Y., De Silva S.S., 2017 paradigm changes in freshwater aquaculture practice in China: moving towards achieving environmental integrity and sustainability. *Ambio* 47 (4): 410-426.
- Xu J., Ji P., Zhao Z., Zhang Y., Feng J., Wang J., Li J., Zhang X., Zhao L., Liu G., Xu P., Sun X., 2012 Genome-wide snp discovery from transcriptome of four common carp strains. *PLoS ONE* 7(10): 1-9.
- Yulinda E., 2012 [The financial analysis of catfish (*Clarias gariepinus*) hatchery business in Lembah Sari village, Rumbai Pesisir district, Pekanbaru, Riau province.] *Jurnal Hasil Riset* 17 (1): 38-55. [in Indonesian]

Received: 29 February 2020. Accepted: 26 March 2020. Published online: xx April 2020.

Authors:

Bambang Poerwanto, Magister Program of Fisheries Science, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: poerwantobambang2@gmail.com

Idiannor Mahyudin, Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: dr\_idiannor@yahoo.co.id


Leila Ariyani Sofia, Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: leila.ariyani@ulm.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Poerwanto B., Mahyudin I., Sofia L. A., 2020 The analysis of factors influencing carp (*Cyprinus carpio*) seed production in Bungur District, Tapin Regency, South Kalimantan. *AAFL Bioflux* 13(2):xxx-xxx.

REVISI 26 Maret 2020



## The analysis of factors influencing carp (*Cyprinus carpio*) seed production in Bungur District, Tapin Regency, South Kalimantan

<sup>1</sup>Bambang Poerwanto, <sup>2</sup>Idianoor Mahyudin, <sup>2</sup>Leila A. Sofia

<sup>1</sup> Magister Program of Fisheries Science, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia; <sup>2</sup> Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia. Corresponding author: L. A. Sofia, leila.ariyani@ulm.ac.id

**Abstract.** Seed production has become one of the factors supporting the continuity of fish farming activity. Carp, *Cyprinus carpio* is one of the largest freshwater fish productions that support the human food needs in the world. This study aims to know factors affecting carp seed production. The study locality was purposively selected in Bungur District, Tapin Regency, South Kalimantan, and samples were collected through census from 42 carp farmers. The primary data were descriptively quantitatively analyzed using multiple regression. Results showed that there were 4 factors affecting the seed production, number of male spawners, feed amount for fry, spawning frequency, and the use frequency of the same female spawners. Carp seed production development could be done by considering supply of enough number and good quality spawners, good protein feeding and efficient feeding technique, and spawning recovery time of the female spawners.

**Key Words:** fish farmer, pond, stocking density, spawners, feeding.

**Introduction.** World human population in 2019 reached 7.7 billion people. Pada tahun 2050 diperkirakan jumlah penduduk dunia akan meningkat sebesar 2 billion menjadi 9.7 billion (United Nations 2019). Increase in human population causes increase in world food needs including fisheries products. In 2016, the world fisheries production also reached 170.9 million tons, from which 151.2 million tons were used for human consumption and 19.7 million tons for non consumption need. This production comprises fishing fisheries and aquaculture. Nevertheless, the world fishing fisheries production has declined from 2011 to 2016 from 92.2 to 90.9 million tons, while world aquacultural production has significantly risen from 61.8 to 80.0 million tons in the same period (FAO 2018). Systematic fishing in tremendously high numbers, weak reporting, and other complex pressures added with ecosystem change consequences have resulted in decline in fishing fisheries production (Allan et al 2005). Hence, fish farming becomes a very important sector that plays important roles in world food security due to increasing human population and food needs (Wang et al 2017).

Indonesia as the fourth highest human population country, 267.6 million people (United Nations 2019) required fish consumption up to 43.94 kg capita<sup>-1</sup> in 2016. This need was supported by 16.68 million tons of aquaculture and 6.83 million tons of fishing fisheries. Total fisheries production of Indonesia in 2016 reached 23.51 million tons and has contributed to 13.75% of world fisheries production (Ministry of Marine Affairs and Fisheries (MMAF) of RI 2018a). The need of Indonesian fisheries products can be achieved by 70% from aquaculture. This condition makes aquaculture be very crucial. There are several fish commodities developed in Indonesia, one of which is carp, *Cyprinus carpio*. This species becomes the fourth largest freshwater aquacultural commodity in Indonesia with a volume of 316,646 tons in 2017 (MMAF RI 2018b). This species also became the third largest fish culture commodity in the world (Xu et al 2012; FAO 2018) and one of the

Commented [indra1]: all the factors or only some of them?

Commented [indra2]: can you update the info for 2020? Or, at least for 2019?

fish species that can reach high biomass and its distribution has been developed by human due to its important economic value (Arteni & Rosca 2010; Saikia & Das 2009).

The magnitude of carp culture commodity needs fish seed availability as important prerequisite of carp cultivation (Bhuyan et al 2012). The supply of good quality fish seeds is necessary for faster development of aquacultural industry. Hatchery efforts are also needed to reduce the dependence of aquaculture upon the use of wild seeds (Ghosh et al 2019). The development of fish seed quality has been conducted through various techniques, such as application of certain treatment to fish seeding production factors. The treatment could be implemented by previously knowing which production factor yields significant effect on the seeding quantity and quality (Mizuno et al 2012; Karim et al 2016; Hariani & Slamet 2019). Studies on the effect of various factors on carp condition have been done a lot, such as how the intestinal microbes affect the fish growth (Li et al 2013) up to how the toxin effect, such as cobalt chloride, on the hematological condition of carp (Saravi et al 2009). Nevertheless, there are very few studies on carp seeding quality and quantity development. Nowadays, the development of good fish seed quality production is needed in order to meet the seed demand in aquaculture and make the public hatchery be a profitable business (Ismi et al 2014).

Tapin Regency, South Kalimantan province, is one of the regions that run carp culture activities selama periode 2013-2017 with average production level up to 13,198,600 individuals.year<sup>-1</sup> (Animal Husbandary and Fisheries Service of Tapin Regency 2018). Bungur District is one of the largest carp seeding localities in Tapin Regency. The seed production must be raised due to increasing human population and food demand, fisheries product in particular. Increased fish seed production is expected to be able to maintain the sustainability of the carp culture industry. Developing the carp seeding production requires a right strategy. This study aims to know factors affecting the carp seed production in Bungur District, Tapin Regency, the province of South Kalimantan.

## Material and Method

**Study site description.** The work was carried out from January to May 2019 in Bungur District, Tapin Regency, South Kalimantan province with an area of 148.98 km<sup>2</sup>. Bungur District is the largest aquacultural area in Tapin Regency, where 75% is located in Bungur District, and most of it is used for fish hatchery business, including carp seeding (Statistic Center of Tapin Regency 2019). This study site selection was done since Bungur District is the fish seeding center of Tapin Regency with the biggest aquacultural area and number of fish farmers.

**Population and samples.** Fish hatchery farmers in Shabah village and Kalumpang village, Bungur District are 42 people. All population members were taken as research samples through census method.

**Data analyses.** Factors influencing carp seed production were analyzed using descriptively quantitative method and OLS multiple regression with Cobb-Douglas-typed production function approach. This function has been widely used in the field of agricultural economy to show the relationship between number of inputs and outputs (Singh 2008; Osuagwu & Olaifa 2018). It was selected for the following potentials: (1) the Cobb-Douglas (C-D) function estimation will give the regression coefficient that indicates the elasticity, (2) the elasticity indicates the return to scale rate, and (3) the calculation is relatively easier than other functions (Soekartawi 1994; Bhanu Murthy 2002; Vergos et al 2010).

The model applied in this study was as follows:

$$Y = a X_1^{b_1} X_2^{b_2} X_3^{b_3} X_4^{b_4} X_5^{b_5} X_6^{b_6} X_7^{b_7} X_8^{b_8} X_9^{b_9} X_{10}^{b_{10}} D_1 e$$

$$\ln Y = \ln a + b_1 \ln X_1 + b_2 \ln X_2 + b_3 \ln X_3 + b_4 \ln X_4 + b_5 \ln X_5 + b_6 \ln X_6 + b_7 \ln X_7 + b_8 \ln X_8 + b_9 \ln X_9 + b_{10} \ln X_{10} + D_1 + e$$

Commented [indra3]: to be supported by references

Commented [indra4]: make it more explicit

Commented [indra5]: for what period of time? Each year?

Commented [indra6]: also add some info about the statistical analysis since you refer to it in one of the tables

where  $a$  is constant,  $b_1$ - $b_{10}$  are regression coefficients,  $Y$  is the fish seeding production (ind year<sup>-1</sup>);  $X_1$  is pond size (m<sup>2</sup>),  $X_2$  is number of female spawners (ind),  $X_3$  is number of males (ind),  $X_4$  is feed volume for spawners (kg),  $X_5$  is total feed volume for fry (kg),  $X_6$  is amount of manure fertilizer (pack),  $X_7$  is spawning frequency (times),  $X_8$  is use frequency of the same female spawners,  $X_9$  is number of labor used (HOK),  $X_{10}$  is fish farmer's experience (year),  $D_1$  (dummy variable) is trained fish farmer = 1, untrained fish farmer = 0, and  $e$  is error term.

The precision of the C-D production function model was tested with classic assumption tests which include test normality, multicollinearity, and heteroskedasticity, then test goodness of Fit (R<sup>2</sup>), simultaneous variable coefficient test (F test), and partial variable coefficient test (t-test) were done. If all requirements are fulfilled, the model could be used for prediction (Gujarati 1988). The regression coefficients of all inputs used in the model were *a priori* assumed as positive (Ogundari & Akinbogun 2010; Asamoah et al 2012), except the frequency of the same female spawners utilization. Dummy variable coefficient, trained fish farmers  $D_1$ , was assumed to be positive (Onianwa et al 2004; Hyuha et al 2011).

## Results

**Seeding activity description.** The Public Hatchery Unit has been done by the farming group in Bungur District for 8 years in average. The hatchery business run for 1-5 years was 48%, 6-10 years was 24%, and more than 10 years was 29%. The construction of carp seeding pond used by fish farmers in Bungur District is soil ponds with area around 1,490-18,212 m<sup>2</sup> and number of ponds managed were 4-29 units hatchery<sup>-1</sup> (Table 1) with total labors of 12-306 working days. Number of labors in each hatchery was 2-3 people depending upon number of spawning ponds operated. The pond localities are generally near the irrigation route so that the ponds are close enough to the water source. Before use, the fish farmers have fertilized the pond with manure. The organic fertilizer is effective to grow natural food that the productivity and fish growth can be increased (Kour et al 2016). The fertilizer applied in the study site was about 160-2,000 bags yr<sup>-1</sup> hatchery<sup>-1</sup> or about 14 bags m<sup>-2</sup>.

Fish spawner is major production factor in seed production, in either quality or quantity. Both females and males were prepared by the fish farmers. The female spawner candidates were 1.5-3 years old with individual weight of about 2-4 kg, the males were older than 6 months old with an individual weight of 0.2-1 kg. In general, number of spawners prepared by the fish farmers were 4-53 females and 14-178 males per business unit. The female and male spawners ratio was 2:8 each spawning.

Table 1  
Carp hatchery production and input production description in Public Hatchery Unit in Bungur district

Description	Unit	Value		
		Maximum	Minimum	Average
Fish seeding production	ind yr <sup>-1</sup>	2,080,000	100,000	340,520
Total pond size	m <sup>2</sup>	18,212	1,490	6,874
Number of ponds	unit	29	4	9
Pond size	m <sup>2</sup> unit <sup>-1</sup>	1,197	298	782
Number of female spawners	ind	53	4	16
Weight of female spawners	kg ind <sup>-1</sup>	4	2	3
Number of male spawners	ind	178	14	61
Weight of male spawners	kg ind <sup>-1</sup>	1	0.2	0.34
Feed volume of spawners	kg yr <sup>-1</sup>	840	40	318
Feed volume for fish fry	kg yr <sup>-1</sup>	1,560	200	647
Volume of manure fertilizer	bags yr <sup>-1</sup>	2,000	160	651
Spawning frequency	no yr <sup>-1</sup>	32	4	7
Use frequency of the same female spawners	no yr <sup>-1</sup>	6	2	4

Number of labors used	working days	306	12	83
Fish farmer's experience	year	18	1	8

Total time utilized for spawning in one-year is about 8 months. One production cycle requires more or less 60 days, and it could be done 1-2 time mo<sup>-1</sup> or 4-32 times yr<sup>-1</sup>. To maintain the quality of the produced seed, the frequency use of the same female spawners was 2-6 times yr<sup>-1</sup>, and every year the new spawners are prepared to replace the old ones. Total number of seeds produced by the hatchery were 100,000-2,080,000 ind yr<sup>-1</sup>.

These are the fry size ranges that are commonly prepared by the hatchery to the fish farmers under different selling price, 2-3 cm, 3-5 cm, and 5-8 cm long that needed about 2 months. In this stage, the seeds were given fine pellet with total feed weight of 200-1560 kg y<sup>-1</sup> per business unit. Similarly, the spawners were fed with artificial feed as much as 40-840 kg yr<sup>-1</sup> per business unit.

**Financial analysis.** Profit is the main goal of a business. The amount of profit gained will determine the fish farmer's decision making in the business production planning. Carp hatchery business needs an investment of IDR 2.9 million – IDR 254.23 million (Table 2) depending upon land size, number of ponds used, and other supporting facilities, such as keeping house and storage, net/happa, kakaban/egg substrate, bucket, sorting box, and scoop net. Each business unit required the operational costs of IDR 21.99 million – IDR 201.6 million yr<sup>-1</sup> with the highest in feed preparation for the seeds, IDR 1.05 million – IDR 94 million yr<sup>-1</sup> depending upon the seed numbers produced. The additional feed was administered at the life stage of about 7-day old larvae (with yolk sac) up to be ready for sale.

The second highest operational cost is labor's wages, IDR 16.4 million – IDR 27 million. It was allocated for pond preparation or processing, dyke improvement, and feeding. The lowest cost was recorded in male spawner supply, IDR 135,000 – IDR 1,750,000 yr<sup>-1</sup>. Previous studies found no enough difference, with total costs of IDR. 66,262,510.89, approximately 31% is use for feed costs (Shivakumar et al 2014). This finding is in agreement with Kholifah et al (2012) that carp hatchery business has good prospectus with a profit of IDR. 7,329,301.90 cycle<sup>-1</sup>. Different from that of Yulinda (2012) concerning the business financial analysis of catfish *Clarias gariepinus* with much lower profit, averagely IDR. 1,745,194 cycle<sup>-1</sup>. It could result from higher market price of carp fry than that of catfish.

Table 2  
Investment, operational costs, and profit of carp hatchery unit in Bungur District

Description	Unit	Value		
		Maximum	Minimum	Mean
Investment (total):	IDR	<b>254,225,000</b>	<b>2,935,000</b>	<b>46,256,475</b>
Pond	IDR	200,000,000	1,800,000	32,222,500
Other supporting facilities	IDR	54,225,000	1,135,000	14,033,975
Operational costs (total):	IDR yr <sup>-1</sup>	<b>201,600,000</b>	<b>21,995,000</b>	<b>61,318,504</b>
Female spawner preparation	IDR yr <sup>-1</sup>	7,500,000	480,000	2,139,500
Male spawner preparation	IDR yr <sup>-1</sup>	1,750,000	135,000	621,038
Spawner's feed preparation	IDR yr <sup>-1</sup>	22,800,000	900,000	6,181,167
Seed feed preparation	IDR yr <sup>-1</sup>	94,000,000	1,050,000	19,699,750
Manure fertilizer	IDR yr <sup>-1</sup>	33,500,000	2,680,000	10,445,300
Other input preparation	IDR yr <sup>-1</sup>	14,750,000	350,000	3,081,750
Labor's wages	IDR yr <sup>-1</sup>	27,300,000	16,400,000	19,150,000
Revenue and profit:				
Revenue	IDR yr <sup>-1</sup>	268,000,000	57,600,000	120,215,750
Business profit	IDR yr <sup>-1</sup>	<b>66,400,000</b>	<b>35,605,000</b>	<b>58,897,246</b>

The revenue (from seed sales) obtained by a hatchery business unit ranged from IDR 57.6 million to IDR 268 million. The amount of revenue is dependent upon the number of production, seed size, and selling price. The larger the seed size is, the more expensive

Commented [indra7]: why three classes of lengths?

Commented [indra8]: which length: standard, total or fork?

Commented [C9]: These are the fry size ranges that are commonly prepared by the hatchery to the fish farmers under different selling price

Commented [indra10]: to be rephrased, it has no meaning, because the verb is missing

Commented [indra11]: check again the calculations, because these totals here are not correct. For example, for the first column the total exceeds 200 million IDR

the seed will be. Under an agreement among the fish farmers, the selling price was IDR 250 ind<sup>-1</sup> for 2-3 cm long seed, IDR 300 ind<sup>-1</sup> for 3-5 cm length seed, and IDR 350 ind<sup>-1</sup> for 5-8 cm length seed, respectively. Therefore, the business profit varied as well from IDR 35,6 million to IDR 66,4 million.yr<sup>-1</sup>.

**Production and production factors.** The effect of production factors on the carp seed production showed that several production factors (input) involved in production function model gave the consistent values with the theoretical estimate. The production factors of positive value and significant effect on production were number of male spawners, feed volume for seeds, and spawning frequency done by the fish farmers. However, the use of the same female spawners for several spawning activities during the reproductive cycle will cause seed production decline as well (Table 3).

Commented [indra12]: we cannot see this value in the table...

Table 3  
Factors influencing the production of common carp hatchery

Variable	Coefficient	SE	t-Statistic	P value
Intercept	7.543	1.264	5.945	0.000
Total pond size (m <sup>2</sup> )	0.059	0.123	0.477	0.637
No. female spawners (ind.)	0.065	0.106	0.620	0.540
No. male spawners (ind.)	0.343	0.116	2.951**	0.006
Feed volume for spawners (kg)	-0.071	0.088	-0.813	0.423
Feed volume for seeds (kg)	0.381	0.138	2.760**	0.010
No. manure fertilizer (bag)	0.094	0.103	0.920	0.365
Spawning frequency (times)	0.910	0.110	8.241**	0.000
Use frequency of the same female spawners (times)	-0.324	0.139	-2.338**	0.026
Labor's working days	-0.204	0.113	-1.802	0.082
Fish farmer's experience (year)	-0.048	0.079	-0.608	0.548
Trained fish farmers (dummy)	0.112	0.139	0.808	0.425
R-squared		0.828		
Adjusted R-squared		0.765		
Standard error		0.321		
Observations		42		
F-statistic		13.154		
Prob (F-statistic)		0.000		
Durbin-Watson		2.190		

Commented [indra13]: you need to say few words about these indices in the Material and Method section - Data analyses

Note: double asterisks (\*\*) denote significance at 1% level, respectively

Hence, to increase the seed production, the fish farmers could increase several inputs and reduce the use frequency of the same female spawners to give them some time to recover and to have perfect maturation.

**Discussion.** Fish gamete quality, male or female, is determined by a number of factors, such as age, management, food, chemical and physical factors, water quality, and so on, that will impact the survivorship of the embryos, larvae, and/or fry in short or long term (Valdebenito et al 2013). Fish spawners are a major production factor in the hatchery business. The use of higher number of spawners in spawning activity will positively influence the number of eggs produced. The more the spawners are spawned, the higher the spawning frequencies could be done and the higher the number of eggs will be produced (Syaifudin et al 2007). The female carp spawner appropriate for spawning should be 1.5-3 years old with minimum weight of 1.5 kg ind<sup>-1</sup>, while the male should be at least 6 months old with minimum weight of 0.5 kg ind<sup>-1</sup> (Mantau et al 2004; Dadras et al 2013). The weight ratio of female and male is 1:1 (kg m<sup>-2</sup>) meaning that a 2 kg ind<sup>-1</sup> female could be balanced by 3 males of 600-700 g ind<sup>-1</sup>.

Some fish hatchery businesses showed that female-male ratio became factor determining the quality and quantity of seeds produced. For instance, in rainbow fish

*(Iriatherina weneri)* spawning, it is known that the use of sex ratio with 1:3 of total weight is more efficient than 1:1 and 1:2 ratios. It is possible that more male spawners will give higher opportunity to fertilize the eggs. Also, the seed quality produced at the spawner sex ratios of 1:2 and 1:3 gave 5-day old seeds with good survival rate (Herjayanto et al 2016). The climbing perch *Anabas testudineus* at 1:4 ratio spawners yields good fecundity (Burmansyah et al 2013). However, the female-male ratio practiced by the fish farmers in the study site indicated that total weight of male spawner was still below the total weight of female spawner. Thus, number and weight of male spawners need to be added in order to increase the seed production.

Commented [indra14]: provide the scientific name too

*C. carpio* naturally showed 2 clear reproductive cycle per year. Each reproductive cycle is divided into 4 phases. Males have growth phase, maturation, spawning, and resting, while females have preparatory phase, pre-spawning, spawning, and post-spawning (Guha & Mukherjee 1991). At 2 years of age, carp has fecundity of 523,500 eggs and at 3 years of age, the fecundity reaches 657,701 eggs (Aliniya et al 2013). Thus, the more the number of spawners spawned are, the more the number of eggs and eventually more larvae are produced. The availability of spawners spawned will also interchangeably give the opportunity to the spawners to have perfect reproductive cycle. The spawning frequency using the same female spawners gave impact on the seed growth. This study found that the seeds obtained from the spawners often used had lower body mass and shorter size. Also, the larvae produced from the often used spawners looked paler than those from the less used spawners.

Commercial feasibility of intensive fish culture depends on market demand and production costs. The largest part of the production costs is feed, so that it is important to know the optimal feeding and the right feed management strategy implementation (Jia et al 2016). For instance, the right feeding frequency can reduce the production costs and prevent water quality degradation (Daudpota et al 2016). Higher feeding frequency gives positive impact on the fish growth and makes the feed conversion ratio value be more efficient (Ganzon-Naret 2013a). Besides that, the percent of feeding becomes important factor for fish growth and the percentage of optimal feeding determines the aquaculture success (Deyab & Hussein 2015; Aryani et al 2017). The percent feed amount given will influence the fish growth rate as weight increment (Ganzon-Naret 2013b; Sofia et al 2013).

Feed is an important component for the spawners in order to produce good quality eggs and seeds in high number. The fish spawners spawned at the high frequency need good quality feed. Specific additional feeding also affects the aquacultural production. For example, addition of tauge extract in gold fish *Carrasius auratus* feed gives faster spawning cycle (Fajrin et al 2012). Addition of probiotic to the feed can increase body weight and fish endurance, so that the production rises and feed costs decline (Aitkaliyeva et al 2019). Besides feeding technique, the fish farmers should understand how to store the feed before use. According to Solomon et al (2016), storage in closed condition can prevent the nutritional damage of the feed.

Carp seeds produced need different feed treatments with life stages divided into 3 nursing phases. **First**, this stage was done for 7-day old larvae (0.6-0.7 cm long) at a stocking density of 100 ind m<sup>-2</sup> and in a pond of about 60 cm deep. At this stage, they were fed with the aquatic biota living in the pond and fine pellet. In about 15 days of rearing, the fish grew to 1-3 cm long. **Second**, this stage was conducted at the stocking density was about 50 ind m<sup>-2</sup> in about 20 days of rearing up to producing 3-5 cm long fish. **Third**, the stocking density of this stage was about 25 ind m<sup>-2</sup> at rearing period of about 30 days until 5-8 cm long-fish were obtained. After these nursing phases, the seeds are ready to sell, and they are sold when reaching the individual size of 3-5 cm and 5-8 cm long.

So far, feeding in each nursing phase still relied upon the artificial feed so that the operational cost for the fry feed was very high. On the other hand, the economic return of the larvae produced in the hatchery is more important than the fry growth. Therefore, alternative additional feed and feed cost efficiency need to be applied in order to increase the survivorship. One of these is bloodworm (*Tubifex sp.*) supplement. According to Rawat et al (2019), 5% tubifex supplement can be used to increase the fish survivorship, especially in indoor seed rearing.

Commented [indra15]: add the Latin name as well



**Conclusions.** There were 4 factors affecting the production of *C. carpio* hatchery, three of which had positive effect on seed production, namely the number of male spawners, the amount of feed for the fry, and the spawning frequency. While the use frequency of female spawners during the production cycle negatively impact the seed production. This finding suggested that the fish farmers need to pay attention on the availability of sufficient number of good fish spawners, good protein feeding, efficient feeding technique, and longer post-spawning recovery time for female spawners.

**Acknowledgements.** We would greatly appreciate the Fisheries Services of Tapin Regency, the Fisheries Consultant for field information and guidance in this study. We would also thank the fish farmers in the Public Hatchery Unit in Bungur District, Tapin Regency for their collaboration.

## References

- Animal Husbandary and Fisheries Service of Tapin Regency, 2018. [Annual report on fisheries of Tapin Regency, South Kalimantan]. 74 pp. [in Indonesian]
- Aitkaliyeva A.A., Alpeisov S.A., Isbekov K.B., Assylbekova S.Z., Badryzlova N.S., 2019 Influence of different feeds and feed additives on fish-breeding and biological indicators at rearing rainbow trout. *EurAsian Journal of BioSciences* 13(1): 437-442.
- Aliniya, M., Khara, H., Noveiri, S.B., & Dadras, H., 2013 Influence of age of common carp (*Cyprinus carpio*) broodstock on reproductive traits and fertilization. *Turkish Journal of Fisheries and Aquatic Sciences* 13: 19-25.
- Allan J. D., Abeli R., Hogan Z., Revenga C., Taylor B. W., Welcomme R.L., Winemiller K., 2005 Overfishing of inland waters. *BioScience* 55 (12): 1041-1051.
- Arteni M. O., Rosca I., 2010 Some morphological feature of carp (*Cyprinus carpio* Linnaeus, 1758) in different stages of development. *Animal Biology and Animal Husbandry Bioflux* 2 (1): 47 -52.
- Aryani N., Azrita, Mardiah A., Syandri H., 2017 Influence of feeding rate on the growth, feed efficiency and carcass composition of the giant gourami (*Osphronemus goramy*). *Pakistan Journal Zoology* 49(5): 1775-1781.
- Asamoah E.K., Nunoo F.K.E., Asare Y.B.O., Addo S., Sumaila U.R., 2012 Aproduction function analysis of pond aquaculture in Southern Ghana. *Aquaculture Economic & Management* 16: 183-201.
- Bhanumurthy K.V., 2002 Arguing a case for the Cobb-Douglas production function. *Review of Commerce Studies*. Delhi, India. Pp. 75-91.
- Bhuyan A., Akther S., Aktar N., 2012 Present status and fish seed production of the hatcheries of six upazilas of Rajshahi District. *University Journal of Zoology, Rajshahi University* 30: 29-32.
- Burmansyah, Muslim, Fitriani M., 2013 [Semi-natural spawning of the climbing perch (*Anabas testudineus*) under different sex ratio.] *Jurnal Akuakultur Rawa Indonesia* 1(1): 23-33. [in Indonesian]
- Dadras H., Golpour A., Zahmatkesh M., Khara H., Noveiri S.B., Siddique M. A. M., 2017 Effects of age on the reproductive performance of different males and females in bighead carp *Hypophthalmichthys nobilis* (Richardson, 1845). *Comparative Clinical Pathology* 26: 1165-1171.
- Daudpota A. M., Abbas G., Kalhoro I. B., Shah S. S. A. Shah, Kalhoro H., Hafeez-ur-Rehman M., Ghaffar A., 2016 Effect of feeding frequency on growth performance, feed utilization and body composition of juvenile nile tilapia, *oreochromis niloticus* (L) reared in low salinity water. *Pakistan Journal Zoologi* 48(1): 171-177.
- Deyab E.S.D.M.S., Hussein E.E.M., 2015 Effects of different feeding rates on growth performance and body composition of red tilapia, *Oreochromis mossambiquese* x *O. niloticus*, Fingerlings, *International Journal of Aquaculture* 5 (12) 1-7.
- Fajrin Ch. N., Buwono I. D., Sriati., 2012 [Addition of tauge extract in feed for spawning success development of goldfish (*Carassius auratus*).] *Jurnal Perikanan & Kelautan* 3(3): 51-60. [in Indonesian]

- FAO, 2018 The state of world fisheries and aquaculture, meeting the sustainable development goals. Food and Agriculture Organization of the United Nations, Rome. 28 pp.
- Ganzon-Naret, E.S., 2013a Effects of feeding frequency on growth, survival rate and body composition in sea bass (*Lates calcarifer*) juveniles fed a commercial diet under laboratory condition. *Animal Biology & Animal Husbandry Bioflux* 5 (2): 175-176.
- Ganzon-Naret, E.S., 2013b Influence of different feeding rate using commercial dry pellets on growth, feed efficiency and survival for hatchery produced sea bass *Lates calcarifer*. *Animal Biology & Animal Husbandry Bioflux* 5 (2): 241-248.
- Ghosh A., Mohapatra B. C., Chakrabarti P. P., Hussan A., Das A., 2019 Induced breeding of *Catla catla* carried out at low temperature in FRP carp hatchery of Arunachal Pradesh, India. *Journal of Environmental Biology* 40 (3): 328-334.
- Guha D., Mukherjee D., 1991 Seasonal cyclical change in the gonadal activity of Common carp, *Cyprinus carpio* Linn. *Indian Journal of Fisheries* 38 (4): 218-223.
- Gujarati, D.N., 1988 *Basic Econometrics*. McGraw-Hill. 705 pp.
- Hariani D., Slamet WK P., 2019 Combination of feed protein level and laserpuncture induction of broodstock catfish (*Clarias* sp.) to increase estrogen, vitellogenin, and egg quality. *EurAsian Journal of BioSciences* 13: 769-779.
- Herjayanto M., Carman O., Soelistyowati D. T., 2016 [Spawning behavior, reproductive potential of females, and spawning technique optimization of rainbow fish *Iriatherina wernerii* Meinken, 1974.] *Jurnal Ikhtologi Indonesia* 16 (2): 171-183. [in Indonesian]
- Hyuha T.S., Bukenya J.O., Twinamasiko J., Moinar J., 2011 Profitability analysis of small scale aquaculture enterprises in Central Uganda. *International Journal of Fisheries and Aquaculture* 2(15): 271-278.
- Ismi S., Asih Y. N., Kusumawati D., 2014 Improvement of seed production and quality of grouper by hybridization program. *Jurnal Oseanologi Indonesia* 1 (1): 1-5. [in Indonesian]
- Jia B., Hilaire St. S., Singh K., Gardner I. A., 2016 Farm level return and costs of yellow catfish (*Pelteobagrus fulvidraco*) aquaculture in Guangdong and Zhejiang provinces, China. *Aquaculture Reports* 4: 48-56.
- Karim M., Keus H. J., Ullah Md. H., Kassam L., Phillips M., Beveridge M., 2016 Investing in carp seed quality improvements in homestead aquaculture: lessons from Bangladesh. *Aquaculture* 453: 19-30.
- Kholifah S., Antara I. M., Dewi R. K., 2012 [Carp (*Cyprinus carpio*) hatchery business feasibility in "Sari Nadi" Group, Tua village, Marga district, Tabanan regency.] *E-Jurnal Agribisnis & Agrowisata* 1 (2): 88-95. [in Indonesian]
- Kour S., Masud S., Khan A., 2016 Effect of organic manure and inorganic fertilizer on the growth and proximate composition of common carp, *Cyprinus carpio*. *Journal of Environmental Biology* 37: 149-153
- Li X. M., Yan Q. Y., Xie S., Hu W., Yu Y., 2013 Gut microbiota contributes to the growth of fast-growing transgenic Common carp (*Cyprinus carpio* L.). *PLoS One* 8 (5): e64577.
- Mantau Z., Rawung J.B.M., Sudarty, 2004 [Effective and efficient carp spawning.] *Jurnal Litbang Pertanian* 23 (2): 68-73. [in Indonesian]
- Ministry of Marine Affairs and Fisheries (MMAF) of Indonesian Republic, 2018a [Fisheries productivity of Indonesia in the forum of Merdeka Barat 9, the ministry of communication and informatics] 49 pp [in Indonesian].
- Ministry of Marine Affairs and Fisheries (MMAF) of Indonesian Republic, 2018b [Volume and aquaculture fisheries production value with major commodities and province of 2017.] *Kelautan dan Perikanan dalam Angka 2018*. 590 pp [in Indonesian]
- Mizuno K., Yamaguchi S. S., Miura C., Miura T., 2012 Method for efficiently obtaining fertilized eggs from the black scraper *Thamnaconus modestus* by natural spawning in captivity. *Fisheries Science* 78: 1059-1064.
- Ogundari K., Akinbogun O.O., 2010 Modeling technical efficiency with production risk: a study of fish farm in Nigeria. *Marine Resource Economics* 25: 295-308.

- Onianwa O., Wheelock G., Gyawali B., Gan J., Dubois M., Schelhas J., 2004 An analysis of factors affecting participation behavior of limited resource farmers in agricultural cost-share programs in Alabama. *Journal of Agribusiness* 22 (1): 17-29.
- Osuagwu E.S, Olaiifa E., 2018 Effects of oil spills on fish production in the Niger Delta. *PLoS ONE* 13(10): e0205114. 14 pp.
- Rawat P., Biswas P., Jena A. K., Patel A. B., Pandey P.K., 2019 Effect of dietary incorporation of natural attractants on growth and survival during seed rearing of Indian butter catfish, *Ompok bimaculatus*. *Journal of Environmental Biology* 40: 661-667.
- Saikia S. K., Das D. N., 2009 Feeding ecology of common carp (*Cyprinus carpio* L.) in a rice-fish culture system of the apatani plateau (Arunachal Pradesh, India). *Aquatic Ecology* 43 (2): 559-568.
- Saravi Saeedi S. S., Karami S., Karami B., Shokrzadeh M., 2009 Toxic effects of cobalt chloride on hematological factors of common carp (*Cyprinus carpio*). *Biological Trace Element Research* 1 (3):144-152.
- Shivakumar M., Bala S., Rajanna C., Naveenkumar B. T., 2014 Economics of seed rearing and farming of carps. *International Journal of Fisheries and Aquatic Studies* 2(1): 42-45.
- Singh K., 2008 Farm specific economic efficiency of fish production in South Tripura District: a stochastic frontier approach. *Indian Journal of Agricultural Economics* 63 (4): 598-613.
- Soekartawi, 1994 [Cobb-Dougllass function-based production economic theory]. PT RajaGrafindo Persada, Jakarta, 258 pp. [in Indonesian]
- Sofia L. A., Muhammad S., Hanani N., Fatah L., 2013 Simulation model of household economy on production and welfare of catfish (*Pangasius hypophthalmus*) farmer in Banjar minapolitan, South Kalimantan. *IOSR Journal* 7(6): 1-8.
- Solomon S.G., Tiamiyu L.O., Okomoda V.T., and Adaga K., 2016 Nutrient profile of commercial aqua-feeds under different storage conditions. *International Journal of Aquaculture* 6(12): 1-11.
- Statistic Center of Tapin Regency, 2019 [Tapin regency in figures 2019. Statistic Center of Tapin regency, South Kalimantan.] 451 pp [in Indonesian]
- Syaifudin M., Aliah R. S., Muslim, Sumantadinata K., 2007 [Relation between broodstock number and spawning frequency and egg production of humpback grouper (*Cromileptes altivelis*).] *Jurnal Akuakultur Indonesia* 6 (2): 191-196. [in Indonesian]
- United Nations, 2019 World Population Prospects 2019. Population Division. Department of Economic and Social Affairs. United Nations. <https://www.un.org/development/desa/publications/world-population-prospects-2019-highlights.html>
- Valdebenito I. I., Gallegos P. C., Effer B. R., 2013 Gamete quality in fish: evaluation paramaters and determining factors. *Zygote* 23: 177-197.
- Vergos K., Christopoulos A., Krystallidis P., Papandroni O., 2010 Economies of scale concentration in the Greek and the Norwegian aquaculture industry. An empirical study. *International Journal of Bussines Management Economic Res.* 1(1): 70-78.
- Wang Q. D., Zhongjie L., Jian F., Liu J., Shao W. Y., Jing Y., De Silva S.S., 2017 paradigm changes in freshwater aquaculture practice in China: moving towards archieving environmental intergrity and sustainability. *Ambio* 47 (4): 410-426.
- Xu J., Ji P., Zhao Z., Zhang Y., Feng J., Wang J., Li J., Zhang X., Zhao L., Liu G., Xu P., Sun X., 2012 Genome-wide snp discovery from transcriptome of four common carp strains. *PLoS ONE* 7(10): 1-9.
- Yulinda E., 2012 [The financial analysis of catfish (*Clarias gariepinus*) hatchery business in Lembah Sari village, Rumbai Pesisir district, Pekanbaru, Riau province.] *Jurnal Hasil Riset* 17 (1): 38-55. [in Indonesian]

Received: 29 February 2020. Accepted: 26 March 2020. Published online: xx April 2020.

Authors:

Bambang Poerwanto, Magister Program of Fisheries Science, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: poerwantobambang2@gmail.com

Idiannor Mahyudin, Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: dr\_idiannor@yahoo.co.id

Leila Ariyani Sofia, Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: leila.ariyani@ulm.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Poerwanto B., Mahyudin I., Sofia L. A., 2020 The analysis of factors influencing carp (*Cyprinus carpio*) seed production in Bungur District, Tapin Regency, South Kalimantan. AACL Bioflux 13(2):xxx-xxx.

On Sunday, April 5, 2020, 11:06:42 PM GMT+8, gavriloaie ionel claudiu <[ionelclaudiu@yahoo.com](mailto:ionelclaudiu@yahoo.com)> wrote:

Dear Dr. Leila Sofia,

There are still two minor comments to be checked and addressed. So, kindly use the document attached.

I edited the whole references list following the journal's format and I also did few corrections.


Also check the title online: <http://www.bioflux.com.ro/home/volume-13-2-2020/>.

With many thanks,  
Claudiu G.

2 Attachments



OPEN ACCESS



Bioflux (publishing house)  
54 Ceahlau Street,  
Cluj-Napoca 400488,  
Romania, European Union

### Certificate/Letter of acceptance

This certificate shows that your manuscript entitled:

**The analysis of factors influencing carp (*Cyprinus carpio*) seed production in Bungur District, Tapin Regency, South Kalimantan**

Authors:

**Bambang Poerwanto, Idianoor Mahyudin, Leila A. Sofia**


was accepted for publication in vol 13, issue 2 (2020) of the scientific/academic journal:  
Aquaculture, Aquarium, Conservation & Legislation –  
International Journal of the Bioflux Society.

Thank you for publishing with us!

Cordially yours,

Editor-in-chief:  
Claudiu Gavriiloaie, PhD





## The analysis of factors influencing carp (*Cyprinus carpio*) seed production in Bungur District, Tapin Regency, South Kalimantan

<sup>1</sup>Bambang Poerwanto, <sup>2</sup>Idianoor Mahyudin, <sup>2</sup>Leila A. Sofia

<sup>1</sup> Magister Program of Fisheries Science, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia; <sup>2</sup> Fisheries Services of Tapin Regency, Jl. Jenderal Sudirman 2.5 km, Rantau, South Kalimantan; <sup>3</sup> Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia. Corresponding author: L. A. Sofia, leila.ariyani@ulm.ac.id

**Abstract.** Seed production has become one of the factors supporting the continuity of fish farming activity. Carp, *Cyprinus carpio* is one of the largest freshwater fish productions that support the human food needs in the world. This study aims to know several factors affecting carp seed production. The study locality was purposively selected in Bungur District, Tapin Regency, South Kalimantan, and samples were collected through census from 42 carp farmers. The primary data were descriptively quantitatively analyzed using multiple regression. Results showed that there were 4 factors affecting the seed production, number of male spawners, feed amount for fry, spawning frequency, and the use frequency of the same female spawners. Carp seed production development could be done by considering the supply of enough number and good quality spawners, good protein feeding and efficient feeding technique, and spawning recovery time of the female spawners.

**Key Words:** fish farmer, pond, stocking density, spawners, feeding.

**Introduction.** World human population in 2019 reached 7.7 billion people and is estimated to be 9.7 billion in 2050 (United Nations 2019). Increase in human population causes increase in world food needs including fisheries products. In 2016, the world fisheries production also reached 170.9 million tons, from which 151.2 million tons were used for human consumption and 19.7 million tons for non consumption need. This production comprises fishing fisheries and aquaculture. Nevertheless, the world fishing fisheries production has declined from 2011 to 2016 from 92.2 to 90.9 million tons, while world aquaculture production has significantly risen from 61.8 to 80.0 million tons in the same period (FAO 2018). Systematic fishing in tremendously high numbers, weak reporting, and other complex pressures added with ecosystem change consequences have resulted in decline in fishing fisheries production (Allan et al 2005). Hence, fish farming becomes a very important sector that plays important roles in world food security due to increasing human population and food needs (Wang et al 2017).

Indonesia as the fourth highest human population country, 267.6 million people (United Nations 2019) required fish consumption up to 43.94 kg capita<sup>-1</sup> in 2016. This need was supported by 16.68 million tons of aquaculture and 6.83 million tons of fishing fisheries. Total fisheries production of Indonesia in 2016 reached 23.51 million tons and has contributed to 13.75% of world fisheries production (Ministry of Marine Affairs and Fisheries (MMAF) of RI 2018a). The need of Indonesian fisheries products can be achieved by 70% from aquaculture. This condition makes aquaculture be very crucial. There are several fish commodities developed in Indonesia, one of which is carp, *Cyprinus carpio*. This species becomes the fourth largest freshwater aquaculture commodity in Indonesia with a volume of 316,646 tons in 2017 (MMAF RI 2018b). This species also became the third largest fish culture commodity in the world (Xu et al 2012; FAO 2018) and one of the fish species that can reach high biomass and its distribution has been developed by human due to its important economic value (Arteni & Rosca 2010; Saikia & Das 2009).

The magnitude of carp culture commodity needs fish seed availability as important prerequisite of carp cultivation (Bhuyan et al 2011). The supply of good quality fish seeds is necessary for faster development of aquaculture industry. Hatchery efforts are also needed to reduce the dependence of aquaculture upon the use of wild seeds (Ghosh et al 2019). The development of carp seed quality has been conducted through various techniques, such as application of certain treatment to carp seeding production factors. The treatment could be implemented by previously knowing which production factor yields significant effect on the seeding quantity and quality (Mizuno et al 2012; Karim et al 2016; Hariani & Pungky Slamet 2019). Studies on the effect of various factors on carp condition have been done a lot, such as how the intestinal microbes affect the fish growth (Li et al 2013) up to how the toxin effect, such as cobalt chloride, on the hematological condition of carp (Saravi et al 2009). Nevertheless, there are very few studies on carp seeding quality and quantity development. Nowadays, the development of good fish seed quality production is needed in order to meet the seed demand in aquaculture and make the public hatchery be a profitable business (Ismi et al 2013).

Tapin Regency, South Kalimantan province, is one of the regions that run carp culture activities during 2013-2017 with mean production level up to 13,198,600 ind.yr<sup>-1</sup> (Animal Husbandary and Fisheries Service of Tapin Regency 2018). Bungur District is one of the largest carp seeding localities in Tapin Regency. The seed production must be raised due to increasing human population and food demand, fisheries product in particular. Increased fish seed production is expected to be able to maintain the sustainability of the carp culture industry. Developing the carp seeding production requires a right strategy. This study aims to know factors affecting the carp seed production in Bungur District, Tapin Regency, the province of South Kalimantan.

## Material and Method

**Study site description.** The work was carried out from January to May 2019 in Bungur District, Tapin Regency, South Kalimantan province with an area of 148.98 km<sup>2</sup>. Bungur District is the largest aquaculture area in Tapin Regency, where 75% is located in Bungur District, and most of it is used for fish hatchery business, including carp seeding (Statistic Center of Tapin Regency 2019). This study site selection was done since Bungur District is the fish seeding center of Tapin Regency with the biggest aquaculture area and number of fish farmers.

**Population and samples.** Fish hatchery farmers in Shabah village and Kalumpang village, Bungur District are 42 people. All population members were taken as research samples through census method.

**Data analyses.** Factors influencing carp seed production were analyzed using descriptively quantitative method and OLS multiple regression with Cobb-Douglas-typed production function approach. This function has been widely used in the field of agricultural economy to show the relationship between number of inputs and outputs (Singh 2008; Osuagwu & Olaifa 2018). It was selected for the following potentials: (1) the Cobb-Douglas (C-D) function estimation will give the regression coefficient that indicates the elasticity, (2) the elasticity indicates the return to scale rate, and (3) the calculation is relatively easier than other functions (Soekartawi 1994; Bhanu Murthy 2002; Vergos et al 2010).

The model applied in this study was as follows:

$$Y = a X_1^{b_1} X_2^{b_2} X_3^{b_3} X_4^{b_4} X_5^{b_5} X_6^{b_6} X_7^{b_7} X_8^{b_8} X_9^{b_9} X_{10}^{b_{10}} D_1 e$$

$$\ln Y = \ln a + b_1 \ln X_1 + b_2 \ln X_2 + b_3 \ln X_3 + b_4 \ln X_4 + b_5 \ln X_5 + b_6 \ln X_6 + b_7 \ln X_7 + b_8 \ln X_8 + b_9 \ln X_9 + b_{10} \ln X_{10} + D_1 + e$$

where a is constant, b<sub>1</sub>-b<sub>10</sub> are regression coefficients, Y is the fish seeding production (ind year<sup>-1</sup>); X<sub>1</sub> is pond size (m<sup>2</sup>), X<sub>2</sub> is number of female spawners (ind), X<sub>3</sub> is number of males (ind), X<sub>4</sub> is feed volume for spawners (kg), X<sub>5</sub> is total feed volume for fry (kg), X<sub>6</sub> is amount of manure fertilizer (pack), X<sub>7</sub> is spawning frequency (times), X<sub>8</sub> is use frequency of the same female spawners, X<sub>9</sub> is number of labor used (HOK), X<sub>10</sub> is fish farmer's experience



(year),  $D_1$  (dummy variable) is trained fish farmer = 1, untrained fish farmer = 0, and  $e$  is error term.

The precision of the C-D production function model was tested with classic assumption tests which include test normality, multicollinearity, and heteroskedasticity, then test goodness of Fit ( $R^2$ ), simultaneous variable coefficient test (F test), and partial variable coefficient test (t-test) were done. If all requirements are fulfilled, the model could be used for prediction (Gujarati 1988). The regression coefficients of all inputs used in the model were *a priori* assumed as positive (Ogundari & Akinbogun 2010; Asamoah et al 2012), except the frequency of the same female spawners utilization. Dummy variable coefficient, trained fish farmers  $D_1$ , was assumed to be positive (Onianwa et al 2004; Hyuha et al 2011).

## Results

**Seeding activity description.** The Public Hatchery Unit has been done by the farming group in Bungur District for 8 years in average. The hatchery business run for 1-5 years was 48%, 6-10 years was 24%, and more than 10 years was 29%. The construction of carp seeding pond used by fish farmers in Bungur District is soil ponds with area around 1,490-18,212  $m^2$  and number of ponds managed were 4-29 units hatchery<sup>-1</sup> (Table 1) with total labors of 12-306 working days. Number of labors in each hatchery was 2-3 people depending upon number of spawning ponds operated. The pond localities are generally near the irrigation route so that the ponds are close enough to the water source. Before use, the fish farmers have fertilized the pond with manure. The organic fertilizer is effective to grow natural food that the productivity and fish growth can be increased (Kour et al 2016). The fertilizer applied in the study site was about 160-2,000 bags yr<sup>-1</sup> hatchery<sup>-1</sup> or about 14 bags  $m^{-2}$ .

Fish spawner is major production factor in seed production, in either quality or quantity. Both females and males were prepared by the fish farmers. The female spawner candidates were 1.5-3 years old with individual weight of about 2-4 kg, the males were older than 6 months old with an individual weight of 0.2-1 kg. In general, number of spawners prepared by the fish farmers were 4-53 females and 14-178 males per business unit. The female and male spawners ratio was 2:8 each spawning.

Table 1  
Carp hatchery production and input production description in Public Hatchery Unit in Bungur district

Description	Unit	Value		
		Maximum	Minimum	Average
Fish seeding production	ind yr <sup>-1</sup>	2,080,000	100,000	340,520
Total pond size	$m^2$	18,212	1,490	6,874
Number of ponds	unit	29	4	9
Pond size	$m^2$ unit <sup>-1</sup>	1,197	298	782
Number of female spawners	ind	53	4	16
Weight of female spawners	kg ind <sup>-1</sup>	4	2	3
Number of male spawners	ind	178	14	61
Weight of male spawners	kg ind <sup>-1</sup>	1	0.2	0.34
Feed volume of spawners	kg yr <sup>-1</sup>	840	40	318
Feed volume for fish fry	kg yr <sup>-1</sup>	1,560	200	647
Volume of manure fertilizer	bags yr <sup>-1</sup>	2,000	160	651
Spawning frequency	no yr <sup>-1</sup>	32	4	7
Use frequency of the same female spawners	no yr <sup>-1</sup>	6	2	4
Number of labor used	working days	306	12	83
Fish farmer's experience	year	18	1	8

Total time utilized for spawning in one-year is about 8 months. One production cycle requires more or less 60 days, and it could be done 1-2 times  $mo^{-1}$  or 4-32 times  $yr^{-1}$ . To maintain the quality of the produced seed, the frequency use of the same female spawners

was 2-6 times yr<sup>-1</sup>, and every year the new spawners are prepared to replace the old ones. Total number of seeds produced by the hatchery were 100,000-2,080,000 ind yr<sup>-1</sup>.

These are the fry size ranges that are commonly prepared by the hatchery to the fish farmers under different selling price, 2-3 cm, 3-5 cm, and 5-8 cm total length that need about 2 months. In this stage, the seeds were given fine pellet with total feed weight of 200-1560 kg yr<sup>-1</sup> per business unit. Similarly, the spawners were fed with artificial feed as much as 40-840 kg yr<sup>-1</sup> per business unit.

**Commented [indra1]:** all sizes need this amount of time? I suppose not. So, make it more clear or simply remove it.

**Fincancial analysis.** Profit is the main goal of a business. The amount of profit gained will determine the fish farmer's decision making in the business production planning. Carp hatchery business needs an investment of IDR 2.9 million – IDR 254.23 million (Table 2) depending upon land size, number of ponds used, and other supporting facilities, such as keeping house and storage, net/happa, kakaban/egg substrate, bucket, sorting box, and scoop net. Each business unit required the operational costs of IDR 29.69 millio – IDR 159.34 million yr<sup>-1</sup> with the highest in feed preparation for the seeds, IDR 1.05 million – IDR 94 million yr<sup>-1</sup> depending upon the seed numbers produced. The additional feed was administered at the life stage of about 7-day old larvae (with yolk sac) up to be ready for sale.

The second highest operational cost is labor's wages, IDR 16.4 million – IDR 27 million. It was allocated for pond preparation or processing, dyke improvement, and feeding. The lowest cost was recorded in male spawner supply, IDR 135,000 – IDR 1,750,000 yr<sup>-1</sup>. Previous studies found no enough difference, with total costs of IDR. 66,262,510.89, approximately 31% is used for feed costs (Shivakumar et al 2014). This finding is in agreement with Kholifah et al (2012) that carp hatchery business has good prospectus with a profit of IDR. 7,329,301.90 cycle<sup>-1</sup>, but different from that of Yulinda (2012) concerning the business financial analysis of catfish *Clarias gariepinus* with much lower profit, averagely IDR. 1,745,194 cycle<sup>-1</sup>. It could result from higher market price of carp fry than that of catfish.

Table 2  
Investment, operational costs, and profit of carp hatchery unit in Bungur District

Description	Unit	Value		
		Maximum	Minimum	Mean
Investment (total):	IDR	<b>254,225,000</b>	<b>2,935,000</b>	<b>46,256,475</b>
Pond	IDR	200,000,000	1,800,000	32,222,500
Other supporting facilities	IDR	54,225,000	1,135,000	14,033,975
Operational costs (total):	IDR yr <sup>-1</sup>	<b>201,600,000</b>	<b>21,995,000</b>	<b>61,318,505</b>
Female spawner preparation	IDR yr <sup>-1</sup>	7,500,000	480,000	2,139,500
Male spawner preparation	IDR yr <sup>-1</sup>	1,750,000	135,000	621,038
Spawner's feed preparation	IDR yr <sup>-1</sup>	22,800,000	900,000	6,181,167
Seed feed preparation	IDR yr <sup>-1</sup>	94,000,000	1,050,000	19,699,750
Manure fertilizer	IDR yr <sup>-1</sup>	33,500,000	2,680,000	10,445,300
Other input preparation	IDR yr <sup>-1</sup>	14,750,000	350,000	3,081,750
Labor's wages	IDR yr <sup>-1</sup>	27,300,000	16,400,000	19,150,000
Revenue and profit:				
Revenue	IDR yr <sup>-1</sup>	268,000,000	57,600,000	120,215,750
Business profit	IDR yr <sup>-1</sup>	66,400,000	35,605,000	58,897,246

The revenue (from seed sales) obtained by a hatchery business unit ranged from IDR 57.6 million to IDR 268 million. The amount of revenue is dependent upon the number of production, seed size, and selling price. The larger the seed size is, the more expensive the seed will be. Under an agreement among the fish farmers, the selling price was IDR 250 ind<sup>-1</sup> for 2-3 cm total length seed, IDR 300 ind<sup>-1</sup> for 3-5 cm total length seed, and IDR 350 ind<sup>-1</sup> for 5-8 cm total length seed, respectively. Therefore, the business profit varied as well from IDR 35,6 to IDR 66,4 million yr<sup>-1</sup>.

**Production and production factors.** The effect of production factors on the carp seed production showed that several production factors (input) involved in the production function model gave the consistent values with the theoretical estimate. The production factors of positive value and significant effect on production were number of male spawners, feed volume for seeds, and spawning frequency done by the fish farmers. However, the use of the same female spawners for several spawning activities during the reproductive cycle will cause seed production decline as well (Table 3).

Table 3  
Factors influencing the production of common carp hatchery

Variable	Coefficient	SE	t-Statistic	P value
Intercept	7.543	1.264	5.945	0.000
Total pond size (m <sup>2</sup> )	0.059	0.123	0.477	0.637
No. female spawners (ind.)	0.065	0.106	0.620	0.540
No. male spawners (ind.)	0.343	0.116	2.951**	0.006
Feed volume for spawners (kg)	-0.071	0.088	-0.813	0.423
Feed volume for seeds (kg)	0.381	0.138	2.760**	0.010
No. manure fertilizer (bag)	0.094	0.103	0.920	0.365
Spawning frequency (times)	0.910	0.110	8.241**	0.000
Use frequency of the same female spawners (times)	-0.324	0.139	-2.338**	0.026
Labor's working days	-0.204	0.113	-1.802	0.082
Fish farmer's experience (year)	-0.048	0.079	-0.608	0.548
Trained fish farmers (dummy)	0.112	0.139	0.808	0.425
R-squared		0.828		
Adjusted R-squared		0.765		
Standard error		0.321		
Observations		42		
F-statistic		13.154		
Prob (F-statistic)		0.000		
Durbin-Watson		2.190		

Note: double asterisks (\*\*) denote significance at 1% level, respectively.

Hence, to increase the seed production, the fish farmers could increase several inputs and reduce the use frequency of the same female spawners to give them some time to recover and to have perfect maturation.

**Discussion.** Fish gamete quality, male or female, is determined by a number of factors, such as age, management, food, chemical and physical factors, water quality, and so on, that will impact the survivorship of the embryos, larvae, and/or fry in short or long term (Valdebenito et al 2013). Fish spawners are a major production factor in the hatchery business. The use of higher number of spawners in spawning activity will positively influence the number of eggs produced. The more the spawners are spawned, the higher the spawning frequencies could be done and the higher the number of eggs will be produced (Syaifudin et al 2007). The female carp spawner appropriate for spawning should be 1.5-3 years old with minimum weight of 1.5 kg ind<sup>-1</sup>, while the male should be at least 6 months old with minimum weight of 0.5 kg ind<sup>-1</sup> (Mantau & Rawung 2004; Aliniya et al 2013). The weight ratio of female and male is 1:1 (kg m<sup>-2</sup>) meaning that a 2 kg ind<sup>-1</sup> female could be balanced by 3 males of 600-700 g ind<sup>-1</sup>.

Some fish hatchery businesses showed that female-male ratio became factor determining the quality and quantity of seeds produced. For instance, in rainbow fish (*Iriatherina werneri*) spawning, it is known that the use of sex ratio with 1:3 of total weight is more efficient than 1:1 and 1:2 ratios. It is possible that more male spawners will give higher opportunity to fertilize the eggs. Also, the seed quality produced at the spawner sex ratios of 1:2 and 1:3 gave 5-day old seeds with good survival rate (Herjayanto et al 2016). The climbing perch *Anabas testudineus* at 1:4 ratio spawners yields good fecundity

(Burmansyah et al 2013). However, the female-male ratio practiced by the fish farmers in the study site indicated that total weight of male spawner was still below the total weight of female spawner. Thus, number and weight of male spawners need to be added in order to increase the seed production.

*C. carpio* naturally showed 2 clear reproductive cycles per year. Each reproductive cycle is divided into 4 phases. Males have growth phase, maturation, spawning, and resting, while females have preparatory phase, pre-spawning, spawning, and post-spawning (Guha & Mukherjee 1991). At 2 years of age, carp has fecundity of 523,500 eggs and at 3 years of age, the fecundity reaches 657,701 eggs (Aliniya et al 2013). Thus, the more the number of spawners spawned are, the more the number of eggs and eventually more larvae are produced. The availability of spawners spawned will also interchangeably give the opportunity to the spawners to have perfect reproductive cycle. The spawning frequency using the same female spawners gave impact on the seed growth. This study found that the seeds obtained from the spawners often used had lower body mass and shorter size. Also, the larvae produced from the often used spawners looked paler than those from the less used spawners.

Commercial feasibility of intensive fish culture depends on market demand and production costs. The largest part of the production costs is feed, so that it is important to know the optimal feeding and the right feed management strategy implementation (Jia et al 2016). For instance, the right feeding frequency can reduce the production costs and prevent water quality degradation (Daudpota et al 2016). Higher feeding frequency gives positive impact on the fish growth and makes the feed conversion ratio value be more efficient (Ganzon-Naret 2013a). Besides that, the percent of feeding becomes important factor for fish growth and the percentage of optimal feeding determines the aquaculture success (Deyab & Hussein 2015; Aryani et al 2017). The percent feed amount given will influence the fish growth rate as weight increment (Ganzon-Naret 2013b; Sofia et al 2013).

Feed is an important component for the spawners in order to produce good quality eggs and seeds in high number. The fish spawners spawned at the high frequency need good quality feed. Specific additional feeding also affects the aquaculture production. For example, addition of tauge extract in gold fish *Carrasius auratus* feed gives faster spawning cycle (Fajrin et al 2012). Addition of probiotic to the feed can also increase body weight and fish endurance, so that the production rises and feed costs decline (Aitkaliyeva et al 2019). Besides feeding technique, the fish farmers should understand how to store the feed before use. According to Solomon et al (2016), storage in closed condition can prevent the nutritional damage of the feed.

Carp seeds produced need different feed treatments with life stages divided into 3 nursing phases. First, this stage was done for 7-day old larvae (0.6-0.7 cm total length) at a stocking density of 100 ind m<sup>-2</sup> and in a pond of about 60 cm deep. At this stage, they were fed with the aquatic biota living in the pond and fine pellet. In about 15 days of rearing, the fish grew to 1-3 cm total length. Second, this stage was conducted at the stocking density of about 50 ind m<sup>-2</sup> in about 20 days of rearing up to producing 3-5 cm total length fish. Third, the stocking density of this stage was about 25 ind m<sup>-2</sup> at rearing period of about 30 days until 5-8 cm total length fish were obtained. After these nursing phases, the seeds are ready to sell, and they are sold when reaching the individual size of 3-5 cm and 5-8 cm long.

So far, feeding in each nursing phase still relied upon the artificial feed so that the operational cost for the fry feed was very high. On the other hand, the economic return of the larvae produced in the hatchery is more important than the fry growth. Therefore, alternative additional feed and feed cost efficiency need to be applied in order to increase the survivorship. One of these is bloodworm (*Tubifex* sp.) supplement. According to Rawat et al (2019), 5% tubifex supplement can be used to increase the fish survivorship, especially in indoor seed rearing.

**Conclusions.** There were 4 factors affecting the production of *C. carpio* hatchery, three of which had positive effect on seed production, namely the number of male spawners, the amount of feed for the fry, and the spawning frequency. While the use frequency of female spawners during the production cycle negatively impact the seed production. This finding suggested that the fish farmers need to pay attention on the availability of sufficient

number of good fish spawners, good protein feeding, efficient feeding technique, and longer post-spawning recovery time for female spawners.

**Acknowledgements.** We would greatly appreciate the Animal Husbandry and Fisheries Services of Tapin Regency, the Fisheries Consultant for field information and guidance in this study. We would also thank the fish farmers in the Public Hatchery Unit in Bungur District, Tapin Regency for their collaboration.

## References

- Aitkaliyeva A. A., Alpeisov S. A., Isbekov K. B., Assylbekova S. Z., Badryzlova N. S., 2019 Influence of different feeds and feed additives on fish-breeding and biological indicators at rearing rainbow trout. *EurAsian Journal of BioSciences* 13(1):437-442.
- Aliniya M., Nezami S., Khara H., Noveiri S. B., Dadras H., Fallah Z., 2013 Influence of age of common carp (*Cyprinus carpio*) broodstock on reproductive traits and fertilization. *Turkish Journal of Fisheries and Aquatic Sciences* 13:19-25.
- Allan J. D., Abeli R., Hogan Z., Revenga C., Taylor B. W., Welcomme R. L., Winemiller K., 2005 Overfishing of inland waters. *BioScience* 55(12):1041-1051.
- Animal Husbandry and Fisheries Service of Tapin Regency, 2018 [Annual report on fisheries of Tapin Regency, South Kalimantan]. 74 pp. [in Indonesian]
- Arteni M. O., Rosca I., 2010 Some morphological features of carp (*Cyprinus carpio* Linnaeus, 1758) in different stages of development. *ABAH Bioflux* 2(1):47-52.
- Aryani N., Azrita, Mardiah A., Syandri H., 2017 Influence of feeding rate on the growth, feed efficiency and carcass composition of the giant gourami (*Osphronemus goramy*). *Pakistan Journal Zoology* 49(5):1775-1781.
- Asamoah E. K., Nunoo F. K. E., Osei-Asare Y. B., Addo S., Sumaila U. R., 2012 A production function analysis of pond aquaculture in Southern Ghana. *Aquaculture Economic and Management* 16(3):183-201.
- Bhanu Murthy K. V., 2002 Arguing a case for the Cobb-Douglas production function. *Review of Commerce Studies* 20-21(1):75-91.
- Bhuyan A., Akther S., Aktar N., 2011 Present status and fish seed production of the hatcheries of six upazilas of Rajshahi District. *University Journal of Zoology, Rajshahi University* 30:29-32.
- Burmansyah, Muslim, Fitriani M., 2013 [Semi-natural spawning of the climbing perch (*Anabas testudineus*) under different sex ratio]. *Jurnal Akuakultur Rawa Indonesia* 1(1):23-33. [in Indonesian]
- Daudpota A. M., Abbas G., Kalhor H., Hafeez-ur-Rehman M., Ghaffar A., 2016 Effect of feeding frequency on growth performance, feed utilization and body composition of juvenile Nile tilapia, *Oreochromis niloticus* (L) reared in low salinity water. *Pakistan Journal of Zoology* 48(1):171-177.
- Deyab E. S. D. M. S., Hussein E. E. M., 2015 Effects of different feeding rates on growth performance and body composition of red tilapia, *Oreochromis mossambiquense* x *O. niloticus*, fingerlings. *International Journal of Aquaculture* 5(12):1-7.
- Fajrin C. N., Buwono I. D., Sriati, 2012 [Addition of tauge extract in feed for spawning success development of goldfish (*Carassius auratus*)]. *Jurnal Perikanan dan Kelautan* 3(3):51-60. [in Indonesian]
- FAO, 2018 The state of world fisheries and aquaculture, meeting the sustainable development goals. FAO, Rome, 463 pp.
- Ganzon-Naret E. S., 2013 Effects of feeding frequency on growth, survival rate and body composition in sea bass (*Lates calcarifer*) juveniles fed a commercial diet under laboratory condition. *ABAH Bioflux* 5(2):175-182.
- Ganzon-Naret E.S., 2013b Influence of different feeding rate using commercial dry pellets on growth, feed efficiency and survival for hatchery produced sea bass *Lates calcarifer*. *ABAH Bioflux* 5(2):241-248.
- Ghosh A., Mohapatra B. C., Chakrabarti P. P., Hussan A., Das A., 2019 Induced breeding of *Catla catla* carried out at low temperature in FRP carp hatchery of Arunachal Pradesh, India. *Journal of Environmental Biology* 40(3):328-334.

- Guha D., Mukherjee D., 1991 Seasonal cyclical change in the gonadal activity of common carp, *Cyprinus carpio* Linn. *Indian Journal of Fisheries* 38(4):218-223.
- Gujarati D. N., 1988 Basic econometrics. McGraw-Hill, 705 pp.
- Hariani D., Pungky Slamet W. K., 2019 Combination of feed protein level and laserpuncture induction of broodstock catfish (*Clarias* sp.) to increase estrogen, vitellogenin, and egg quality. *EurAsian Journal of BioSciences* 13:769-779.
- Herjayanto M., Carman O., Soelistyowati D. T., 2016 [Spawning behavior, female reproductive potential, and breeding technique optimize of threadfin rainbowfish *Iriatherina wernerii* Meinken, 1974]. *Jurnal Ikhtologi Indonesia* 16(2):171-183. [in Indonesian]
- Huaha T. S., Bukenya J. O., Twinamasiko J., Molnar J., 2011 Profitability analysis of small scale aquaculture enterprises in Central Uganda. *International Journal of Fisheries and Aquaculture* 2(15):271-278.
- Ismi S., Asih Y. N., Kusumawati D., 2013 [Improvement of seed production and quality of grouper by hybridization program]. *Jurnal Ilmu dan Teknologi Keleutan Tropis* 5(2):333-342. [in Indonesian]
- Jia B., St-Hilaire S., Singh K., Gardner I. A., 2016 Farm-level returns and costs of yellow catfish (*Pelteobagrus fulvidraco*) aquaculture in Guangdong and Zhejiang provinces, China. *Aquaculture Reports* 4:48-56.
- Karim M., Keus H. J., Ullah M. H., Kassam L., Phillips M., Beveridge M., 2016 Investing in carp seed quality improvements in homestead aquaculture: lessons from Bangladesh. *Aquaculture* 453:19-30.
- Kholifah S., Antara I. M., Dewi R. K., 2012 [Carp (*Cyprinus carpio*) hatchery business feasibility in "Sari Nadi" Group, Tua village, Marga district, Tabanan regency]. *E-Jurnal Agribisnis & Agrowisata* 1(2):88-95. [in Indonesian]
- Kour S., Masud S., Khan A., 2016 Effect of organic manure and inorganic fertilizer on the growth and proximate composition of common carp, *Cyprinus carpio*. *Journal of Environmental Biology* 37(1):149-153.
- Li X., Yan Q., Xie S., Hu W., Yu Y., Hu Z., 2013 Gut microbiota contributes to the growth of fast-growing transgenic common carp (*Cyprinus carpio* L.). *PLoS ONE* 8(5):e64577.
- Mantau Z., Rawung J. B. M., 2004 [Effective and efficient carp spawning]. *Jurnal Litbang Pertanian* 23(2):68-73. [in Indonesian]
- Ministry of Marine Affairs and Fisheries (MMAF) of Indonesian Republic, 2018a [Fisheries productivity of Indonesia in the forum of Merdeka Barat 9, the Ministry of Communication and Informatics]. 49 pp. [in Indonesian]
- Ministry of Marine Affairs and Fisheries (MMAF) of Indonesian Republic, 2018b [Volume and aquaculture fisheries production value with major commodities and province of 2017]. *Kelautan dan Perikanan dalam Angka 2018*, 590 pp. [in Indonesian]
- Mizuno K., Shimizu-Yamaguchi S., Miura C., Miura T., 2012 Method for efficiently obtaining fertilized eggs from the black scraper *Thamnaconus modestus* by natural spawning in captivity. *Fisheries Science* 78:1059-1064.
- Ogundari K., Akinbogun O. O., 2010 Modeling technical efficiency with production risk: a study of fish farms in Nigeria. *Marine Resource Economics* 25:295-308.
- Onianwa O., Wheelock G., Gyawali B., Gan J., Dubois M., Schelhas J., 2004 An analysis of factors affecting participation behavior of limited resource farmers in agricultural cost-share programs in Alabama. *Journal of Agribusiness* 22(1):17-29.
- Osuagwu E. S., Olaiya E., 2018 Effects of oil spills on fish production in the Niger Delta. *PLoS ONE* 13(10):e0205114.
- Rawat P., Biswas P., Jena A. K., Patel A. B., Pandey P. K., 2019 Effect of dietary incorporation of natural attractants on growth and survival during seed rearing of Indian butter catfish, *Ompok bimaculatus*. *Journal of Environmental Biology* 40(4):661-667.
- Saikia S. K., Das D. N., 2009 Feeding ecology of common carp (*Cyprinus carpio* L.) in a rice-fish culture system of the Apatani plateau (Arunachal Pradesh, India). *Aquatic Ecology* 43(2):559-568.

Commented [indra2]: I was not able to find this reference anywhere on the Internet

- Saravi Saeedi S. S., Karami S., Karami B., Shokrzadeh M., 2009 Toxic effects of cobalt chloride on hematological factors of common carp (*Cyprinus carpio*). *Biological Trace Element Research* 132:144-152.
- Shivakumar M., Bala S., Rajanna C., Naveenkumar B. T., 2014 Economics of seed rearing and farming of carps. *International Journal of Fisheries and Aquatic Studies* 2(1):42-45.
- Singh K., 2008 Farm specific economic efficiency of fish production in South Tripura District: a stochastic frontier approach. *Indian Journal of Agricultural Economics* 63 (4):598-613.
- Soekartawi, 1994 [Cobb-Douglas function-based production economic theory]. PT RajaGrafindo Persada, Jakarta, 258 pp. [in Indonesian]
- Sofia L. A., Muhammad S., Hanani N., Fatah L., 2013 Simulation model of household economy on production and welfare of catfish (*Pangasianodon hypophthalmus*) farmer in Banjar Minapolitan, South Kalimantan. *IOSR Journal of Environmental Science, Toxicology and Food Technology* 7(6):1-8.
- Solomon S. G., Tiarniyu L. O., Okomoda V. T., Adaga K., 2016 Nutrient profile of commercial aqua-feeds under different storage conditions. *International Journal of Aquaculture* 6(12):1-11.
- Statistic Center of Tapin Regency, 2019 [Tapin Regency in figures 2019]. Statistic Center of Tapin Regency, South Kalimantan, 451 pp. [in Indonesian]
- Syaifudin M., Aliah R. S., Muslim, Sumantadinata K., 2007 [Relation between broodstock number and spawning frequency and egg production of humpback grouper (*Cromileptes altivelis*)]. *Jurnal Akuakultur Indonesia* 6(2):191-196. [in Indonesian]
- United Nations, 2019 World Population Prospects 2019. Population Division. Department of Economic and Social Affairs. United Nations. <https://www.un.org/development/desa/publications/world-population-prospects-2019-highlights.html>.
- Valdebenito I. I., Gallegos P. C., Effer B. R., 2013 Gamete quality in fish: evaluation parameters and determining factors. *Zygote* 23:177-197.
- Vergos K., Christopoulos A., Krystallidis P., Papandroni O., 2010 Economies of scale and concentration in the Greek and the Norwegian aquaculture industry: an empirical study. *International Journal of Business Management and Economic Research* 1(1):70-78.
- Wang Q., Li Z., Gui J. F., Liu J., Ye S., Yuan J., De Silva S. S., 2017 Paradigm changes in freshwater aquaculture practice in China: moving towards achieving environmental integrity and sustainability. *Ambio* 47(4):410-426.
- Xu J., Ji P., Zhao Z., Zhang Y., Feng J., Wang J., Li J., Zhang X., Zhao L., Liu G., Xu P., Sun X., 2012 Genome-wide SNP discovery from transcriptome of four common carp strains. *PLoS ONE* 7(10):e48140.
- Yulinda E., 2012 [Financial analysis of catfish (*Clarias gariepinus*) hatchery business in Lembah Sari, Rumbai Pesisir district, Pekanbaru, the Province of Riau]. *Jurnal Perikanan dan Kelautan* 17(1):38-55. [in Indonesian]

Received: 29 February 2020. Accepted: 26 March 2020. Published online: 05 April 2020.

Authors:

Bambang Poerwanto, Magister Program of Fisheries Science, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: poerwantobambang2@gmail.com

Idiannor Mahyudin, Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: dr\_idiannor@yahoo.co.id

Leila Ariyani Sofia, Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: leila.ariyani@ulm.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Poerwanto B., Mahyudin I., Sofia L. A., 2020 The analysis of factors influencing carp (*Cyprinus carpio*) seed production in Bungur District, Tapin Regency, South Kalimantan. *AAAL Bioflux* 13(2):804-812.

Search all conversations



Active



367 of 415 < >


On Tuesday, April 7, 2020, 6:07:32 AM wrote:

Dear Dr. Gavrioloie,  
I herewith send you the revised final form of Poerwanto et al. References were also added. I put all revisions in yellow. Thank you

Sincere,



REVISI 7 APRIL 2020



## The analysis of factors influencing carp (*Cyprinus carpio*) seed production in Bungur District, Tapin Regency, South Kalimantan

<sup>1</sup>Bambang Poerwanto, <sup>2</sup>Idianoor Mahyudin, <sup>2</sup>Leila A. Sofia

<sup>1</sup> Magister Program of Fisheries Science, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia; <sup>2</sup> Fisheries Services of Tapin Regency, Jl. Jenderal Sudirman 2.5 km, Rantau, South Kalimantan; <sup>3</sup> Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia. Corresponding author: L. A. Sofia, leila.ariyani@ulm.ac.id

**Abstract.** Seed production has become one of the factors supporting the continuity of fish farming activity. Carp, *Cyprinus carpio* is one of the largest freshwater fish productions that support the human food needs in the world. This study aims to know several factors affecting carp seed production. The study locality was purposively selected in Bungur District, Tapin Regency, South Kalimantan, and samples were collected through census from 42 carp farmers. The primary data were descriptively quantitatively analyzed using multiple regression. Results showed that there were 4 factors affecting the seed production, number of male spawners, feed amount for fry, spawning frequency, and the use frequency of the same female spawners. Carp seed production development could be done by considering the supply of enough number and good quality spawners, good protein feeding and efficient feeding technique, and spawning recovery time of the female spawners.

**Key Words:** fish farmer, pond, stocking density, spawners, feeding.

**Introduction.** World human population in 2019 reached 7.7 billion people and is estimated to be 9.7 billion in 2050 (United Nations 2019). Increase in human population causes increase in world food needs including fisheries products. In 2016, the world fisheries production also reached 170.9 million tons, from which 151.2 million tons were used for human consumption and 19.7 million tons for non consumption need. This production comprises fishing fisheries and aquaculture. Nevertheless, the world fishing fisheries production has declined from 2011 to 2016 from 92.2 to 90.9 million tons, while world aquaculture production has significantly risen from 61.8 to 80.0 million tons in the same period (FAO 2018). Systematic fishing in tremendously high numbers, weak reporting, and other complex pressures added with ecosystem change consequences have resulted in decline in fishing fisheries production (Allan et al 2005). Hence, fish farming becomes a very important sector that plays important roles in world food security due to increasing human population and food needs (Wang et al 2017).

Indonesia as the fourth highest human population country, 267.6 million people (United Nations 2019) required fish consumption up to 43.94 kg capita<sup>-1</sup> in 2016. This need was supported by 16.68 million tons of aquaculture and 6.83 million tons of fishing fisheries. Total fisheries production of Indonesia in 2016 reached 23.51 million tons and has contributed to 13.75% of world fisheries production (Ministry of Marine Affairs and Fisheries (MMAF) of RI 2018a). The need of Indonesian fisheries products can be achieved by 70% from aquaculture. This condition makes aquaculture be very crucial. There are several fish commodities developed in Indonesia, one of which is carp, *Cyprinus carpio*. This species becomes the fourth largest freshwater aquaculture commodity in Indonesia with a volume of 316,646 tons in 2017 (MMAF RI 2018b). This species also became the third largest fish culture commodity in the world (Xu et al 2012; FAO 2018) and one of the fish species that can reach high biomass and its distribution

has been developed by human due to its important economic value (Arteni & Rosca 2010; Saikia & Das 2009).

The magnitude of carp culture commodity needs fish seed availability as important prerequisite of carp cultivation (Bhuyan et al 2011). The supply of good quality fish seeds is necessary for faster development of aquaculture industry. Hatchery efforts are also needed to reduce the dependence of aquaculture upon the use of wild seeds (Ghosh et al 2019). The development of carp seed quality has been conducted through various techniques, such as application of certain treatment to carp seeding production factors. The treatment could be implemented by previously knowing which production factor yields significant effect on the seeding quantity and quality (Mizuno et al 2012; Karim et al 2016; Hariani & Pungky Slamet 2019). Studies on the effect of various factors on carp condition have been done a lot, such as how the intestinal microbes affect the fish growth (Li et al 2013) up to how the toxin effect, such as cobalt chloride, on the hematological condition of carp (Saravi et al 2009). Nevertheless, there are very few studies on carp seeding quality and quantity development. Nowadays, the development of good fish seed quality production is needed in order to meet the seed demand in aquaculture and make the public hatchery be a profitable business (Ismi et al 2013).

Tapin Regency, South Kalimantan province, is one of the regions that run carp culture activities during 2013-2017 with mean production level up to 13,198,600 ind.yr<sup>-1</sup> (Animal Husbandary and Fisheries Service of Tapin Regency 2018). Bungur District is one of the largest carp seeding localities in Tapin Regency. The seed production must be raised due to increasing human population and food demand, fisheries product in particular. Increased fish seed production is expected to be able to maintain the sustainability of the carp culture industry. Developing the carp seeding production requires a right strategy. This study aims to know factors affecting the carp seed production in Bungur District, Tapin Regency, the province of South Kalimantan.

## Material and Method

**Study site description.** The work was carried out from January to May 2019 in Bungur District, Tapin Regency, South Kalimantan province with an area of 148.98 km<sup>2</sup>. Bungur District is the largest aquaculture area in Tapin Regency, where 75% is located in Bungur District, and most of it is used for fish hatchery business, including carp seeding (Statistic Center of Tapin Regency 2019). This study site selection was done since Bungur District is the fish seeding center of Tapin Regency with the biggest aquaculture area and number of fish farmers.

**Population and samples.** Fish hatchery farmers in Shabah village and Kalumpang village, Bungur District are 42 people. All population members were taken as research samples through census method.

**Data analyses.** Factors influencing carp seed production were analyzed using descriptively quantitative method and OLS multiple regression with Cobb-Douglas-typed production function approach. This function has been widely used in the field of agricultural economy to show the relationship between number of inputs and outputs (Singh 2008; Osuagwu & Olaifa 2018). It was selected for the following potentials: (1) the Cobb-Douglas (C-D) function estimation will give the regression coefficient that indicates the elasticity, (2) the elasticity indicates the return to scale rate, and (3) the calculation is relatively easier than other functions (Soekartawi 1994; Bhanu Murthy 2002; Vergos et al 2010).

The model applied in this study was as follows:

$$Y = a X_1^{b_1} X_2^{b_2} X_3^{b_3} X_4^{b_4} X_5^{b_5} X_6^{b_6} X_7^{b_7} X_8^{b_8} X_9^{b_9} X_{10}^{b_{10}} D_1 e$$

$$\ln Y = \ln a + b_1 \ln X_1 + b_2 \ln X_2 + b_3 \ln X_3 + b_4 \ln X_4 + b_5 \ln X_5 + b_6 \ln X_6 + b_7 \ln X_7 + b_8 \ln X_8 + b_9 \ln X_9 + b_{10} \ln X_{10} + D_1 + e$$

where a is constant, b<sub>1</sub>-b<sub>10</sub> are regression coefficients, Y is the fish seeding production (ind year<sup>-1</sup>); X<sub>1</sub> is pond size (m<sup>2</sup>), X<sub>2</sub> is number of female spawners (ind), X<sub>3</sub> is number of

males (ind),  $X_4$  is feed volume for spawners (kg),  $X_5$  is total feed volume for fry (kg),  $X_6$  is amount of manure fertilizer (pack),  $X_7$  is spawning frequency (times),  $X_8$  is use frequency of the same female spawners,  $X_9$  is number of labor used (HOK),  $X_{10}$  is fish farmer's experience (year),  $D_1$  (dummy variable) is trained fish farmer = 1, untrained fish farmer = 0, and  $e$  is error term.

The precision of the C-D production function model was tested with classic assumption tests which include test normality, multicollinearity, and heteroskedasticity, then test goodness of Fit ( $R^2$ ), simultaneous variable coefficient test (F test), and partial variable coefficient test (t-test) were done. If all requirements are fulfilled, the model could be used for prediction (Gujarati 1988). The regression coefficients of all inputs used in the model were *a priori* assumed as positive (Ogundari & Akinbogun 2010; Asamoah et al 2012), except the frequency of the same female spawners utilization. Dummy variable coefficient, trained fish farmers  $D_1$ , was assumed to be positive (Onianwa et al 2004; Hyuha et al 2011).

## Results

**Seeding activity description.** The Public Hatchery Unit has been done by the farming group in Bungur District for 8 years in average. The hatchery business run for 1-5 years was 48%, 6-10 years was 24%, and more than 10 years was 29%. The construction of carp seeding pond used by fish farmers in Bungur District is soil ponds with area around 1,490-18,212  $m^2$  and number of ponds managed were 4-29 units hatchery<sup>-1</sup> (Table 1) with total labors of 12-306 working days. Number of labors in each hatchery was 2-3 people depending upon number of spawning ponds operated. The pond localities are generally near the irrigation route so that the ponds are close enough to the water source. Before use, the fish farmers have fertilized the pond with manure. The organic fertilizer is effective to grow natural food that the productivity and fish growth can be increased (Kour et al 2016). The fertilizer applied in the study site was about 160-2,000 bags  $yr^{-1}$  hatchery<sup>-1</sup> or about 14 bags  $m^{-2}$ .

Fish spawner is major production factor in seed production, in either quality or quantity. Both females and males were prepared by the fish farmers. The female spawner candidates were 1.5-3 years old with individual weight of about 2-4 kg, the males were older than 6 months old with an individual weight of 0.2-1 kg. In general, number of spawners prepared by the fish farmers were 4-53 females and 14-178 males per business unit. The female and male spawners ratio was 2:8 each spawning.

Table 1  
Carp hatchery production and input production description in Public Hatchery Unit in Bungur district

Description	Unit	Value		
		Maximum	Minimum	Average
Fish seeding production	ind $yr^{-1}$	2,080,000	100,000	340,520
Total pond size	$m^2$	18,212	1,490	6,874
Number of ponds	unit	29	4	9
Pond size	$m^2$ unit <sup>-1</sup>	1,197	298	782
Number of female spawners	ind	53	4	16
Weight of female spawners	kg ind <sup>-1</sup>	4	2	3
Number of male spawners	ind	178	14	61
Weight of male spawners	kg ind <sup>-1</sup>	1	0.2	0.34
Feed volume of spawners	kg $yr^{-1}$	840	40	318
Feed volume for fish fry	kg $yr^{-1}$	1,560	200	647
Volume of manure fertilizer	bags $yr^{-1}$	2,000	160	651
Spawning frequency	no $yr^{-1}$	32	4	7
Use frequency of the same female spawners	no $yr^{-1}$	6	2	4
Number of labor used	working days	306	12	83
Fish farmer's experience	year	18	1	8

Total time utilized for spawning in one-year is about 8 months. One production cycle requires more or less 60 days, and it could be done 1-2 times mo<sup>-1</sup> or 4-32 times yr<sup>-1</sup>. To maintain the quality of the produced seed, the frequency use of the same female spawners was 2-6 times yr<sup>-1</sup>, and every year the new spawners are prepared to replace the old ones. Total number of seeds produced by the hatchery were 100,000-2,080,000 ind yr<sup>-1</sup>.

These are the fry size ranges that are commonly prepared by the hatchery to the fish farmers under different selling price, 2-3 cm ( $\pm 15$  days), 3-5 cm ( $\pm 20$  days), and 5-8 cm total length that need about  $\pm 30$  days. In this stage, the seeds were given fine pellet with total feed weight of 200-1560 kg yr<sup>-1</sup> per business unit. Similarly, the spawners were fed with artificial feed as much as 40-840 kg yr<sup>-1</sup> per business unit.

**Financial analysis.** Profit is the main goal of a business. The amount of profit gained will determine the fish farmer's decision making in the business production planning. Carp hatchery business needs an investment of IDR 2.9 million – IDR 254.23 million (Table 2) depending upon land size, number of ponds used, and other supporting facilities, such as keeping house and storage, net/happa, kakaban/egg substrate, bucket, sorting box, and scoop net. Each business unit required the operational costs of IDR 29.69 million – IDR 159.34 million yr<sup>-1</sup> with the highest in feed preparation for the seeds, IDR 1.05 million – IDR 94 million yr<sup>-1</sup> depending upon the seed numbers produced. The additional feed was administered at the life stage of about 7-day old larvae (with yolk sac) up to be ready for sale.

The second highest operational cost is labor's wages, IDR 16.4 million – IDR 27 million. It was allocated for pond preparation or processing, dyke improvement, and feeding. The lowest cost was recorded in male spawner supply, IDR 135,000 – IDR 1,750,000 yr<sup>-1</sup>. Previous studies found no enough difference, with total costs of IDR. 66,262,510.89, approximately 31% is used for feed costs (Shivakumar et al 2014). This finding is in agreement with Kholifah et al (2012) that carp hatchery business has good prospectus with a profit of IDR. 7,329,301.90 cycle<sup>-1</sup>, but different from that of Yulinda (2012) concerning the business financial analysis of catfish *Clarias gariepinus* with much lower profit, averagely IDR. 1,745,194 cycle<sup>-1</sup>. It could result from higher market price of carp fry than that of catfish.

Table 2  
Investment, operational costs, and profit of carp hatchery unit in Bungur District

Description	Unit	Value		
		Maximum	Minimum	Mean
Investment (total):	IDR	<b>254,225,000</b>	<b>2,935,000</b>	<b>46,256,475</b>
Pond	IDR	200,000,000	1,800,000	32,222,500
Other supporting facilities	IDR	54,225,000	1,135,000	14,033,975
Operational costs (total):	IDR yr <sup>-1</sup>	<b>201,600,000</b>	<b>21,995,000</b>	<b>61,318,505</b>
Female spawner preparation	IDR yr <sup>-1</sup>	7,500,000	480,000	2,139,500
Male spawner preparation	IDR yr <sup>-1</sup>	1,750,000	135,000	621,038
Spawner's feed preparation	IDR yr <sup>-1</sup>	22,800,000	900,000	6,181,167
Seed feed preparation	IDR yr <sup>-1</sup>	94,000,000	1,050,000	19,699,750
Manure fertilizer	IDR yr <sup>-1</sup>	33,500,000	2,680,000	10,445,300
Other input preparation	IDR yr <sup>-1</sup>	14,750,000	350,000	3,081,750
Labor's wages	IDR yr <sup>-1</sup>	27,300,000	16,400,000	19,150,000
Revenue and profit:				
Revenue	IDR yr <sup>-1</sup>	268,000,000	57,600,000	120,215,750
Business profit	IDR yr <sup>-1</sup>	66,400,000	35,605,000	58,897,246

The revenue (from seed sales) obtained by a hatchery business unit ranged from IDR 57.6 million to IDR 268 million. The amount of revenue is dependent upon the number of production, seed size, and selling price. The larger the seed size is, the more expensive the seed will be. Under an agreement among the fish farmers, the selling price was IDR

250 ind<sup>-1</sup> for 2-3 cm total length seed, IDR 300 ind<sup>-1</sup> for 3-5 cm total length seed, and IDR 350 ind<sup>-1</sup> for 5-8 cm total length seed, respectively. Therefore, the business profit varied as well from IDR 35,6 to IDR 66,4 million yr<sup>-1</sup>.

**Production and production factors.** The effect of production factors on the carp seed production showed that several production factors (input) involved in the production function model gave the consistent values with the theoretical estimate. The production factors of positive value and significant effect on production were number of male spawners, feed volume for seeds, and spawning frequency done by the fish farmers. However, the use of the same female spawners for several spawning activities during the reproductive cycle will cause seed production decline as well (Table 3).

Table 3  
Factors influencing the production of common carp hatchery

Variable	Coefficient	SE	t-Statistic	P value
Intercept	7.543	1.264	5.945	0.000
Total pond size (m <sup>2</sup> )	0.059	0.123	0.477	0.637
No. female spawners (ind.)	0.065	0.106	0.620	0.540
No. male spawners (ind.)	0.343	0.116	2.951**	0.006
Feed volume for spawners (kg)	-0.071	0.088	-0.813	0.423
Feed volume for seeds (kg)	0.381	0.138	2.760**	0.010
No. manure fertilizer (bag)	0.094	0.103	0.920	0.365
Spawning frequency (times)	0.910	0.110	8.241**	0.000
Use frequency of the same female spawners (times)	-0.324	0.139	-2.338**	0.026
Labor's working days	-0.204	0.113	-1.802	0.082
Fish farmer's experience (year)	-0.048	0.079	-0.608	0.548
Trained fish farmers (dummy)	0.112	0.139	0.808	0.425
R-squared		0.828		
Adjusted R-squared		0.765		
Standard error		0.321		
Observations		42		
F-statistic		13.154		
Prob (F-statistic)		0.000		
Durbin-Watson		2.190		

Note: double asterisks (\*\*) denote significance at 1% level, respectively.

Hence, to increase the seed production, the fish farmers could increase several inputs and reduce the use frequency of the same female spawners to give them some time to recover and to have perfect maturation.

**Discussion.** Fish gamete quality, male or female, is determined by a number of factors, such as age, management, food, chemical and physical factors, water quality, and so on, that will impact the survivorship of the embryos, larvae, and/or fry in short or long term (Valdebenito et al 2013). Fish spawners are a major production factor in the hatchery business. The use of higher number of spawners in spawning activity will positively influence the number of eggs produced. The more the spawners are spawned, the higher the spawning frequencies could be done and the higher the number of eggs will be produced (Syaifudin et al 2007). The female carp spawner appropriate for spawning should be 1.5-3 years old with minimum weight of 1.5 kg ind<sup>-1</sup>, while the male should be at least 6 months old with minimum weight of 0.5 kg ind<sup>-1</sup> (Mantau & Rawung 2004; Aliniya et al 2013). The weight ratio of female and male is 1:1 (kg m<sup>-2</sup>) meaning that a 2 kg ind<sup>-1</sup> female could be balanced by 3 males of 600-700 g ind<sup>-1</sup>.

Some fish hatchery businesses showed that female-male ratio became factor determining the quality and quantity of seeds produced. For instance, in rainbow fish (*Iriatherina wernerii*) spawning, it is known that the use of sex ratio with 1:3 of total

weight is more efficient than 1:1 and 1:2 ratios. It is possible that more male spawners will give higher opportunity to fertilize the eggs. Also, the seed quality produced at the spawner sex ratios of 1:2 and 1:3 gave 5-day old seeds with good survival rate (Herjayanto et al 2016). The climbing perch *Anabas testudineus* at 1:4 ratio spawners yields good fecundity (Burmansyah et al 2013). However, the female-male ratio practiced by the fish farmers in the study site indicated that total weight of male spawner was still below the total weight of female spawner. Thus, number and weight of male spawners need to be added in order to increase the seed production.

*C. carpio* naturally showed 2 clear reproductive cycles per year. Each reproductive cycle is divided into 4 phases. Males have growth phase, maturation, spawning, and resting, while females have preparatory phase, pre-spawning, spawning, and post-spawning (Guha & Mukherjee 1991). At 2 years of age, carp has fecundity of 523,500 eggs and at 3 years of age, the fecundity reaches 657,701 eggs (Aliniya et al 2013). Thus, the more the number of spawners spawned are, the more the number of eggs and eventually more larvae are produced. The availability of spawners spawned will also interchangeably give the opportunity to the spawners to have perfect reproductive cycle. The spawning frequency using the same female spawners gave impact on the seed growth. This study found that the seeds obtained from the spawners often used had lower body mass and shorter size. Also, the larvae produced from the often used spawners looked paler than those from the less used spawners.

Commercial feasibility of intensive fish culture depends on market demand and production costs. The largest part of the production costs is feed, so that it is important to know the optimal feeding and the right feed management strategy implementation (Jia et al 2016). For instance, the right feeding frequency can reduce the production costs and prevent water quality degradation (Daudpota et al 2016). Higher feeding frequency gives positive impact on the fish growth and makes the feed conversion ratio value be more efficient (Ganzon-Naret 2013a). Besides that, the percent of feeding becomes important factor for fish growth and the percentage of optimal feeding determines the aquaculture success (Deyab & Hussein 2015; Aryani et al 2017). The percent feed amount given will influence the fish growth rate as weight increment (Ganzon-Naret 2013b; Sofia et al 2013).

Feed is an important component for the spawners in order to produce good quality eggs and seeds in high number. The fish spawners spawned at the high frequency need good quality feed. Specific additional feeding also affects the aquaculture production. For example, addition of tauge extract in gold fish *Carrasius auratus* feed gives faster spawning cycle (Fajrin et al 2012). Addition of probiotic to the feed can also increase body weight and fish endurance, so that the production rises and feed costs decline (Aitkaliyeva et al 2019). Besides feeding technique, the fish farmers should understand how to store the feed before use. According to Solomon et al (2016), storage in closed condition can prevent the nutritional damage of the feed.

Carp seeds produced need different feed treatments with life stages divided into 3 nursing phases. First, this stage was done for 7-day old larvae (0.6-0.7 cm total length) at a stocking density of 100 ind m<sup>-2</sup> and in a pond of about 60 cm deep. At this stage, they were fed with the aquatic biota living in the pond and fine pellet. In about 15 days of rearing, the fish grew to 1-3 cm total length. Second, this stage was conducted at the stocking density of about 50 ind m<sup>-2</sup> in about 20 days of rearing up to producing 3-5 cm total length fish. Third, the stocking density of this stage was about 25 ind m<sup>-2</sup> at rearing period of about 30 days until 5-8 cm total length fish were obtained. After these nursing phases, the seeds are ready to sell, and they are sold when reaching the individual size of 3-5 cm and 5-8 cm long.

So far, feeding in each nursing phase still relied upon the artificial feed so that the operational cost for the fry feed was very high. On the other hand, the economic return of the larvae produced in the hatchery is more important than the fry growth. Therefore, alternative additional feed and feed cost efficiency need to be applied in order to increase the survivorship. One of these is bloodworm (*Tubifex* sp.) supplement. According to Rawat et al (2019), 5% tubifex supplement can be used to increase the fish survivorship, especially in indoor seed rearing.

**Conclusions.** There were 4 factors affecting the production of *C. carpio* hatchery, three of which had positive effect on seed production, namely the number of male spawners, the amount of feed for the fry, and the spawning frequency. While the use frequency of female spawners during the production cycle negatively impact the seed production. This finding suggested that the fish farmers need to pay attention on the availability of sufficient number of good fish spawners, good protein feeding, efficient feeding technique, and longer post-spawning recovery time for female spawners.

**Acknowledgements.** We would greatly appreciate the Animal Husbandry and Fisheries Services of Tapin Regency, the Fisheries Consultant for field information and guidance in this study. We would also thank the fish farmers in the Public Hatchery Unit in Bungur District, Tapin Regency for their collaboration.

## References

- Aitkaliyeva A. A., Alpeisov S. A., Isbekov K. B., Assylbekova S. Z., Badryzlova N. S., 2019 Influence of different feeds and feed additives on fish-breeding and biological indicators at rearing rainbow trout. *EurAsian Journal of BioSciences* 13(1):437-442.
- Aliniya M., Nezami S., Khara H., Noveiri S. B., Dadras H., Fallah Z., 2013 Influence of age of common carp (*Cyprinus carpio*) broodstock on reproductive traits and fertilization. *Turkish Journal of Fisheries and Aquatic Sciences* 13:19-25.
- Allan J. D., Abeli R., Hogan Z., Revenga C., Taylor B. W., Welcomme R. L., Winemiller K., 2005 Overfishing of inland waters. *BioScience* 55(12):1041-1051.
- Animal Husbandary and Fisheries Service of Tapin Regency, 2018 [Annual report on fisheries of Tapin Regency, South Kalimantan]. 74 pp. [in Indonesian]
- Arteni M. O., Rosca I., 2010 Some morphological features of carp (*Cyprinus carpio* Linnaeus, 1758) in different stages of development. *ABAH Bioflux* 2(1):47-52.
- Aryani N., Azrita, Mardiah A., Syandri H., 2017 Influence of feeding rate on the growth, feed efficiency and carcass composition of the giant gourami (*Osphronemus goramy*). *Pakistan Journal Zoology* 49(5):1775-1781.
- Asamoah E. K., Nunoo F. K. E., Osei-Asare Y. B., Addo S., Sumaila U. R., 2012 A production function analysis of pond aquaculture in Southern Ghana. *Aquaculture Economic and Management* 16(3):183-201.
- Bhanu Murthy K. V., 2002 Arguing a case for the Cobb-Douglas production function. *Review of Commerce Studies* 20-21(1):75-91.
- Bhuyan A., Akther S., Aktar N., 2011 Present status and fish seed production of the hatcheries of six upazilas of Rajshahi District. *University Journal of Zoology, Rajshahi University* 30:29-32.
- Burmansyah, Muslim, Fitriani M., 2013 [Semi-natural spawning of the climbing perch (*Anabas testudineus*) under different sex ratio]. *Jurnal Akuakultur Rawa Indonesia* 1(1):23-33. [in Indonesian]
- Daudpota A. M., Abbas G., Kalhoro I. B., Shah S. S. A. Shah, Kalhoro H., Hafeez-ur-Rehman M., Ghaffar A., 2016 Effect of feeding frequency on growth performance, feed utilization and body composition of juvenile Nile tilapia, *Oreochromis niloticus* (L) reared in low salinity water. *Pakistan Journal of Zoology* 48(1):171-177.
- Deyab E. S. D. M. S., Hussein E. E. M., 2015 Effects of different feeding rates on growth performance and body composition of red tilapia, *Oreochromis mossambiquise* x *O. niloticus*, fingerlings. *International Journal of Aquaculture* 5(12):1-7.
- Fajrin C. N., Buwono I. D., Sriati, 2012 [Addition of tauge extract in feed for spawning success development of goldfish (*Carassius auratus*)]. *Jurnal Perikanan dan Kelautan* 3(3):51-60. [in Indonesian]
- FAO, 2018 The state of world fisheries and aquaculture, meeting the sustainable development goals. FAO, Rome, 463 pp.
- Ganzon-Naret E. S., 2013 Effects of feeding frequency on growth, survival rate and body composition in sea bass (*Lates calcarifer*) juveniles fed a commercial diet under laboratory condition. *ABAH Bioflux* 5(2):175-182.

- Ganzon-Naret E.S., 2013b Influence of different feeding rate using commercial dry pellets on growth, feed efficiency and survival for hatchery produced sea bass *Lates calcarifer*. *ABAH Bioflux* 5(2):241-248.
- Ghosh A., Mohapatra B. C., Chakrabarti P. P., Hussan A., Das A., 2019 Induced breeding of *Catla catla* carried out at low temperature in FRP carp hatchery of Arunachal Pradesh, India. *Journal of Environmental Biology* 40(3):328-334.
- Guha D., Mukherjee D., 1991 Seasonal cyclical change in the gonadal activity of common carp, *Cyprinus carpio* Linn. *Indian Journal of Fisheries* 38(4):218-223.
- Gujarati D. N., 1988 Basic econometrics. McGraw-Hill, 705 pp.
- Hariani D., Pungky Slamet W. K., 2019 Combination of feed protein level and laserpuncture induction of broodstock catfish (*Clarias* sp.) to increase estrogen, vitellogenin, and egg quality. *EurAsian Journal of BioSciences* 13:769-779.
- Herjayanto M., Carman O., Soelistyowati D. T., 2016 [Spawning behavior, female reproductive potential, and breeding technique optimize of threadfin rainbowfish *Iriatherina wernerii* Meinken, 1974]. *Jurnal Ikhtologi Indonesia* 16(2):171-183. [in Indonesian]
- Hyuha T. S., Bukenya J. O., Twinamasiko J., Molnar J., 2011 Profitability analysis of small scale aquaculture enterprises in Central Uganda. *International Journal of Fisheries and Aquaculture* 2(15):271-278.
- Ismi S., Asih Y. N., Kusumawati D., 2013 [Improvement of seed production and quality of grouper by hybridization program]. *Jurnal Ilmu dan Teknologi Keleutan Tropis* 5(2):333-342. [in Indonesian]
- Jia B., St-Hilaire S., Singh K., Gardner I. A., 2016 Farm-level returns and costs of yellow catfish (*Pelteobagrus fulvidraco*) aquaculture in Guangdong and Zhejiang provinces, China. *Aquaculture Reports* 4:48-56.
- Karim M., Keus H. J., Ullah M. H., Kassam L., Phillips M., Beveridge M., 2016 Investing in carp seed quality improvements in homestead aquaculture: lessons from Bangladesh. *Aquaculture* 453:19-30.
- Kholifah S., Antara I. M., Dewi R. K., 2012 [Carp (*Cyprinus carpio*) hatchery business feasibility in "Sari Nadi" Group, Tua village, Marga district, Tabanan regency]. *E-Jurnal Agribisnis & Agrowisata* 1(2):88-95. <https://ojs.unud.ac.id/index.php/JAA/article/view/4862> [in Indonesian]
- Kour S., Masud S., Khan A., 2016 Effect of organic manure and inorganic fertilizer on the growth and proximate composition of common carp, *Cyprinus carpio*. *Journal of Environmental Biology* 37(1):149-153.
- Li X., Yan Q., Xie S., Hu W., Yu Y., Hu Z., 2013 Gut microbiota contributes to the growth of fast-growing transgenic common carp (*Cyprinus carpio* L.). *PLoS ONE* 8(5):e64577.
- Mantau Z., Rawung J. B. M., 2004 [Effective and efficient carp spawning]. *Jurnal Litbang Pertanian* 23(2):68-73. [in Indonesian]
- Ministry of Marine Affairs and Fisheries (MMAF) of Indonesian Republic, 2018a [Fisheries productivity of Indonesia in the forum of Merdeka Barat 9, the Ministry of Communication and Informatics]. 49 pp. [in Indonesian]
- Ministry of Marine Affairs and Fisheries (MMAF) of Indonesian Republic, 2018b [Volume and aquaculture fisheries production value with major commodities and province of 2017]. *Kelautan dan Perikanan dalam Angka 2018*, 590 pp. [in Indonesian]
- Mizuno K., Shimizu-Yamaguchi S., Miura C., Miura T., 2012 Method for efficiently obtaining fertilized eggs from the black scraper *Thamnaconus modestus* by natural spawning in captivity. *Fisheries Science* 78:1059-1064.
- Ogundari K., Akinbogun O. O., 2010 Modeling technical efficiency with production risk: a study of fish farms in Nigeria. *Marine Resource Economics* 25:295-308.
- Onianwa O., Wheelock G., Gyawali B., Gan J., Dubois M., Schelhas J., 2004 An analysis of factors affecting participation behavior of limited resource farmers in agricultural cost-share programs in Alabama. *Journal of Agribusiness* 22(1):17-29.
- Osuagwu E. S., Olaifa E., 2018 Effects of oil spills on fish production in the Niger Delta. *PLoS ONE* 13(10):e0205114.

Commented [indra1]: I was not able to find this reference anywhere on the Internet



- Rawat P., Biswas P., Jena A. K., Patel A. B., Pandey P. K., 2019 Effect of dietary incorporation of natural attractants on growth and survival during seed rearing of Indian butter catfish, *Ompok bimaculatus*. *Journal of Environmental Biology* 40(4):661-667.
- Saikia S. K., Das D. N., 2009 Feeding ecology of common carp (*Cyprinus carpio* L.) in a rice-fish culture system of the Apatani plateau (Arunachal Pradesh, India). *Aquatic Ecology* 43(2):559-568.
- Saravi Saeedi S. S., Karami S., Shokrzadeh M., 2009 Toxic effects of cobalt chloride on hematological factors of common carp (*Cyprinus carpio*). *Biological Trace Element Research* 132:144-152.
- Shivakumar M., Bala S., Rajanna C., Naveenkumar B. T., 2014 Economics of seed rearing and farming of carps. *International Journal of Fisheries and Aquatic Studies* 2(1):42-45.
- Singh K., 2008 Farm specific economic efficiency of fish production in South Tripura District: a stochastic frontier approach. *Indian Journal of Agricultural Economics* 63 (4):598-613.
- Soekartawi, 1994 [Cobb-Douglas function-based production economic theory]. PT RajaGrafindo Persada, Jakarta, 258 pp. [in Indonesian]
- Sofia L. A., Muhammad S., Hanani N., Fatah L., 2013 Simulation model of household economy on production and welfare of catfish (*Pangasianodon hypophthalmus*) farmer in Banjar Minapolitan, South Kalimantan. *IOSR Journal of Environmental Science, Toxicology and Food Technology* 7(6):1-8.
- Solomon S. G., Tiamiyu L. O., Okomoda V. T., Adaga K., 2016 Nutrient profile of commercial aqua-feeds under different storage conditions. *International Journal of Aquaculture* 6(12):1-11.
- Statistic Center of Tapin Regency, 2019 [Tapin Regency in figures 2019]. Statistic Center of Tapin Regency, South Kalimantan, 451 pp. [in Indonesian]
- Syaifudin M., Aliah R. S., Muslim, Sumantadinata K., 2007 [Relation between broodstock number and spawning frequency and egg production of humpback grouper (*Cromileptes altivelis*)]. *Jurnal Akuakultur Indonesia* 6(2):191-196. [in Indonesian]
- United Nations, 2019 World Population Prospects 2019. Population Division. Department of Economic and Social Affairs. United Nations. <https://www.un.org/development/desa/publications/world-population-prospects-2019-highlights.html>.
- Valdebenito I. I., Gallegos P. C., Effer B. R., 2013 Gamete quality in fish: evaluation parameters and determining factors. *Zygote* 23:177-197.
- Vergos K., Christopoulos A., Krystallidis P., Papandroni O., 2010 Economies of scale and concentration in the Greek and the Norwegian aquaculture industry: an empirical study. *International Journal of Business Management and Economic Research* 1(1):70-78.
- Wang Q., Li Z., Gui J. F., Liu J., Ye S., Yuan J., De Silva S. S., 2017 Paradigm changes in freshwater aquaculture practice in China: moving towards achieving environmental integrity and sustainability. *Ambio* 47(4):410-426.
- Xu J., Ji P., Zhao Z., Zhang Y., Feng J., Wang J., Li J., Zhang X., Zhao L., Liu G., Xu P., Sun X., 2012 Genome-wide SNP discovery from transcriptome of four common carp strains. *PLoS ONE* 7(10):e48140.
- Yulinda E., 2012 [ Financial analysis of catfish (*Clarias gariepinus*) hatchery business in Lembah Sari, Rumbai Pesisir district, Pekanbaru, the Province of Riau]. *Jurnal Perikanan dan Kelautan* 17(1):38-55. [in Indonesian]

Received: 29 February 2020. Accepted: 26 March 2020. Published online: 05 April 2020.

Authors:

Bambang Poerwanto, Magister Program of Fisheries Science, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: poerwantobambang2@gmail.com  
 Idiannor Mahyudin, Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: dr\_idiannor@yahoo.co.id  
 Leila Ariyani Sofia, Study Program of Fisheries Agribusiness, Fisheries and Marine Faculty, Lambung Mangkurat University, Jl. A. Yani Km. 36 Banjarbaru, South Kalimantan, Indonesia, e-mail: leila.ariyani@ulm.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Poerwanto B., Mahyudin I., Sofia L. A., 2020 The analysis of factors influencing carp (*Cyprinus carpio*) seed production in Bungur District, Tapin Regency, South Kalimantan. AACL Bioflux 13(2):804-812.

Search all conversations



Active



UNIVERSITAS  
JAMBUJANG



367 of 415 < >

Here you have the final form of the paper, which has also ben uploaded on the journal's website: <http://www.bioflux.com.ro/home/volume-13-2-2020/>.  
Thank you very much for paying so much attention to all the details!

Cordially yours,  
Claudiu G.

On Wednesday, April 8, 2020, 1:55:58 AM GMT+3, wrote:

Dear Dr. Gavrioloie,  
I have rechecked the whole document of Poerwanto et al's final form and made some minor corrections (yellow). Now, it is ready for uploading it. Thank you  
Sincerely,

On Wednesday, April 8, 2020, 01:39:07 AM GMT+8, gavrioloie ionel claudiu <[ioneiclaudiu@yahoo.com](mailto:ioneiclaudiu@yahoo.com)> wrote:

Dear,

Here you have the final form of the paper. I hope everything is good now. Just let me know when to upload it on the journal's website.  
Thank you!

Yours,  
Claudiu G.

Search all conversations



Active



363 of 415

Fwd: Fw: the almost final form of the paper of Poerwanto et al - The analysis of factors influencing carp (Cyprinus carpio) seed production ... Inbox



----- Forwarded message -----

Date: Kam, 9 Apr 2020 21:36

Subject: Fw: the almost final form of the paper of Poerwanto et al - The analysis of factors influencing carp (Cyprinus carpio) seed production ...

----- Forwarded Message -----

**From:** gavriloaie ionel claudiu <[ionelclaudiu@yahoo.com](mailto:ionelclaudiu@yahoo.com)>

**Sent:** Thursday, April 9, 2020, 09:59:02 PM GMT+8

**Subject:** Re: the almost final form of the paper of Poerwanto et al - The analysis of factors influencing carp (Cyprinus carpio) seed production ...

Dear,

I was able to do the correction. If you would have asked me to do it after more than 3-4 days it would not have been possible. You should see the new version online in up to 24 hours.

Thank you very much!

Cordially yours,

Claudiu G.

On Thursday, April 9, 2020, 8:51:01 AM GMT+3, wrote:

