

Jurnal Web of Science (Thyroid Dysfunction and Infertility of Women of Reproductive Age)

by Turnitin.com ®

Submission date: 09-Apr-2022 09:21PM (UTC+0700)

Submission ID: 1806069426

File name: id_Dysfunction_and_Infertility_of_Women_of_Reproductive_Age.pdf (198.92K)

Word count: 5104

Character count: 28633



25

Journal of Pharmaceutical Research International**32(23): 124-132, 2020; Article no. JPRI.61511****ISSN: 2456-9119***(Past name: British Journal of Pharmaceutical Research, Past ISSN: 2231-2919, NLM ID: 101631759)*

Thyroid Dysfunction and Infertility of Women of Reproductive Age

Ekaterina Andreevna Sterlyova^{1*}, Tatiana Kharitonovna Timokhina²,
Yuriy Viktorovich Sivkov³, Alexandr Anatolyevich Markov^{2,3}, Huldani⁴,
Vitaly V. Goncharov⁵ and Harun Achmad⁶

¹Stavropol State Medical University, Russia.

²Tyumen State Medical University, Tyumen, Russian Federation, Russia.

³Tyumen Industrial University, Tyumen, Russian Federation, Russia.

⁴Department of Physiology, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, South Kalimantan, Indonesia.

⁵Department of State and International Law, Kuban State Agrarian University Named after I. T. Trubilin, Krasnodar, Russia.

⁶Department of Pediatric Dentistry, Faculty of Dentistry, Hasanuddin University, Indonesia.

Authors' contributions

This work was carried out in collaboration among all authors. Author EAS designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors TKT and YVS managed the analyses of the study. Authors AAM, Huldani, VVG and HA managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2020/v32i2330799

Editor(s):

(1) Dr. Wenbin Zeng, Xiangya School of Pharmaceutical Sciences, Central South University, China.

Reviewers:

(1) C. Muhas, KTN College of Pharmacy, India.

(2) Happy Chuta, North Eastern Indira Gandhi Regional Institute of Health and Medical Sciences, India.

(3) A Julie Christy, Shri Sathya Sai Medical College and Research Institute, Sri Balaji Vidyapeeth University, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/61511>

Original Research Article

Received 18 July 2020

Accepted 23 September 2020

Published 01 October 2020

ABSTRACT

Autoimmunity of the thyroid gland (TAI) or its dysfunction is quite common among women of reproductive age, and there are suggestions in the literature that they are associated with an unfavorable level of fertility and a negative outcome of pregnancy, as in the case of spontaneous conception or after assisted reproductive technologies (ART).

This assumption makes it necessary to screen autoantibodies to thyrotropin (TSH) and thyroid peroxidase among infertile women who have made a number of attempts to become pregnant.

*Corresponding author: E-mail: ya.nk999@yandex.ru;

44

Some authors have conducted a number of studies where they have examined the relationship between autoimmunity of the thyroid gland, thyroid function, and fertility. However, there is currently no consensus on the upper limit of the norm for TSH to determine thyroid dysfunction and the limits for intervention. Despite the recent update of the American thyroid Association (ATA) on guidelines for the diagnosis and treatment of thyroid diseases during pregnancy and the postpartum period, many issues remain unresolved in ART. The author came to the following conclusions: open thyroid dysfunction often leads to menstrual disorders, fertility problems, and pregnancy complications, and therefore should be treated accordingly. Currently, there is little evidence to recommend treatment with levothyroxine at TSH levels between 2.5 and 4.0 MMU / l, given the possible side effects of overtreatment, especially in patients with mild thyroid dysfunction. We suggest careful longitudinal monitoring, especially in the presence of thyroid antibodies in women undergoing ART. The 4 MMU / l limit for TSH appears as the intervention level for SCH treatment in women with and without transabdominal ultrasound in ART.

Keywords: Thyroid; autoimmunity; infertility; assisted reproductive technologies.

1. INTRODUCTION

Birth is a fundamental evolutionary process for sustaining life. In this regard, it is very important to take into account the complex regulatory endocrine and immune factors that determine fertility and implantation.

The international committee for monitoring assisted reproductive technologies (ICMART) and the world health organization (WHO) define infertility (clinical definition) in the revised Glossary of ART terminology "as a disease of the reproductive system defined by the inability to achieve clinical pregnancy after 12 months or more of regular unprotected sexual intercourse [1]."

It has been determined that approximately one in six of all couples will have difficulty conceiving (primary infertility) or conceiving the desired number of children (secondary infertility) [2]. The world health organization (WHO) task force on infertility diagnosis and treatment conducted a study of 8,500 infertile couples in developed countries [3]. Female-related infertility accounted for 37%, and combined male and female factors accounted for 35% of the causes of infertility. The most common causes of female infertility were: ovulation disorders (25%), endometriosis (15%), pelvic adhesions (12%), blocked fallopian tubes (11%), other tube abnormalities (11%), and hyperprolactinemia (7%).

When faced with fertility problems, couples often seek help from fertility specialists. Over the past two decades, the field of fertility has been studied quite intensively, and researchers have tried to

identify curable risk factors that contribute to infertility. These risk factors include the presence of thyroid autoimmunity and thyroid dysfunction. Consequently, screening for TSH and anti-thyroperoxidase antibodies is usually a part of the initial study [4].

2. METHODS OF RESEARCH

This article is a review and is based on the analysis of data from the literature sources of specialists in the field of the problem under study. In the process of writing, we studied the publications of specialists in the field of thyroid dysfunction and infertility over the past 15 years. The type of research is analytical. In this work, we have studied the literature related to the problem of the effect of thyroid dysfunction on female infertility. Comparative and analytical research methods were applied. In addition, a statistical study of indicators related to the research topic and presented in various sources was carried out, as well as their description and systematization.

3. RESULTS AND DISCUSSION

Literature analysis showed a high prevalence of thyroid disorders (dysfunction and autoimmunity) among women of reproductive age [5]. The incidence of hypothyroidism ranges from 2% to 4% and is largely due to autoimmune thyroid disease (TAI) [6].

Subclinical thyroid abnormalities, including subclinical hypothyroidism, hypothyroxinemia, and / or isolated TAI, are much more common than clinical thyroid dysfunction.

The importance of thyroid hormones has been emphasized since the discovery of TSH, its receptors, and thyroid hormone receptors (TR- α 1 and TR- β 1) on the epithelium of the ovarian surface and in the oocytes of the primary, and secondary follicles. Thyroid hormones appear to be involved in the complex regulation of ovarian function. Thyroid hormones can also indirectly affect aspects of fertility by altering gonadotropin-releasing hormone (GnRH) and prolactin secretion, levels of sex hormone-binding globulin (SHBG), and coagulation factors.

Taking into account the importance of thyroid hormones, even mild thyroid insufficiency has been identified as one of the possible causes of adverse fertility and pregnancy outcomes.

The extent to which hyperthyroidism is associated with fertility problems is not clearly established due to limited data in the literature. However, in general, in the absence of adequate treatment, hyperthyroidism is associated with early pregnancy loss.

Thyrotoxicosis increases serum levels of sex hormone-binding globulin (SHBG) and estradiol (E2) compared to euthyroid women. The latter may be the result of an increase in SHBG or an increase in the production of E2 and androgen in combination with an increase in the degree of conversion to estrone and E2. In addition, LH secretion is increased among patients with Graves' disease compared to patients with euthyroidism.

Some authors have noted that primary or secondary infertility occurs among 5.8% of patients with hyperthyroidism [7]. Menstrual disorders are common among women with hyperthyroidism. Early data indicate menstrual irregularities of up to 65%, compared with 17% of healthy people in the control group. More recent data showed a significantly lower prevalence of menstrual disorders - about 22% compared to 8% of healthy people in the control group. This contradiction is related to different approaches to the diagnosis and treatment of clinical hyperthyroidism. Hypomenorrhea, polymenorrhea, oligomenorrhea, and hypermenorrhea are the most common disorders of the menstrual cycle.

Open hypothyroidism is associated with an increased risk of fertility problems and adverse complications of early and late pregnancy [8]. Hypothyroidism leads to a number of hormonal

changes. The rate of metabolic clearance of both Androstenedione and estrone decreases, and peripheral aromatization increases. In addition, SHBG plasma binding activity is reduced. Consequently, plasma concentrations of both total testosterone and E2 decrease, and their unbound fraction increases.

Hypothyroidism can also lead to a blunted LH response, which stimulates TRH secretion and increases serum prolactin levels. Since prolactin impairs the pulsating secretion of gonadotropin-releasing hormone (GnRH), this can lead to ovulation dysfunction, including yellow body insufficiency with low levels of progesterone secretion in the luteal phase of the cycle [9]. Therefore, clinical hypothyroidism leads to a number of ovulation disorders among women of fertile age. There may be changes in the length of the cycle, as well as in the volume of menstrual bleeding secondary to breakthrough bleeding after anovulation and / or violation of hemostatic factors associated with hypothyroidism.

Most authors do not always link subclinical hypothyroidism (SCH) and infertility. This may be partly due to different thresholds used to determine the upper limit of normal TSH concentration, as well as the lack of well-planned prospective studies.

In a large retrospective cross-sectional study of 11,254 women in Denmark, impaired fertility was associated with subclinical hypothyroidism, defined as TSH > 3.7 mU / l [10]. A recent cross-sectional study also showed higher normal TSH levels in women with unexplained infertility compared to the fertility control [9].

Based on these data, it can be said that the levels of TSH used in various studies to determine the relationship of thyroid function with fertility problems varied significantly. In general, the Association with adverse fertility outcomes appears to occur at TSH levels above 4.0 mU / l.

TAI is the most common autoimmune disease of women of childbearing age and increases the risk of thyroid dysfunction. The prevalence of TAI is usually estimated at about 10% and is more common in women seeking infertility consultation.

Other causes of infertility have also been linked to TAI. Women with polycystic ovary syndrome

(PCOS) were found to have an increased prevalence of TAI. A possible explanation may be polymorphisms of the PCOS-related fibrillin 3 gene, which affects the activity of TGF- β , a key regulator of immune tolerance.

Together with lower TGF- β , low vitamin D levels, and a high estrogen-to-progesterone ratio, these factors may contribute to autoimmunity [11].

As for endometriosis, the data are more contradictory, with conflicting results of increased prevalence of TAI. There is evidence that endometriosis is associated with various immunological changes, including antibodies to endometrial antigens.

In general, high levels of TPO-abs have been identified as the most sensitive marker of TAI and are associated with the risk of subclinical hypothyroidism. Therefore, most studies examining the prevalence of TAI among fertile women or any association between TAI and pregnancy outcomes are based on the presence of only TPO-abs and do not take into account autoantibodies to thyroglobulin (Tg-abs).

Also in the literature, the Association of TAI with adverse pregnancy outcomes, with an increased risk of miscarriage and premature birth in spontaneous pregnancy, as well as in pregnancy after ART. Since the late 1990s, several studies have been published on the effects of TAI on post-ART outcomes. The results of these studies, however, are contradictory, probably due to different study plans, a small number of registered patients with a low absolute number of analyzed events, and the use of surrogate endpoints for the fertility result (Table 1).

Research by Zhong et al. comparison of IVF results in TAI-positive and TAI-negative women showed that TAI-positive women had significantly lower fertilization (64.3% vs. 74.6%), implantation (17.8% vs. 27.1%), and pregnancy rate (33.3% vs. 46.7. %) and a higher risk of pregnancy termination (26.9 vs. 11.8%) after IVF compared to their negative TAI counterparts. Two meta-analyses showed an increase in the frequency of miscarriages of TAI-positive women with subfertility. In the first meta-analysis of four prospective studies conducted by Toulis, women with TAI were twice as likely to experience a miscarriage after IVF pregnancy.

10

The actual difference in the number of miscarriages between euthyroid women with TAI who underwent IVF and the control group (absolute risk) remained small, which led to a negligible impact on the frequency of clinical pregnancies and deliveries.

In the second meta-analysis of 12 studies, the risk of miscarriage was significantly higher among women with TAI, resulting in a lower birth rate. However, the amount of TAI did not affect the number of eggs extracted and the frequency of fertilization.

More recent studies on this topic have not been able to confirm the link between TAI and their negative impact on fertility. In a retrospective study, Lukaszuk et al. compared pregnancy outcome between 114 TAI-positive and 495 TAI-negative infertile women. No significant differences were found in fertilization, implantation, pregnancy rates, live birth, or higher risk of miscarriage. A prospective study by Sakar et al. showed comparable rates of pregnancy and miscarriage between 49 positive and 202 negative women after IVF.

According to Tan et al. any previous association between TAI-positive women and a negative pregnancy outcome may be confused due to selection bias, particularly the choice of female patients who seek treatment for fertility problems. To avoid this bias, the author selected couples with male infertility and studied female partners with TAI, but without known female infertility. They found that the pregnancy outcome was comparable among women with and without thyroid autoimmunity after intracytoplasmic sperm injection [12].

The main pathophysiological mechanisms that link the possible negative impact of TAI on the outcome of pregnancy after ART remain unknown. The presence of TAI may represent a peripheral marker of a general immune imbalance, which can lead to insufficient fertilization, insufficient implantation, and prolonged pregnancy. The mechanism by which TAI can act is a change in endometrial susceptibility that affects the fetal allograft. Quantitative and qualitative changes in the profile of endometrial T cells with reduced IL-4 and IL-10 secretion along with increased interferon- γ secretion have been reported.

Hyperactivity and increased migration of cytotoxic natural killer cells can also alter the immune and hormonal response of the uterus in women with TAI. Compared to women who have had one or no miscarriages, women with multiple miscarriages appear to have an increased number of CD5 / 20 + B cells with a higher prevalence in women with thyroid antibodies [13]. Activation of polyclonal B cells is more often observed in TAI and is associated with an increase in the titers of inorganic specific autoantibodies. Limited data indicate the presence of antiphospholipid (APL) and / or anti-nuclear (ANA) antibodies in the population of infertile women with TAI. Anti-lupus antibodies and anti-cardiolipin antibodies are significantly more common in women with infertility. In general, the influence of various associated autoimmune diseases on reproductive health indicators is largely ignored, and it requires further study.

Alternatively, anti-thyroid antibodies may be the direct cause, since the presence of anti-thyroid antibodies can have an adverse effect on the quality of oocytes and embryos. Indeed, the authors also found significantly lower oocyte fertilization and the percentage of class A embryos when comparing infertile women who underwent IVF with thyroid autoimmunity with negative controls. For the first time, this study documented the presence of thyroid antibodies in ovarian follicular fluid. The authors suggested that thyroid antibodies may pass the "follicle-blood" barrier during secondary follicle maturation. The cytotoxicity of these antibodies can then damage the maturing oocyte and ultimately reduce oocyte quality and fertilization potential.

On the other hand, a positive TAI status increases the risk of developing (sub) clinical hypothyroidism. Women with TAI have an increased risk of developing (sub) clinical hypothyroidism during spontaneous pregnancy [14]. In the case of assisted reproductive technology (ART) and ovarian hyperstimulation, it has been reported that TSH levels rise significantly to levels above 2.5 Mme / l before pregnancy, and even more so in the presence of TAI [15]. Since thyroid hormones play an important role in oocyte maturation and implantation, it has been suggested that a decrease in thyroid function caused by the stimulation Protocol in women with TAI may negatively affect the pregnancy rate during ART.

In the literature there are also studies on approaches to treating hypothyroidism in order to increase fertility in women. Detailed information about the effect of hypothyroidism on the results of controlled ovarian hyperstimulation (HOG) and IVF is limited, as there are no available randomized controlled trials. The reason lies in the unethical question of refusing treatment to patients with clinical obvious hypothyroidism. Numerous negative effects of maternal overt hypothyroidism on pregnancy outcomes and adverse neurocognitive effects of offspring have been documented in the literature. Treatment of hypothyroidism with levothyroxine usually restores normal menstrual structure, eliminates hormonal changes, and improves fertility [16]. However, some women with treated hypothyroidism still cannot get pregnant and seek infertility treatment, including controlled ovarian hyperstimulation and / or IVF. Treatment of apparent hypothyroidism in pregnant women is mandatory and includes levothyroxine therapy adapted to achieve normal trimester serum levels that stimulate thyroid-stimulating hormone (TSH).

The latest ATA recommendations call for achieving TSH concentrations below 2.5 Mme / l in patients with clinical hypothyroidism. An absolute inability to produce thyroid hormones during COX can actually have a detrimental effect on the oocyte and / or endometrium. However, the level of TSH that should be reached before starting ART remains controversial.

A meta-analysis of 3 RCTS showed a positive effect of levothyroxine treatment on pregnancy after ART in women with subclinical hypothyroidism. Although no benefit was shown for clinical pregnancy rates (combined relative risk 1.75, 95% CI 0.90–3.38), there was an increase in delivery rates (combined relative risk 2.76, 95% CI 1.20–6.44). However, the included studies differed with respect to the upper TSH limit for determining SCH. This limit often exceeded 4 Mme / l. In fact, most of the available data conclude that there is no difference in the outcome of ART in women with euthyroid disease with TSH levels <2.5 Mme / l and in women with a slight increase in TSH levels from 2.5 to 5.0 Mme / l.

The use of appropriate control ranges should be based on local control values for the population and laboratory. Moreover, in pregnancy, in accordance with the ATA recommendations, it is recommended to use TSH concentrations based

on reference ranges typical for pregnancy [17]. If this is not possible, it is recommended to use a fixed upper limit of 4.0 MMU / l, which usually corresponds to a decrease of 0.5 MMU / l compared to the negligent TSH reference range. These recommendations for the diagnosis and treatment of SCH seem reasonable in the case of ART, since adverse events were mainly reported in patients with TSH values above 4.0 Mme / l.

We have studied four randomized controlled trials that evaluated the effect of levothyroxine treatment in SCH on various aspects of spontaneous pregnancy outcomes. Thus, some authors have noted that treatment with levothyroxine improved adverse pregnancy outcomes in women with a positive response to TPO, the level of TSH exceeded 2.5 Mme / l.

In another study, there was also a decrease in preterm birth in women with a positive response to TPO when treated with levothyroxine, but only in patients with TSH levels exceeding 4.0 Mme / l [16]. The same authors also confirmed only the positive effect of LT4 therapy in reducing preterm birth in women with a negative reaction to TPO and TSH levels above 4 Mme / l.

In a meta-analysis presented in one of the studies, a subgroup analysis involving two studies the effect of non-intervention treatment for SCH on the risk of miscarriage was compared. The results showed that compared with patients who received effective medication, the risk of miscarriage increased significantly in patients without intervention with a total total HR of 1.50 (95% CI 1.03–2.19, P = 0.04). However, the latest study needs to be interpreted with caution due to possible selection bias, different diagnostic criteria for SCH, and different definitions used for pregnancy outcome.

Taking into account the inconsistencies and limitations of available interventional studies, treatment with levothyroxine seems to be appropriate in women with a positive response to TPO when TSH levels exceed the reference range unrelated to pregnancy. However, there is insufficient data in the literature to initiate treatment with levothyroxine in women with a positive response to TPO with TSH levels from 2.5 to 4 Mme / l. In addition, based on a recent study on women with a negative reaction to TPO, treatment can only be recommended for TSH levels exceeding 4 Mme / l. Current recommendations suggest treating SCH with levothyroxine during pregnancy in order to

maintain TSH <2.5 Mme / l. On the other hand, studies show that an increase in TSH in pregnant women should be the upper limit of the reference pregnancy interval, rather than 2.5 Mme / l, and this can also be applied during pregnancy obtained after ART.

Currently, the benefit of levothyroxine treatment for pregnancy outcomes in women with euthyroid disease TAI both during spontaneous pregnancy and after ART is questionable.

In the case of ART, treatment with levothyroxine does not appear to be beneficial for TAI-positive euthyroid women. In specialized studies, there are no improvements in TAI-positive women who have undergone IVF. These results were confirmed by RCTS in China, where treatment with levothyroxine compared to treatment with no levothyroxine did not reduce the frequency of miscarriages and did not increase the number of live births among women with normal thyroid function who tested positive for thyroid autoantibodies undergoing in vitro fertilization and embryo transfer.

The results of the TABLET study were also examined as part of the study. This multicenter, double-blind, randomized controlled trial examined the effects of levothyroxine at a dose of 50 mcg per day in women who tested positive for TPO-abs before conception. Treatment with levothyroxine did not lead to a higher rate of live births or pregnancy compared to placebo, which confirms the findings of previous RCTS.

Despite the low level of evidence, it is recommended to start treatment with any increase in TSH > 2.5 Mme / l before ART, given its potential advantages over minimal risk. However, some potential dangers have recently been noted, indicating that excessive treatment of the thyroid gland in the mother in early pregnancy is associated with harmful effects on IQ and brain morphology in children in childhood. Recently, there have been additional concerns that levothyroxine treatment may increase the risk of preterm birth, gestational diabetes, and preeclampsia.

Most authors agree that all patients seeking medical advice about infertility should be screened for the presence of an underlying thyroid disease, taking into account the potential harmful effect of thyroid dysfunction on the outcome of fertility. A thorough medical history and medical examination should be the first step

Table 1. Main characteristics of research on the relations between TAI and IVF / ICSI results

Author	Year	Design of the study	Purpose of the study	Tests of the thyroid gland	TAI Number +	TAI Number -	State of the thyroid function	Main conclusion
Muller et al.	1999	Retrospective cohort study	LBR; CPR; F-H	TPOAbs; Tg Abs	143	730	euthyroid	Does not affect the outcome of pregnancy
Poppe et al.	1999	Prospective cohort study	CPR; F-H; OPR	TPOAbs	25	148	euthyroid	Does not affect the outcome of pregnancy
Negro et al.	2003	Prospective cohort study	LBR; CPR; F-H	TPOAbs	32	202	euthyroid	Lower LBR; increased MR
Negro et al.	2005	Prospective cohort study	LBR; CPR; F-H; NOR	TPOAbs	43	576	euthyroid	Lower LBR; does not affect CPR
Kilic et al.	2007	Retrospective cohort study	LBR; CPR; F-H; NOR	TPOAbs	42	374	euthyroid	Does not affect the outcome of pregnancy
Zhong et al.	2008	Prospective cohort study	CPR; F-H; NOR	TPOAbs; Tg Abs	23	31	euthyroid	below CPR
Karacan et al.	2012	Retrospective cohort study	CPR; F-H; IK; NOR; FR	TPOAbs; Tg Abs	90	676	He ykazano	below CPR, FR, IR and above MR
Mintzioti et al.	2013	Prospective cohort study	NOR; FR; IK; CPR; F-H; OPR	TPOAbs; Tg Abs	34	219	euthyroid	Does not affect the outcome of pregnancy
Tan et al.	2014	Retrospective cohort study	LBR; NOR; CPR; F-H	TPOAbs; Tg Abs	15	67	euthyroid	Does not affect the outcome of pregnancy
Chai et al.	2014	Retrospective cohort study	LBR; CPR; F-H	TPOAbs; Tg Abs	110	725	euthyroid	Does not affect the outcome of pregnancy
Lukasuk et al.	2014	Retrospective cohort study	LBR; CPR; F-H	TPOAbs; Tg Abs	89	419	euthyroid	Does not affect the outcome of pregnancy
Litwicka et al.	2015	Retrospective cohort study	LBR; CPR; F-H; NOR; infrared	TPOAbs	114	551	euthyroid	Does not affect the outcome of pregnancy
Sakar et al.	2015	Prospective cohort study	LBR; CPR; MR, NOR	TPOAbs; Tg Abs	60	134	euthyroid	Lower LBR; increased MR
Unuane et al.	2016	Prospective cohort study	CPR; OPR; F-H; NOR	He ykazano	49	202	unspecified	Does not affect the outcome of pregnancy
Muller et al.	2016	Retrospective cohort study	LBR; CPR; F-H	TPOAbs	333	2019	euthyroid	Does not affect the outcome of pregnancy

*LBR = lifetime fertility rate; CPR = clinical pregnancy rate; OPR = current pregnancy; MR = miscarriage rate; IR = implantation rate; NOR = number of oocytes extracted; FR = fertilization rate

before starting treatment. In addition to more obvious symptoms such as weight changes, fatigue, irritability, and heart palpitations, one should also pay attention to symptoms such as menstrual irregularities that may indicate thyroid dysfunction. The laboratory assessment should include at least TSH and TPO-abs.

In case of obvious thyroid dysfunction, appropriate therapy should be started immediately. SCH is best treated when TSH values exceed the normal upper value determined by population non-pregnant values, or in the absence of these control ranges with a fixed threshold of 4 Mm / l.

Taking into account the uncertainty in the literature regarding the harm-benefit ratio in patients with TSH cut-off levels of 2.5 to 4.0 Mm / l, intervention in these cases should be limited and discussed with patients regarding the best available data. Currently, there is not enough time to start treatment with levothyroxine at pre-conception TSH levels of 2.5 to 4.0 Mm / l, especially in the case of ART.

4. CONCLUSION

The interaction between thyroid disease and fertility is complex. Open thyroid dysfunction often leads to menstrual disorders, fertility problems, and pregnancy complications, and therefore should be treated accordingly. There is currently little evidence to recommend treatment with levothyroxine with TSH levels between 2.5 and 4.0 Mm / l, given the possible side effects of overtreatment, especially in patients with mild thyroid dysfunction. We offer thorough longitudinal monitoring, especially in the presence of anti-thyroid antibodies among women undergoing ART. The 4 Mm / l limit for TSH appears as an intervention level for SCH treatment among women with and without TAI in ART.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Evers JLH. Female subfertility. *Lancet*. 2002;360:151-159.
2. Poppe K, Glinooer D, Van Steirteghem A, et al. Thyroid dysfunction and autoimmunity in infertile women *Thyroid*. 2002;12:997-1001.
3. Iravani AT, Saeedi MM, Pakravesh J, et al. Thyroid autoimmunity and recurrent spontaneous abortion in Iran: A case-control study *Endocr Pract*. 2008;14:458-464.
4. Alexander EK, Elizabeth N, Pearce EN, et al. Guidelines of the American thyroid association for the diagnosis and management of thyroid disease during pregnancy and the postpartum *Thyroid*. 2017;27:315-389.
5. Aghajanova L, Lindeberg M, Carlsson IB, et al. Receptors for thyroid-stimulating hormone and thyroid hormones in human ovarian tissue *Reprod Biomed Online*. 2009;18:337-347.
6. Millar LK, Wing DA, Leung AS, et al. Low birth weight and preeclampsia, in pregnancies complicated by hyperthyroidism *Obstet Gynecol*. 1994;84: 946-949.
7. McElduff A, Morris J. Thyroid function tests and thyroid autoantibodies in an unselected population of women undergoing first trimester screening for aneuploidy *Aust N Z J Obstet Gynaecol*. 2008;48:478-480.
8. Krassas GE, Poppe K, Glinooer D. Thyroid function and human reproductive health *Endocr Rev*. 2010;31: 702-755.
9. Poppe K, Glinooer D, Tournaye H, et al. Impact of ovarian hyperstimulation on thyroid function in women with – and without thyroid autoimmunity *J Clin Endocrinol Metab*. 2004;89:3808-3812.
10. Achmad H, Hendra Chandha, Sri Harun, Sudjarwo I, Yunus M, Rahma K. Rusdi, Khairunnisa P. Prevalence of Medically Compromised Children Regarding Dental Caries and Treatment Needs in Wahidin Sudirohusodo Hospital, *Journal of International Dental and Medical Research*. 2017;10(3):915-920. ISSN 1309-100X.
11. Markov Alexander. Results of the Experimental Research and Clinical Application of Cannulated Screw with

- Bioactive Coverage on the Basis of Natural Hydroxiapatite during Osteosynthesis of Medial Fractures of Femur's Neck. Journal of Biomimetics, Biomaterials and Biomedical Engineering Trans Tech Publications Ltd, Switzerland. 2020;46:52-59. ISSN: 2296-9845.
12. Unuane D, Velkeniers B, Anckaert E, et al. Thyroglobulin autoantibodies: is there any added value in the detection of thyroid autoimmunity in women consulting for fertility treatment? *Thyroid*. 2013;23:1022-1028.
 13. Roberts J, Jenkins C, Wilson R, et al. Recurrent miscarriage is associated with increased numbers of CD5/20 positive lymphocytes and an increased incidence of thyroid antibodies *Eur J Endocrinol*. 1996;134:84-86.
 14. Poppe K, Glinoeer D, Tournaye H, et al. Impact of the ovarian hyperstimulation syndrome on thyroid function *Thyroid*. 2008;18:801-802.
 15. Subclinical hypothyroidism in the infertile female population: a guideline. Practice Committee of the American Society for Reproductive Medicine *Fertil Steril*. 2015; 104:545-555.
 16. Kutteh WH, Schoolcraft WB, Scott Jr RT. Antithyroid antibodies do not affect pregnancy outcome in women undergoing assisted reproduction *Hum Reprod*. 1999; 14:2886-2890.
 17. Tuncay G, Karaer A, Coskun EI, et al. The impact of thyroid-stimulating hormone levels in euthyroid women on intrauterine insemination outcome *BMC Wom Health*. 2018;18:51. DOI:10.1186/s12905-018-0541-0

© 2020 Sterlyova et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/61511>

Jurnal Web of Science (Thyroid Dysfunction and Infertility of Women of Reproductive Age)

ORIGINALITY REPORT

19%

SIMILARITY INDEX

10%

INTERNET SOURCES

16%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | Bert Scoccia. " <i>In Vitro</i> Fertilization Pregnancy Rates in Levothyroxine-Treated Women With Hypothyroidism Compared to Women Without Thyroid Dysfunction Disorders", <i>Thyroid</i> , 04/27/2012
Publication | 1% |
| 2 | journalair.com
Internet Source | 1% |
| 3 | pubmed.ncbi.nlm.nih.gov
Internet Source | 1% |
| 4 | Submitted to Nassau Community College
Student Paper | 1% |
| 5 | Alban Deroux, Chantal Dumestre-Perard, Camille Dunand-Faure, Laurence Bouillet, Pascale Hoffmann. "Female Infertility and Serum Auto-antibodies: a Systematic Review", <i>Clinical Reviews in Allergy & Immunology</i> , 2016
Publication | 1% |
-

6	journals.eco-vector.com Internet Source	1 %
7	cyberleninka.org Internet Source	1 %
8	Andrea Weghofer, Eric Himaya, Vitaly A. Kushnir, David H. Barad, Norbert Gleicher. "The impact of thyroid function and thyroid autoimmunity on embryo quality in women with low functional ovarian reserve: a case-control study", <i>Reproductive Biology and Endocrinology</i> , 2015 Publication	<1 %
9	Gesthimani Mintziori, Dimitrios G. Goulis, Konstantinos A. Toulis, Christos A. Venetis et al. "Thyroid function during ovarian stimulation: a systematic review", <i>Fertility and Sterility</i> , 2011 Publication	<1 %
10	www.eje-online.org Internet Source	<1 %
11	Na Young Kim. "Thyroid Autoimmunity and its Association with Cellular and Humoral Immunity in Women with Reproductive Failures : THYROID AUTOIMMUNITY AND REPRODUCTIVE FAILURES", <i>American Journal Of Reproductive Immunology</i> , 01/2011 Publication	<1 %

12 Nkiruka Rose Ukibe, Solomon Nwabueze Ukibe, Obiageli Fidelia Emelumadu, Chinedum Charles Onyenekwe et al. "Impact of thyroid function abnormalities on reproductive hormones during menstrual cycle in premenopausal HIV infected females at NAUTH, Nnewi, Nigeria", PLOS ONE, 2017
Publication

13 www.thieme-connect.de
Internet Source

14 Kris Poppe, Candice Autin, Flora Veltri, Georgiana Sitoris, Pierre Kleynen, Jean-Philippe Praet, Serge Rozenberg. " Thyroid Disorders and Outcomes of Assisted Reproductive Technology: An Unfortunate Combination? ", Thyroid, 2020
Publication

15 doaj.org
Internet Source

16 Submitted to American Public University System
Student Paper

17 Weiping Teng. "Pregnancy and Thyroid Diseases in China", Thyroid, 10/2011
Publication

18 Mestman, Jorge H.. "Should Infertile Women Receive Thyroid-Replacement Therapy If Their

Serum TSH \square Is >2.5 mIU/L in the Prepregnancy State?", *Clinical Thyroidology*, 2015.

Publication

19

Paolo Giovanni Artini, Patrizia Monteleone. "Chapter 6 Female Infertility and Autoimmunity", Springer Science and Business Media LLC, 2018

Publication

20

Anne-Dorthe Feldthusen, Palle L. Pedersen, Jacob Larsen, Tina Toft Kristensen, Christina Ellervik, Jan Kvetny. "Impaired Fertility Associated with Subclinical Hypothyroidism and Thyroid Autoimmunity: The Danish General Suburban Population Study", *Journal of Pregnancy*, 2015

Publication

21

onlinelibrary.wiley.com

Internet Source

22

Michael H. Berger, Marisa Messori, Alexander W. Pastuszak, Ranjith Ramasamy. "Association Between Infertility and Sexual Dysfunction in Men and Women", *Sexual Medicine Reviews*, 2016

Publication

23

Felipe A. Morales-Martínez, Luis H. Sordia-Hernández, Martha Merino Ruiz, Selene Garcia-Luna et al. "Association between

<1 %

<1 %

<1 %

<1 %

<1 %

thyroid autoimmunity and ovarian reserve in women with hypothyroidism", Thyroid Research, 2021

Publication

24

K. Łukaszuk, M. Kunicki, P. Kulwikowska, J. Liss, E. Pastuszek, M. Jaszczółt, B. Męczekalski, K. Skowroński. "The impact of the presence of antithyroid antibodies on pregnancy outcome following intracytoplasmatic sperm injection-ICSI and embryo transfer in women with normal thyreotropine levels", Journal of Endocrinological Investigation, 2015

Publication

<1 %

25

Submitted to Nha Trang University

Student Paper

<1 %

26

Reshma M. Ansari. "Potential use of durian fruit (*Durio zibenthinus* Linn) as an adjunct to treat infertility in polycystic ovarian syndrome", Journal of Integrative Medicine, 2016

Publication

<1 %

27

Submitted to University of Kent at Canterbury

Student Paper

<1 %

28

Submitted to Upper Iowa University

Student Paper

<1 %

29

Petrache Vartej, Ioana Vartej. "Chapter 5 Thyroid Diseases and Female Infertility", Springer Science and Business Media LLC, 2018

Publication

<1 %

30

Yingying Zhang, Ling Wu, Tin Chiu Li, Chi Chiu Wang, Tao Zhang, Jacqueline Pui Wah Chung. "Systematic review update and meta-analysis of randomized and non-randomized controlled trials of ovarian stimulation versus artificial cycle for endometrial preparation prior to frozen embryo transfer in women with polycystic ovary syndrome", Reproductive Biology and Endocrinology, 2022

Publication

<1 %

31

www.bmj.com

Internet Source

<1 %

32

www.nature.com

Internet Source

<1 %

33

www.sciendo.com

Internet Source

<1 %

34

Submitted to President University

Student Paper

<1 %

35

Sima Nazarpour, Fahimeh Ramezani Tehrani, Masoumeh Simbar, Maryam Tohidi, Sonia Minooe, Maryam Rahmati, Fereidoun Azizi.

<1 %

"Effects of Levothyroxine on pregnant women with subclinical hypothyroidism, negative for thyroid peroxidase antibodies", The Journal of Clinical Endocrinology & Metabolism, 2017

Publication

36

Sofie Bliddal, Malene Boas, Linda Hilsted, Lennart Friis-Hansen et al. "Increase in thyroglobulin antibody and thyroid peroxidase antibody levels, but not preterm birth-rate, in pregnant Danish women upon iodine fortification", European Journal of Endocrinology, 2017

Publication

<1 %

37

Submitted to Temple University

Student Paper

<1 %

38

epdf.pub

Internet Source

<1 %

39

www.gonadotropin.org

Internet Source

<1 %

40

www.researchsquare.com

Internet Source

<1 %

41

"42nd Annual Meeting of the European Thyroid Association. Budapest, Hungary. September 7–10, 2019", European Thyroid Journal, 2019

Publication

<1 %

42 "Acute Cardiovascular Care 2018", European Heart Journal: Acute Cardiovascular Care, 2018
Publication <1 %

43 "Poster Presentations", Thyroid, 09/2005
Publication <1 %

44 "Subsection Reports", The Journal of Clinical Endocrinology & Metabolism, 2007
Publication <1 %

45 Allan C. Dong, Jessica Morgan, Monica Kane, Alex Stagnaro-Green, Mary D. Stephenson. "Subclinical hypothyroidism and thyroid autoimmunity in recurrent pregnancy loss: a systematic review and meta-analysis", Fertility and Sterility, 2020
Publication <1 %

46 Maarten Buimer. "Transient Hypothyroxinemia in Severe Hypertensive Disorders of Pregnancy :", Obstetrics and Gynecology, 11/2005
Publication <1 %

47 assets.researchsquare.com
Internet Source <1 %

48 iopscience.iop.org
Internet Source <1 %

49 www.scribd.com
Internet Source <1 %

50 " 89 Annual Meeting of the American Thyroid Association ", Thyroid, 2019 <1 %
Publication

51 Batuhan Turgay, Yavuz Emre Şükür, Hasan Ulubaşoğlu, Murat Sönmezer et al. "The association of thyroid stimulating hormone levels and intrauterine insemination outcomes of euthyroid unexplained subfertile couples", European Journal of Obstetrics & Gynecology and Reproductive Biology, 2019 <1 %
Publication

52 Francesco Latrofa. "Autoimmune Hypothyroidism", Contemporary Endocrinology, 2008 <1 %
Publication

53 Submitted to University of Central Florida <1 %
Student Paper

54 ecerm.org <1 %
Internet Source

55 healthcare.walgreens.com <1 %
Internet Source

56 jdgstage.thyroidmanager.org <1 %
Internet Source

57 medcraveonline.com <1 %
Internet Source

58

Internet Source

<1 %

59

www.mdpi.com
Internet Source

<1 %

60

E. van den Boogaard. "Significance of (sub)clinical thyroid dysfunction and thyroid autoimmunity before conception and in early pregnancy: a systematic review", Human Reproduction Update, 09/01/2011
Publication

<1 %

61

Arfan Utiahman, Ayad F. Alkaim, Aseel M. Aljeboree, Smirnova Zhanna Venediktovna et al. "Role of elastostatic loading and cyclic cryogenic treatment on relaxation behavior of Ce-based amorphous alloy", Materials Today Communications, 2021
Publication

<1 %

62

Submitted to University of Aberdeen
Student Paper

<1 %

Exclude quotes On
Exclude bibliography On

Exclude matches Off

Jurnal Web of Science (Thyroid Dysfunction and Infertility of Women of Reproductive Age)

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
