

# A Review of Bandotan Leaf Extract (*Ageratum conyzoides* L.) in Inhibition Test to the Growth of Bacteria (*Porphyromonas* *gingivalis*) Case of Periodontitis Disease

*by* drg bayuindra

---

**Submission date:** 29-Jul-2020 10:35PM (UTC-0400)

**Submission ID:** 1358040839

**File name:** teria\_Porphyromonas\_gingivalis\_Case\_of\_Periodontitis\_Disease.doc (732K)

**Word count:** 3238

**Character count:** 22081

## A Review of Bandotan Leaf Extract (*Ageratum conyzoides* L.) in Inhibition Test to the Growth of Bacteria (*Porphyromonas gingivalis*) Case of Periodontitis Disease

<sup>1</sup>Harun Achmad, <sup>2</sup>Andi Mardiana Adam, <sup>3</sup>Aulia Azizah, <sup>4</sup>Bayu Indra Sukmana, <sup>5</sup>Huldani, <sup>6</sup>Shafira Nurul Khera, <sup>7</sup>Yunita Feby Ramadhany

<sup>1</sup>Department of Pediatric Dentistry, Dentistry Faculty of Hasanuddin University, Makassar, South Sulawesi, Indonesia

<sup>2</sup>Department of Periodontology, Dentistry Faculty of Hasanuddin University, Makassar, South Sulawesi, Indonesia

<sup>3</sup>Department of Preventive and Public Health Dentistry, Faculty of Dentistry, Lambung Mangkurat University, Banjarmasin, Indonesia.

<sup>4</sup>Department of Dental Radiology, Faculty of Dentistry, Lambung Mangkurat University, Banjarmasin Indonesia.

<sup>5</sup>Department of Physiology, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin Indonesia.

<sup>6</sup>Clinical Dental Student, Dentistry Faculty of Hasanuddin University, Makassar, South Sulawesi, Indonesia

<sup>7</sup>Dentist, Dentistry Faculty of Hasanuddin University, Makassar, South Sulawesi, Indonesia

E-mail: [harunachmader@gmail.com](mailto:harunachmader@gmail.com).

Article History:

Submitted: 10.01.2020

Revised: 12.03.2020

Accepted: 24.04.2020

### ABSTRACT

**Introduction:** Periodontitis is defined as an inflammatory disease in the supporting tissues of teeth which is caused by microorganisms of certain groups.

**Objective:** This study aims to know the inhibition power of bandotan leaf extract (*Ageratum conyzoides* L.) against the growth of bacteria causing periodontitis (*Porphyromonas gingivalis*).

**Method:** The type of study used in this research is experimental laboratory. The design of this study used is post test only control group design used the method of dilution and disk-diffusion method. **Results:** The activities of bacteria can only be observed with the method of diffusion to see the inhibition zone. An extract of bandotan leaf (*Ageratum conyzoides* L.) with the concentration of 100% has the largest inhibition zone namely 16,25±0,66 mm and the inhibition zone is larger than control positive namely metrodinazole used as a comparison antibiotic from the extract of bandotan leaf (*Ageratum*

*conyzoides* L.). This research has proven that the extract of bandotan leaf extract (*Ageratum conyzoides* L.) are effective as inhibitory properties against *Porphyromonas gingivalis*.

**Conclusion:** The extract of bandotan leaf (*Ageratum conyzoides* L.) is effective as the inhibition power against *Porphyromonas gingivalis*.

**Keywords:** *Ageratum conyzoides* L, *Porphyromonas Gingivalis*, Periodontitis

**Correspondence:**

Harun Achmad

Department of Pediatric Dentistry, Dentistry Faculty of Hasanuddin University

Makassar, South Sulawesi, Indonesia

E-mail: [harunachmader@gmail.com](mailto:harunachmader@gmail.com)

DOI: [10.31838/srp.2020.4.58](https://doi.org/10.31838/srp.2020.4.58)

@Advanced Scientific Research. All rights reserved

### INTRODUCTION

Dental health disease is the sixth highest health problem often complained by Indonesian people.<sup>1</sup>

Periodontal disease is one of the dental and oral diseases that develops very commonly, and affects about 10.5% to 12% of the world population.<sup>2</sup> In Indonesia periodontal disease has a fairly high prevalence suffered by humans almost all over the world and reaches 50% of the adult population.<sup>3</sup>

Periodontal disease is an inflammation that occurs in the supporting tissues of teeth, including alveolar bone and periodontal ligaments. Periodontal disease often found is gum inflammation or gingivitis and periodontitis.<sup>4</sup>

Periodontitis is defined as an inflammatory disease in the supporting tissues of the teeth caused by specific microorganisms of certain groups, which results in progressive damage of the periodontal ligament and alveolar bone by increasing the formation of probing depths, recessions, or both.<sup>2</sup>

A number of experimental evidence has shown that the primary etiological agent of periodontal disease is a generally Gram-negative stem bacterium which includes *P.gingivalis*. The main habitat of *P. gingivalis* is the subgingival sulcus of the human oral cavity. These periodontopathic bacteria are found in 85.75% of subgingival plaque samples from patients with chronic periodontitis. Oral bacteria and especially pathogenic bacteria, such as

*Porphyromonas gingivalis* have great virulence factors, one of which is the ability to penetrate hard intraoral surfaces.<sup>5</sup>

Efforts to overcome these problems, usually use the concept of antimicrobial treatment. Antibiotics are a group of antimicrobial agents, which also consist of antiviral, antifungal, and antiparasitic chemicals. Antibiotics can kill or suppress live bacteria but they cannot eliminate bacterial calculus and residues, which are traditionally considered an important part of periodontal therapy.<sup>6</sup>

When used appropriately, antibiotics provide unquestionable benefits. However, if used or prescribed incorrectly (irrational prescribing) then it can cause extensive losses in terms of health, economy and even for future generations. The emergence of pathogenic germs that are resistant to one antimicrobial resistance or certain types of antibiotics (multiple drug resistance) greatly complicates the treatment process.<sup>7</sup> The effort to find alternative materials that have no side effects is a solution to the above problems.

The correct alternative material to be the solution to the problem is herbal. Bandotan plants (*Ageratum conyzoides* L.) in Indonesia are classified as weeds so they are often destroyed. But some groups of our people use this plant as traditional medicine to cure various diseases.<sup>8</sup> Some of the uses of this plant are as a drug for new wounds, bleeding wounds, ulcers, eczema, and treating diseases caused by bacterial infections.

Bandotan leaves and flowers (*Ageratum conyzoides* L.) contain saponins, flavonoids, polyphenols, and essential oils. Phenol compounds are generally known as disinfectants used to kill pathogenic microorganisms. Polyphenol compounds have been shown to have antibacterial activity.<sup>9</sup> Phytochemical analysis of *Ageratum conyzoides* L. which has been done by Amadi et al shows that the main compounds in *Ageratum conyzoides* L. plants, namely alkaloids and flavonoids, accumulate on the leaves.<sup>10</sup> The usefulness and efficacy that can be obtained from this plant is very much, but so far there is still little research on Bandotan plants. Therefore this study aims to know the inhibitory test on the extract of bandotan plant (*Ageratum conyzoides* L.) on the growth of bacteria that cause periodontitis (*Porphyromonas gingivalis*).

### MATERIALS AND METHOD

This type of research used in this study is an experimental laboratory research. The research design used is post test only control group design using the dilution method and disk-diffusion method. This research was conducted at the Organic Chemistry Laboratory, Chemical Engineering Department, Ujung Pandang State Polytechnic and the Microbiology Laboratory Faculty of Medicine, Hasanuddin University. The study was conducted in March 2020.

There are three variables in this study, the independent variables are the leaves of the bandotan plant (*Ageratum Conyzoides* L.) starting from a concentration of 15%, 25%, 50%, 75%, 100%. and the dependent variable was the Minimum Inhibitory Concentration (MIC) and the inhibition zone of bacterial growth of *Porphyromonas gingivalis*. Control variables were time,

culture medium, and temperature. The research sample was *Porphyromonas gingivalis* bacteria that have been bred and the leaves of bandotan plant (*Ageratum Conyzoides* L.) that have been extracted by maceration method.

Antibacterial activity testing was done by disk-diffusion method or Kirby-Bauer diffusion method using discs. On the media which was dense then spread the bacteria as much as 0.02 mL, which has been adjusted to the 0.5 McFarland standards evenly using a spreader bar. The first MHA media was divided into seven sections, each of which was placed a disc containing bandotan leaf extract, metronidazole, and negative controls.

The treatment was carried out up to four times. Then incubated at 37°C for 1x24 hours and observed bacterial growth. The length of the inhibition zone formed was measured using calipers in millimeters. The data obtained were analyzed descriptively and displayed in tables and figures. Analysis of the results of research data was carried out with the One Way Anova test using the Statistical Package for the Social Sciences application (SPSS).

### Tools and Materials

Analytical scales, Test tubes, Test tube racks, Glass jars, Rotary evaporators, Petri dishes, Micropipettes, Paper disks, Caliper, Measuring cups, Incubators, Autoclaves, Tweezers, Bunsen, Filter paper, Erlenmeyer flasks, 1 ml spoit, Paper label, Funnel, Handscoon and mask, Vial bottle, GM Towel Bacterial isolates of *Porphyromonas gingivalis*, Bandotan leaves, Metronidazole, Medium Muller Hinton, Agar (MHA), Brain Heart Infusion Broth (BHIB) Media, Methylated spirit, 96% Ethanol, Aluminum foil, Sterile Aquades.

### RESULTS

**Table 1:** Results of Measurement of *Porphyromonas gingivalis* Bacterial Inhibition Zone Diameters

Type of Intervention	Concentration (%)	Inhibition (mm)				Average
		I	II	III	IV	
Leaves Extract Bandotan plants ( <i>Ageratum conyzoides</i> L.)	15	7,7	8,7	9,8	8,2	8,60±0,89
	25	9,5	9,8	10,9	12,6	10,70±1,40
	50	13,3	12,2	13,9	13,1	13,12±0,70
	75	14,3	14,2	14,9	15,3	14,67±0,51
	100	16,7	16,7	15,3	16,3	16,25±0,66
Control (+) Metronidazole		13,2	13,8	13,7	15,3	14,00±0,90
Control (-) Aquades		6,2	6,2	6,2	6,3	6,20±0,08

From Table 1. it can be seen descriptively that the extract of the bandotan plant leaf (*Ageratum conyzoides* L.) concentration of 15% has the lowest average inhibition zone of 8.60 ± 0.89 mm when compared to the entire concentration of the bandotan plant leaf extract (*Ageratum*

*conyzoides* L.). The inhibition zone at 100% concentration of bandotan leaves plant extract (*Ageratum conyzoides* L.) had the largest inhibition zone of 16.25 ± 0.66 mm.

**Table 2:** Test Results of Inhibition Zone Statistics of *Porphyromonas gingivalis* Bacteria

Type of Intervention	Concentration	N	Normality test*	Comparison test**
Leaves Extract	15	4	0.796	
Plants	25	4	0.467	
Bandotan	50	4	0.846	
( <i>Ageratum conyzoides L.</i> )	75	4	0.467	
	100	4	0.117	0.000
Control (+)Metronidazole		4	0.284	
Control (-) Aquades		4	0.683	

\*. The mean difference is significant at the 0.05 level.

Based on Shaphiro-Wilk statistical test results to know the normality value obtained  $p > 0.05$  which means the data was normally distributed thus the test was continued with the parametric test namely OnewayAnova (Table 2).

Based on the One-way Anova statistical test it was found that the significance value was 0,000 ( $p < 0.05$ ) which means that there are significant differences between treatment groups.

**Table 3:** Results of Post Hoc LSD (Least Significant Difference) Inhibition Zone of Porphyromonasgingivalis Bacteria (15%, 25% and 50%)

TreatmentGroup (I)	Comparison(J)	Mean Difference (I-J)	p-value
Concentration 15%	25%	-2.10	0.002
	50%	-4.52	0.000
	75%	-6.07	0.000
	100%	-7.65	0.000
	control+	-5.40	0.000
	control-	2.40	0.001
Concentration 25%	15%	2.10	0.002
	50%	-2.42	0.000
	75%	-3.97	0.000
	100%	-5.55	0.000
	control+	-3.30	0.000
	control-	4.50	0.000
Concentration 50%	15%	4.52	0.000
	25%	2.42	0.000
	75%	-1.55	0.015
	100%	-3.12	0.000
	control+	-0.87	0.150
	control-	6.92	0.000

\*. The mean difference is significant at the 0.05 level.

At a concentration of 15% the extracts of bandotan leaves plant (*Ageratum conyzoides L.*) when compared with the extracts of bandotan leaves plant (*Ageratum conyzoides L.*) the concentrations of 25%, 50%, 75%, 100%, positive and negative controls had  $p < 0, 05$  which means there are significant differences or have different effects.

At a concentration of 25% the extracts of bandotan leaves plant (*Ageratum conyzoides L.*) when compared with the extracts of bandotan leaves plant (*Ageratum conyzoides L.*) concentration of 15%, 50%, 75%, 100%, positive and

negative controls had a value of  $p < 0, 05$  which means there are significant differences or have different effects.

At a concentration of 50% the extracts of bandotan leaves plant (*Ageratum conyzoides L.*) when compared with the extracts of bandotan leaves plant (*Ageratum conyzoides L.*)

75% concentration had a  $p = 0.015 > 0.05$  and on a positive control that had a  $p \text{ value} = 0.150 > 0.05$  which means that all three have no significant difference or have the same effect. When compared to 15%, 25%, 100%, and negative controls with the value of  $p < 0.05$  which means there are significant differences or have different effects.

**Table 4:** Results of Post Hoc LSD (Least Significant Difference) Inhibition Zone of Porphyromonasgingivalis Bacteria (75% and 100%)

Treatment Group (I)	Comparison (J)	Mean Difference (I-J)	p-value
Concentration 75%	15%	6.07	0.000
	25%	3.97	0.000
	50%	1.55	0.015
	100%	-1.57	0.014
	control+	0.67	0.262
	control-	8.47	0.000
Concentration 100%	15%	7.65	0.000
	25%	5.55	0.000
	50%	3.12	0.000
	75%	1.57	0.014
	control+	2.25	0.001
	control-	10.05	0.000

\*. The mean difference is significant at the 0.05 level.

At a concentration of 75% the extracts of bandotan leaves plant (*Ageratum conyzoides* L.) when compared to the extracts of bandotan leaves plant (*Ageratum conyzoides* L.) concentration of 50% had a value of  $p = 0.015 > 0.05$ , 100% concentration had a value of  $p = 0.014 > 0.05$ , and positive control had a value of  $p = 0.262 > 0.05$  which means that all four have no significant difference or have the same effect. When compared with 15%, 25%, and negative controls with  $p < 0.05$  which means there are significant differences or have different effects.

At a concentration of 100% the extracts of bandotan leaves plant (*Ageratum conyzoides* L.) when compared with the extracts of bandotan leaves plant (*Ageratum conyzoides* L.) 75% concentration had a p value =  $0.014 > 0.05$ , which means that there are no significant differences or have the same significant effect. When compared to 15%, 25%, 50%, positive and negative controls had a p value  $< 0.05$  which means there are significant differences or have different effects.

**Table 5:** Results of Post Hoc LSD (Least Significant Difference) Inhibition Zone of *Porphyromonas gingivalis* Bacteria (Positive Control and Negative Control)

Treatment Group (I)	Comparison (J)	Mean Difference (I-J)	p-value
Control (+) Metronidazole	15%	5.40	0.000
	25%	3.30	0.000
	50%	0.87	0.150
	75%	-0.67	0.262
	100%	-2.25	0.001
	control-	7.80	0.000
Control (-) Aquades	15%	-2.40	0.001
	25%	-4.50	0.000
	50%	-6.92	0.000
	75%	-8.47	0.000
	100%	-10.05	0.000
	control+	-7.80	0.000

\*. The mean difference is significant at the 0.05 level.

In the positive control of metronidazole when compared with the extract of bandotan leaves plant (*Ageratum conyzoides* L.) the concentration of 50% had a value of  $p = 0.015 > 0.05$ , a concentration of 75% had a value of  $p = 0.262 > 0.05$ , which means that there are no significant differences or have the same effect. When compared with 15%, 25%, 100% and negative controls that had a  $p < 0.05$  which means there are significant differences or have different effects.

In negative control aquades when compared with concentrations of 15%, 25%, 50%, 75%, 100%, positive control and negative control had a  $p < 0.05$  which means there are significant differences or have different effects. Based on the results of the Post Hoc LSD test the inhibition zone between treatment groups on the average *Porphyromonas gingivalis* bacteria showed a significant value ( $p < 0.05$ ).



## DISCUSSION

In the study of inhibition zone test using the extracts of bandotan leaves plant (*Ageratum conyzoides* L.) on the growth of *Porphyromonas gingivalis*, it was seen that the inhibition zone formed increased in proportion to the increasing concentration of the extract. *Ageratum conyzoides* L. has many useful phytochemical compounds. It contains saponins, flavonoids, polyphenols, and essential oils.<sup>11,12,13</sup>

Phenol compounds are generally known as disinfectants used to kill pathogenic microorganisms. Flavonoids are the largest phenol compounds in nature and have been known to have biological activities as antioxidants, antimelanogenesis and antimutagenesis. Polyphenol compounds have been shown to have antibacterial activity. In addition, the leaves also contain essential oils and there are also coumarin.<sup>14,15</sup>

*Ageratum conyzoides* L. is known to have a good antibacterial effect. Antibacterial activity was demonstrated by the AC-1 component isolated from the leaves of *Ageratum conyzoides* L. In Mitra P's study, the AC-1 component of the bandotan plant showed an antibacterial effect on 4 types of gram-negative bacteria and 4 types of gram-positive bacteria. Based on research by Odeleye, et al., Ethanol extract of *Ageratum conyzoides* L. has potential antibacterial effects for use in medication.<sup>11,16,17</sup>

According to research conducted by Kamboj, antibacterial activity of aqueous extract were tested against three Gram-positive bacteria and seven Gram-negative bacteria and evaluated by the Piter paper disc diffusion method. Results showed a significant control of the growth of *A. viscolactis*, *K. aerogenes*, *B. cereus* and *S. pyogenes*. In an investigation for in vitro anti-methicillin-resistant *Staphylococcus aureus* (MRSA) activity found that MIC range of 55.4-71.0 mcg/ml were recorded for ethanol and water extracts of AC. The concentrations were too high to be considered active and AC were found to be ineffective in vitro in the study; therefore, suggest the immediate stoppage of their traditional use against MRSA-associated diseases in Lagos, Nigeria.<sup>18,19</sup>

According to research conducted by Budiman, the results show that both *Ageratum conyzoides* L. and *Piper Betle* L. extracts have antibacterial activity against *Staphylococcus aureus* with MIC value of 2 % and 5 %, respectively. Then, the gel containing 4 % sodium CMC showed the best physical stability, either containing *Ageratum conyzoides* L. or *Piper Betle* L. extract. The gel dosage forms of both extracts did not show any difference in organoleptic properties, pH and viscosity after 28 days storage. The gel dosage forms of *Ageratum conyzoides* L. and *Piper Betle* L. extracts have antibacterial activity with inhibition zone of 20.3 mm  $\pm$  1.3mm and 15.21  $\pm$  1.3 mm, respectively. The antibacterial activity of *Ageratum conyzoides* L. extract was higher compared to that of *Piper Betle* L. extract in the gel dosage form.<sup>20,21</sup>

According to research conducted by Garg, the results obtained from this study indicated that *A. conyzoides* shows potential antibacterials to againts five human pathogenic bacteria i.e gram + (*Bacillus subtilis*,

*Staphylococcus aureus*) and gram - (*Escherichia coli*, *Klebsiellapneumoniae* and *Pseudomonas aeruginosa*).<sup>22,23,24</sup>

Thus, bacterial activity can only be observed in diffusion method to see inhibition zones. Bandotan leaves plant extract (*Ageratum conyzoides* L.) at a concentration of 100% had the greatest inhibition zone of 16.25  $\pm$  0.66 mm and the inhibition zone was greater than the positive control of metrodinazole which was used as a comparative antibiotic from the extracts of bandotan leaves plant (*Ageratum conyzoides* L.). This research has proven that the extracts of bandotan leaves plant (*Ageratum conyzoides* L.) are effective as an inhibitory power against *Porphyromonas gingivalis*. Thus the research hypothesis was accepted.

## CONCLUSION

1. The extracts of Bandotan leaves plant (*Ageratum conyzoides* L.) are effective as an inhibitory against *Porphyromonas gingivalis*.
2. The extracts of Bandotan leaves plant (*Ageratum conyzoides* L.) at a concentration of 15% has the smallest average inhibition zone as much as 8.60  $\pm$  0.89 mm when compared to the entire concentration of bandotan leaves plant extract (*Ageratum conyzoides* L.). The inhibition zone at the concentration of 100% of the extracts of bandotan leaves plant (*Ageratum conyzoides* L.) has the biggest inhibition zone of 16.25  $\pm$  0.66 mm.
3. Minimal Inhibitory Concentration (MIC) of each of the extracts of bandotan leaves plant (*Ageratum conyzoides* L.) cannot be determined because the color of the extract is too concentrated thus it cannot be clearly seen the difference between clear and turbid tubes.

## REFERENCES

1. Dewi SRP, Marlamsya DO, Bikarindrasari R. (2017). Anti-caries effect of gambier extract in male wistar rats. *Dentino Journal*. 3(2): 83.
2. Newman GM, Takei HH, Klokkevold PR, Carranza FA. (2018). Newman and Carranza's Clinical Periodontology. 13th Ed. China: Elsevier. Pp. 62, 112, 342
3. Pratiwi EW, Praharani D, Arina YMD. (2015). Inhibitory power of papaya leaf extract (*Carica papaya* L.) against the adhesion of *Porphyromonas gingivalis* bacteria to neutrophils. *Medical Library Journal* 3(2): 194.
4. Newman GM, Takei HH, Klokkevold PR, Carranza FA. (2012). Carranza's Clinical Periodontology. 11th Ed. China: Elsevier. p. 41.
5. How KY, Song KP, Chan KG. (2016). *Porphyromonas gingivalis*: an overview of periodontopathic pathogen below the gum line. *Frontiers in Microbiology*. 7(53): 1, 4.
6. Lang NP, Lindhe J. *Clinical Periodontology and Implant Dentistry*. 6th Ed. Oxford: John Wiley & Sons. 2015. 837.

7. Utami ER, El-Hayah. (2011). Antibiotics, resistance, and therapeutic rationality. 1(4): 191.
8. Fitriani A. *Ageratum conyzoides* L. (2014). Alkaloid activity in the growth of staphylococcus aureus bacteria in vitro. Proceedings of the Research Symposium on Natural Medicinal Materials (SPBOA) XVI & Mukhtar XII PERHIPBA. p. 67.
9. Astuti H. (2015). Antibacterial activity test of ethanol extract and water extract of bandotan leaves (*Ageratum conyzoides* L.) on staphylococcus aureus and escherichia coli. Pharmaceutical Journal. 11(1): 291.
10. Safani EE, Kunharjito WAC, Lestari A, Purnama ER. (2019). Potential of bandotan leaf extract (*Ageratum conyzoides* L.) as a spray for the recovery of wound of diabetic mice infected with staphylococcus aureus. BIOTROPIC The Journal of Tropical Biology. 3(1): 69.
11. Melissa, Muchtaridi. (2017). Review: active compounds and pharmacological benefits of *ageratum conyzoides*. Pharmaceutical Journal. 15(1): 200-5.
12. Achmad H, Sri Ramadhany, Suryajaya FE. (2019). Streptococcus Colonial Growth of Dental Plaque Inhibition Using Flavonoid Extract of Ants Nest (*Myrmecodiapendans*): An in Vitro Study. Pesquisa Brasileira em Odontopediatria e Clínica Integrada. 19(1): e4250
13. Achmad, H. et al. (2018). Apoptosis Induction (Caspase-3,-9) and Human Tongue Squamous Cell Carcinoma VEGF Angiogenesis Inhibition using Flavonoid's Ethyl Acetate Fraction of Papua Ant Hill (*Myrmecodiapendans*). Journal of International Dental and Medical Research. 10(1): 276-84
14. Mardiana Adam, Achmad H. (2018). The Relationship of Mineral Fluor Exposure in Water with The Presence of Gingivitis (Study Case in Subdistrict of Tempe, Sengkang City, Wajo District). Journal of International Dental and Medical Research. (2) pp.470-476.
15. Djais, Arnilrawaty, et al. (2019). Differences of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of Moringa Leaf Extract (*Moringa Oleifera* L.) on Bacteria *Aggregatibacter Actinomycetemcomitans* and *Porphyromonas Gingivalis*. JIPHRD. 10(8): 896-900.
16. Adam, A. M., Thahir, H., & Achmad, H. (2019). Effect of *Channa Striata* Extract on the Expression of Tumor Necrosis Factor- $\alpha$  (TNF- $\alpha$ ) in Rat Experience Periodontitis. Indian JIPHRD. 10(9): 1391-1393.
17. Ramadhany S, Achmad H, Handayani H, Tanumihardja M, Singgih MF, Inayah NH, Ramadhany YF. (2020). Formulation of Ethanol Extract (*Myrmecodiapendans*) as an Antibacterial *Streptococcus mutans* in Chewable Lozenges for Children with Early Childhood Caries. Sys Rev Pharm. 11(4): 252-257.
18. Kamboj A, Saluja AK. (2008). *Ageratum conyzoides* L.: A review on its phytochemical and pharmacological profile. International Journal of Green Pharmacy. p.59-68. DOI: 10.4103/0973-8258.41171.
19. Achmad H, Tanumihardja M, Sartini, Ramadhany S, Singgih MF, Ramadhany YF, Mutmainnah N. (2020). Chewable Lozenges using White Shrimp Waste (*Litopenaeus vannamei*) in Reduce Colonization of Bacteria *Streptococcus mutans* in the Case of Early Childhood Caries. Sys Rev Pharm. 11(4): 293-299
20. Djais, A. I., Thahir, H., Hatta, M., Achmad, H, & Sari, M. (2019). Effect of Moringa Leaf Extract (*Moringa Oleifera*) on increasing the Number of Osteoblasts as a Marker of Bone Remodeling. JIPHRD. 2019;10(9):1394-1398.
21. Budiman A, Aulifa DL. (2020). A Study Comparing Antibacterial Activity of *Ageratum Conyzoides* L. Extract and Piper Betle L. Extract in Gel Dosage Forms Against *Staphylococcus Aureus*. Pharmacogn J. 12(3): 473-477.
22. Achmad H, Adam AM, Mappangara S, Oktawati S, Sjahril R, Singgih MF, Neormansyah I, Siswanto H. (2020). Identification and Antimicrobial Susceptibility of *Granulicatella adiacens* Isolated from Periodontal Pocket. Sys Rev Pharm. 11(4): 324-331.
23. Garg P, Grewal A. (2015). In Vitro Antibacterial Activity of *Ageratum Conyzoides* L. (Asteraceae). World Journal of Pharmacy and Pharmaceutical Sciences. 4(7): 893-897.
24. Achmad H, Thahir H, Rieuwpassa I, Mardiana AA, Oktawati S, Samad R, Djais AI, Gani A, Singgih MF, Madjid F, Admy SC. (2020). The Effectiveness of *Channa striata* Extract Antimicrobial Effect on Periopathogen Bacteria (*Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans*). Sys Rev Pharm. 11(4): 319-323.

# A Review of Bandotan Leaf Extract (*Ageratum conyzoides* L.) in Inhibition Test to the Growth of Bacteria (*Porphyromonas gingivalis*) Case of Periodontitis Disease

## ORIGINALITY REPORT

9%

SIMILARITY INDEX

6%

INTERNET SOURCES

6%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.greenpharmacy.info">www.greenpharmacy.info</a> Internet Source	3%
2	Submitted to Universitas Katolik Indonesia Atma Jaya Student Paper	1%
3	Submitted to Universitas Jenderal Achmad Yani Student Paper	1%
4	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	1%
5	<a href="http://hdl.handle.net">hdl.handle.net</a> Internet Source	1%
6	<a href="http://scienceandnature.org">scienceandnature.org</a> Internet Source	1%
7	<a href="http://konyvtar.uni-pannon.hu">konyvtar.uni-pannon.hu</a> Internet Source	<1%
8	Anton Bahtiar, Mutiara Nurazizah, Tirza Roselina, Anita Paulina Tambunan, Ade	<1%



Arsianti. "Ethanollic extracts of babandotan leaves ( *Ageratum conyzoides* L.) prevents inflammation and proteoglycan degradation by inhibiting TNF- $\alpha$  and MMP-9 on osteoarthritis rats induced by monosodium iodoacetate", *Asian Pacific Journal of Tropical Medicine*, 2017

Publication

---

9

Harel, Dipak, Dirk Schepmann, Helge Prinz, Reto Brun, Thomas J. Schmidt, and Bernhard Wünsch. "Natural Product Derived Antiprotozoal Agents: Synthesis, Biological Evaluation, and Structure–Activity Relationships of Novel Chromene and Chromane Derivatives", *Journal of Medicinal Chemistry*, 2013.

Publication

---

<1%

10

ROBERTO GHERZI. "The Human Homeodomain Protein OTX2 Binds to the Human Tenascin-C Promoter and Trans-Represses Its Activity in Transfected Cells", *DNA and Cell Biology*, 05/1997

Publication

---

<1%

11

Submitted to Federal University of Technology

Student Paper

---

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

