The Occurrences of Heavy Metals in Water, Sediment and Wild_Hayati J Biosciences 29 5 2022_Herliwati et al pdf

by Sp_jasa Cek Plagiarisme Wa: 085935293540

Submission date: 23-Nov-2022 07:35PM (UTC-0800)

Submission ID: 1962392255

File name: ment and Wild Hayati | Biosciences 29 5 2022 Herliwati et al.pdf (326.08K)

Word count: 3296
Character count: 16661



EISSN: 2086-4094

The Occurrences of Heavy Metals in Water, Sediment and Wild Shrimps Caught from Barito Estuary, South Kalimantan, Indonesia

Herliwati¹, Mijani Rahman¹, Achmad Syamsu Hidayat¹, Ulil Amri¹, Ika Sumantri²*

- ¹Faculty of Fishery and Marine Science, University of Lambung Mangkurat, Banjarbaru, Indonesia
- ²Department of Animal Science, Faculty of Agriculture, University of Lambung Mangkurat, Banjarbaru, Indonesia

ARTICLE INFO

Article history: Received December 19, 2019 Received in revised form December 22, 2021 Accepted January 18, 2022

KEYWORDS: Heavy metal, Toxic metal bioaccumulation. Wild shrimps

ABSTRACT

The research investigated the occurrences of heavy metal (Pb, Cu, and Cd) in waters, sediments, and wild shrimps collected from the Barito River estuary. Water and sediment samples were collected from 11 study sites by purposive sampling. At the same time, shrimps samples were captured around the sites. The study showed low levels of heavy metals in water samples; those were: Cd<0.0019 mg/L, Cu<0.001 mg/L, and Pb<0.0019 mg/L. Low levels of heavy metals were also detected in sediment samples, those were Cd <0.24 mg/kg, Cu <0.013-0.69 mg/kg, and Pb<0.024 mg/kg. Heavy metals contaminations were detected in Parapenaeopsis sculptilis, those were Cd = 2.802 mg/kg; Cu = 3.399 mg/kg, and Pb = 1.294 mg/kg. In Acetes japonicus, the heavy metals concentration were Cd = 2.127 mg./kg, Cu = 5.518 mg/ kg, and Pb = 2.723 mg/kg. In Penaeus merguiensi, the heavy metals concentrations were Cd = 8.598 mg/kg, Cu = 6.403 mg/kg, and Pb = 5.433 mg/kg. This study indicated the increases of heavy metals concentrations from water to sediment and finally into the shrimps. The presence of heavy metals in shrimps indicated the bioaccumulation of toxic metals, especially for Pb and Cd concentrations which exceeded the tolerable limit according to JECFA.

1. Introduction

Barito River is one of the major rivers in the South Kalimantan and Central Kalimantan Provinces. This main river of Barito flows from Muller Mountain into the Java Sea, and it has a length of about 900 km, a wide average of about 800 m, and a depth average of about 8 m (Arisanty and Saputra 2017). The role of the Barito River for human life is increasing in line with the population increase. An increase in population and various activities can cause an increase in the amount of waste, including those containing heavy metals (Khan et al. 2018).

As environmental pollutants and potentially toxic materials, many heavy metals have drawn great attention worldwide (Liu et al. 2009; Montuori et al. 2013; Zhang et al. 2012). Water containing heavy metals is very harmful to human health. Heavy metals that enter the water will undergo various processes, including transport by tidal currents, dilution, associated with suspended material, coagulation and sedimentation to the bottom, associated with organic sediment material, absorbed by plankton, and in

* Corresponding Author E-mail Address: isumantri@ulm.ac.id turn, will enter the food chain (Eisler 1986; Pandey et al. 2003). Human activity has a major impact on the acceleration of heavy metals accumulation in the food chain. World Health Organization estimated 60-80% of 60 to 80% of the body burden of toxic metals in people living in industrialized or urbanized areas is mainly caused by the intake of metals via food consumption (Pullina et al. 2014).

The concentrations of heavy metals will increase if urban, agricultural, and industrial wastes which contain these pollutants enter the sea through rivers and settle to the bottom of the waters, which eventually become toxic to marine organisms (Hawker and Connell 1992). According to Yu et al. (2011), metals in sediments can be in various forms and bonds, among others, as free ions and bonded to carbonates, which are very unstable; thus, they are easily released into the water then absorbed by organisms.

The occurrences of heavy metals in river mouth ecology are a threat to aquatic organisms and human health. Some heavy metals, such as Cd and Pb, induce inflammation and increase free radicals and oxidants that damage various organs, including the liver and kidneys (Suhartono et al. 2015). This study aims to investigate the concentrations of heavy metals (Pb, Cu,

644 Herliwati et al.

and Cd) in water and sediments and the occurrence of these heavy metals in the shrimps captured around the Barito estuary. Wild shrimps were chosen as samples of aquatic organisms in this present study because shrimp is part of net food connected with water and sediment as its habitat.

2. Materials and Methods

2.1. Study Area and Sampling Site

This research was conducted at the Barito estuary, a traffic lane for coal cargo ships and other cargo vessels. Samplings of heavy metals in water, biota, and sediments were carried out in the waters of the Barito estuary. The number of water sampling points is set up at 11 points spreading in the river mouth. Whereas for sediment, samplings were conducted at 5 points. Water and sediment samples were taken directly at the research location, while shrimps samples were obtained from fishers who captured shrimp around the Barito estuary. The location of the study is illustrated in Figure 1.

2.2. Sample Preparation and Analysis

Five hundred milliliters of water samples from each sampling site were filtered through Whatman filter paper no. 42. Detailed preparation of water samples and determination of Pb, Cu, and Cd using atomic absorption spectrophotometry (Perkin Palmer 3110) were conducted according to the protocol of Indonesian Standard of Industry (SNI) no. 6989-16:2009 for Pb, no. 6989-6:2009 for Cd, and no. 6989-8:2009 for Cu.

In the laboratory, sediment samples were dried at 80°C to constant weight and then ground and filtered through a 63 um sieve. Detailed preparation of sediment samples and determination of Pb, Cu, and Cd using atomic absorption spectrophotometry (Perkin Palmer 3110) were conducted according to the protocol of BBTKL no. IKM.FR.14.BBTKL.

Shrimps samples were dried at 105°C to constant weight, and then the samples were ground and sieved. Detailed preparation of shrimps samples and determination of Pb, Cu, and Cd using atomic



Figure 1. Site sampling for water and sediment in the Barito river estuary

absorption spectrophotometry (Perkin Palmer 3110) were conducted according to the protocol of Indonesian Standard of Industry (SNI) no. 6989-82:2018.

2.3. Data Analysis

Data were analyzed statistically to determine the mean and standard deviation. Safety of shrimp consumption (crustaceans) in the Barito estuary waters is based on the calculation of PTWI (Provisional Torelable Weekly Intake), which is the maximum temporary amount of a substance in milligrams per kilogram of body weight that can be consumed in a week without causing adverse effects on health. Determination of pollution status of heavy metal content to the level of heavy metal pollution is done by Metal Pollution Index (MPI) based on the formulas of Usero et al. (1997) and Giusti et al. (1999).

3. Results

3.1. The Concentration of Heavy Metals in Water

Results showed water samples collected from 11 sites of Barito river estuary containing a low concentration of heavy metals, namely Cd <0.0019 mg/L, Cu <0.001 mg/L, and Pb <0.0019 mg/L (Table 1).

Table 1. Heavy metals concentration in water samples (mg/L)

(IIIg/L)			
Sampling site	Heavy metal (mg/L)		
Sampling site	Cd	Cu	Pb
03°27'00,1" S-114°30'56,1"	<0.0019	<0.001	<0.0019
03°28'24,1" S-114°30'23,7"	< 0.0019	< 0.001	<0.0019
03°30'22,7" S-114°28'27,3"	< 0.0019	< 0.001	<0.0019
03°33'57,2" S-114°31'12,2"	< 0.0019	< 0.001	<0.0019
03°31'02,0" S-114°22'29,1"	< 0.0019	< 0.001	<0.0019
03°32'01,0" S-114°24'56,9"	< 0.0019	< 0.001	<0.0019
03°34'50,2" S-114°29'15,9"	< 0.0019	< 0.001	<0.0019
03°39'58,6" S-114°28'25,4"	< 0.0019	< 0.001	<0.0019
03°39'58,6" S-114°28'35,9"	< 0.0019	< 0.001	<0.0019
03°34'46,1" S-114°31'26,7"	< 0.0019	< 0.001	<0.0019
03°33'16,7" S-114°31'27,7"	<0.0019	< 0.001	<0.0019

3.2. The Concentration of Heavy Metals in Sediment

Analysis of sediment samples showed that Cadmium, Copper, and lead were not detected in three of five sampling sites. Those heavy metals were only found in one site, whereas Cadmium was found in two sites (Table 2).

3.3. The Concentration of Heavy Metals in Shrimps

Determination of heavy metals in the biota (shrimps) showed increases in Cadmium, Copper, and lead concentrations than in water and sediment. As shown in the Table 3, the average levels for each heavy metal in the shrimps were Pb = 3.150 mg/kg, Cu = 5.107 mg/kg, and Cd = 4.509 mg/kg.

Determination of heavy metals in shrimps species captured around sampling sites indicated an increase in the levels of heavy metals from water to the sediments and finally in the biota of sediment (shrimp). The presence of heavy metals in shrimps suggested the occurrence of bioaccumulation of this toxic substance, as shown in Figure 2.

Table 2. Heavy metals concentration in sediment samples (mg/L)

Sampling site	Heavy metal (mg/L)		
Sampling site	Cd	Cu	Pb
03°27'00,1" S-114°30'56,1"	0.00	0.00	0.00
03°28'24,1" S-114°30'23,7"	0.61	0.26	0.69
03°30'22,7" S-114°28'27,3"	0.00	0.00	0.00
03°33'57,2" S-114°31'12,2"	0.35	0.00	0.00
03°31'02,0" S-114°22'29,1"	0.00	0.00	0.00

Table 3. The concentration of heavy metals in shrimps samples (mg/kg)

Shrimp species	Heavy metal (mg/L)			
Silitinp species	Cd	Cu	Pb	
Penaeus merguiensi	8.598±0.749	6.403±1.079	5.433±0.685	
Parapenaeopsis sculptilis	2.802±0.084	3.399±0.118	1.294±0.086	
Acetes japonicus	2.127±0.041	5.518±0.083	2.723±0.027	

646 Herliwati et al.




Figure 2. Heavy metals concentrations in water (mg/L), sediment (mg/kg), and shrimps (mg/kg) of Barito river

4. Discussion

Various human activities result in heavy metals accumulation in the water, such as industrial, agricultural, and domestic sewage (Liu et al. 2016). This study found low Cadmium, Copper, and lead concentrations in water samples of the Barito river mouth. This finding was in agreement with the statement of Gaur et al. (2005) that only a small portion of free heavy metal ions can remain in the dissolved phase in aquatic systems, as most parts settle in the sediment by adsorption, hydrolysis, co-precipitation, and other physical and chemical processes. Sofarini et al. (2010) found heavy metals in water samples of Barito River are low, except for containing Hg (0.28 mg/L) and Pb (0.18 mg/L). Sediments samples analysis indicated the locality and low levels of heavy metals occurrences in the Barito river's sediments. Only from a site sampling was containing all of the studied heavy metals. Previous studies by Sofarini et al. (2010) notified accumulation of heavy metals, especially Cd levels, in the sediment of Barito River. The occurrence of heavy metals in the sediment was worrying considering the possible bioaccumulation of heavy metals in the river bed biota. Further analysis showed the concentrations of heavy metals in the biota (shrimps) were higher than the concentrations found in the water and sediment. The average levels for each heavy metal in the shrimp were Pb = 3.150 mg/kg, Cu = 5.107 mg/kg, and Cd =

4.509 mg/kg. This result is in agreement with the study of Rahman (2006) that found high contamination of Pb (42-105 ml/kg) and Cd (8-17 mg/kg) in crustacean samples collected from the coastal area connected to the Barito River mouth. However, the levels of these three heavy metals were still low compared to Ahmad et al. (2010) in the waters of the Buriganga River, Bangladesh.

Several aquatic biotas have been used for biomonitoring purposes. These organisms accumulate heavy metals in their bodies and are sensitive to chemical pollutions as they directly contact the polluted part of the habitat (Khan *et al.* 2018). Determination of heavy metals in shrimps species captured around sampling sites indicated accumulation of heavy metals from water to the sediments and finally in the biota of sediment (shrimp). The occurrences and levels of Cadmium, Copper, and Lead in shrimps of this study were in harmful levels for human consumption because they have exceeded the tolerance threshold (Provisional Torelable Weekly Intake) according to JECFA (2010) in CCCF (2011), namely 0.025-0.050 mg Pb per kg body weight and 25 µg Cd per kg body weight per month.

The anthropogenic release has been considered the main source of heavy metals accumulation in water, sediment, and aquatic organisms (Savorelli et al. 2017; Aliko et al. 2018). The consumption of these metals at high levels could adversely impact human health. High exposure to Cadmium through ingestion causes immediate poisoning and damage to the liver and

kidneys. In addition, compounds containing Cadmium are also carcinogenic (El-Moselhy et al. 2000). Excessive intake of Copper might cause vomiting, hematemesis (vomiting blood), and gastrointestinal distress (Arvind 2002). Children are the most vulnerable group to the toxic effects of Lead and can suffer permanent adverse health effects, particularly the brain and nervous system development. In adults, lead causes long-term harm, including increased risks of high blood pressure, kidney damage, and neurological effects (Voegborlo et al. 2012).

In conclusion, this study indicated that the occurrences in the low levels of heavy metals, particularly Cadmium, Copper, and Lead in the water and sediment of the Barito river, might result in bioaccumulation of these toxic substances into the wild shrimps to reach the harmful levels for human consumption. Therefore any efforts have to be initiated to reduce these heavy metals levels in the human food chain.

Acknowledgements

The authors would like to thank the University of Lambung Mangkurat for supporting the research through the PNBP Fundamental Research fund.

References

- Ahmad, M.K., Islam, S., Rahman, S., Haque, M.R., Islam, M.M., 2010. Heavy metals in water, sediment and some fishes of Buriganga river, Bangladesh. *International Journal of Environmental Research*. 4, 321–332.
- Aliko, V., Qirjo, M., Sula, E., Morina, V., Faggio, C., 2018. Antioxidant defense system, immune response and erythron profile modulation in Gold fish, Carassius auratus, after acute manganese treatment. Fish Shellfish Immunology. 76, 101-109. https://doi. org/10.1016/j.fsi.2018.02.042
- Arisanty, D., Saputra, A.N., 2017. Remote sensing studies of suspended sediment concentration variation in Barito Delta. *IOP Conf. Series: Earth and Environmental Science.* 98, 012058. https://doi.org/10.1088/1755-1315/98/1/012058
- Arvind, K., 2002. Ecology of Polluted Waters, A.P.H Publishing corporation, Ganja-New Delhi.
- CCCF., 2011. Working Document for Information and Use In Discussion Related to Contaminants and Toxins in the GSCTFF, Joint FAO/WHO Food Standards Programme Codex Committee on Contaminant in Foods, Fifth Session, The Hague.
- Session, The Hague.
 Eisler, R., 1986. Zinc hazards to fish, wildlife and invertebrates:
 a synoptic review. Reproductive Biology. 85, 1–6.
 El-Moselhy, K.M., 2000. Accumulation of Copper, Cadmium
 and lead in some fish from the Gulf of Suez. Egyptian
 Journal of Aquatic Biology and Fisheries. 1, 13–18.
 Gaur, V.K., Gupta, S.K., Pandey, S.D., Gopal, K., Misra, V., 2005.
 Distribution of heavy metals in sediment and water
 of Privar Combi. Environ Monit Assess 102, 410, 422.
- of River Gomti, Environ, Monit, Assess, 102, 419-433.
- https://doi.org/10.1007/s10661-005-6395-6 Giusti, L., Williamson, A.C., Mistry, A., 1999. Biologically available trace metals in Mytilus edulis from the coast of Northern England. Environmental International. 25, 969-981. https://doi.org/10.1016/ S0160-4120(99)00066-5
- Hawker, D.W., Connell, D.W., 1992. Pollution in Tropical Aquatic Systems, CRC Press, Inc., London.

- [JECFA] Joint Expert Committee on Food Additives, 2010. FAO JECFA Monograph: Compendium of Food Additive Specifications, JECFA 73rd Meeting 2010, FAO/WHO,
- Khan, M.I., Khisroon, M., Khan, A., Gulfam, N., Siraj, M., Zaidi, F., Ahmadullah, A., Fatima, S.H., Noreen, S., Hamidullah, S.Z.A., Qadir, F., 2018. Bioaccumulation of heavy metals in water, sediments, and tissues and their histopathological effects on *Anodonta cygnea* (Linea, 1876) in Kabul River, Khyber Pakhtunkhwa, Pakistan. *BioMed Research International.* 6, 1-10. https://doi.org/10.1155/2018/1910274

 Liu, J.L., Li Y.L., Zhang, B., Cao, J.L., Cao, Z.G., Domagalski, J., 2009. Ecological risk of heavy metals in sediments of the Luan Riversource water. *Ecotoxicology*, 18, 748–758. https://doi.org/10.1007/s10646-009-0345-y

 Liu, X.B., Li, D., Song, G., 2016. Assessment of heavy metal levels in surface sediments of estuaries and adjacent
- levels in surface sediments of estuaries and adjacent coastal areas in China. Front. Earth Sci. 11, 85-94. https://doi.org/10.1007/s11707-016-0569-0 Montuori, P., Lama, P., Aurino, S., Naviglio, D., Triassi, M., 2013.
- Metals loads into the Mediterranean sea: estimate of Sarno River inputs and ecological risk. Ecotoxicology 22, 295-307. https://doi.org/10.1007/s10646-012-1026-9
- Pandey, S., Parvez, S., Sayeed, I., Haque, R., Bin-Hafeez, B., Raisuddin, S., 2003. Biomarkers of oxidative stress: a comparative study of river Yamuna fish *Wallago attu* (Bl. and Schn.). *Science of the Total Environment.* 309, 105–115. https://doi.org/10.1016/S0048-9697(03)00006-8
- Pullina, G., Battacone, G., Brambilla, G., Cheli, F., Danieli, P.P., Masoero, F., Pietri, A., Ronchi, B., 2014. An update on the safety of foods of animal origin and feeds. *Italian Journal of Animal Science*, 13, 845-856. https://doi.org/10.4681/lijes.2014.2571
- org/10.4081/ijas.2014.3571 Rahman, A., 2006. Kandungan Cd Timbal (Pb) dan Kadmium (Cd)pada beberapa jenis krustasea di Pantai Batakan dan Takisung Kabupaten Tanah laut Kalimantan Selatan. *Bioscientiae* 3, 93-101.
- Savorelli, F., Manfra, L., Croppo, M., Tornambè, A., Palazzi, D., Canepa, S., Trentini, P.L., Cicero, A.M., Faggio, C., 2017. Fitness evaluation of *Ruditapes philippinarum* exposed to Ni. Biology Trace Element Research. 177 384-393. https://doi.org/10.1007/s12011-016-0885-y Sofarini, D., Abdurrahman, Ichsan, R., 2010. Studi analisis
- pengujian Cd pada badan air, biota, dan sedimen di Perairan Muara DAS Barito. *Jurnal Bumi Lestari*. 10,

- 28-37.

 Suhartono, E., Triawanti, Leksono, A.S., Djati, M.S. 2015.
 Cadmium exposure on rat kidney: lipid peroxidation and chlorinative stress. J. Exp. Life Science. 5, 1-5. https://doi.org/10.21776/ub.jels.2015.005.01.01

 Usero, J., Gonzales-Regalado, E., Gracia, I., 1997. Trace metals in bivalve mollusks Ruditapes decussatus and Ruditapes philippinarum from the Atlantic coast of southern Spain. Environment International. 23, 291-298. https://doi.org/10.1016/S0160-4120(97)00030-5

 Voegborlo, R.B., Atta, A., Agorku, E.S., 2012. Total mercury distribution in different tissues of six species of freshwater fish from the Kpong hydroelectric reservoir in Ghana. Environmental Modeling and Assessment. 184, 3259–3265. https://doi.org/10.1007/s10661-011-2186-4

 Yu, X., Yana, Y., Wang, W., 2011. The distribution and
- Yu, X., Yana, Y., Wang, W., 2011. The distribution and speciation of trace metals in surface sediments from the Pearl River Estuary and the Daya Bay, Southern China. Marine Pollution Bulletin. 60, 1364–137. https:// doi.org/10.1016/j.marpolbul.2010.05.012
- Zhang, G.S., Liu, D.Y., Wu, H.F., Chen, L.F., Han, Q.X., 2012. Heavy metal contamination in the marine organisms in Yantai coast, northern Yellow Sea of China. Ecotoxicology. 21, 1726–1733. https://doi.org/10.1007/ s10646-012-0958-4

The Occurrences of Heavy Metals in Water, Sediment and Wild_Hayati J Biosciences 29 5 2022_Herliwati et al pdf

	_ , ,	Biosciences 29 5	2022_Herliwa	ti et al pdf	_
ORIGINA	ALITY REPORT				—
SIMILA	8 ARITY INDEX	11% INTERNET SOURCES	13% PUBLICATIONS	2% STUDENT PAPERS	
PRIMAR	Y SOURCES				
1	Stojic, M "Evaluat	Ailenkovic, Jelen Iira Pucarevic, S ion of heavy me nd seafood prod	nezana Strbacetals and radio	c. onuclides	%
2	WWW.UN			1	%
3	kuojs.lib Internet Sourc	.ku.ac.th		1	%
4	samafin Internet Source	d.sama.gov.sa		1	%
5	www.km	nae-journal.org		1	%
6	publishir Internet Source	ngrealtime.com		1	%
7		ashid, Rouf Ahr Iohammad Ane	•		%

Shafiq-ur-Rehman. "Environmental and

socioeconomic factors induced blood lead in children: an investigation from Kashmir, India", Environmental Monitoring and Assessment, 2019

Publication

Submitted to Mahidol University <1% 8 Student Paper Submitted to iGroup Student Paper Ahmed Moustafa Nemr. "Total and Leachable 10 Heavy Metals in Muddy and Sandy Sediments of Egyptian Coast Along Mediterranean Sea", Environmental Monitoring and Assessment, 05/08/2007 Publication Mohiuddin, KM, MM Alam, I Ahmed, HM Zakir, <1% 11 and AK Chowdhury. "Physicochemical Properties and Metallic Constituent Load in the Water Samples of the Buriganga of Bangladesh", Journal of Environmental Science and Natural Resources, 2016. Publication Submitted to The Open University of Hong <1% Kong Student Paper bbrc.in Internet Source

Debo Zhao, Shiming Wan, Zhaojie Yu, Jie Huang. "Distribution, enrichment and sources of heavy metals in surface sediments of Hainan Island rivers, China", Environmental Earth Sciences, 2015

<1%

- Publication
- Haifeng Xiao, Shuying Zang, Ying Guan, Shaojun Liu, Yan Gao, Qingzhan Sun, Haifeng Xu, Miao Li, Jingjing Wang, Xueyuan Pei.
 "Assessment of potential risks associated with heavy metal contamination in sediment in Aobaopao Lake, China, determined from sediment cores", Ecotoxicology, 2014

<1%

Publication

Joyce Ayuba Ramadan, Ahmad Isah Haruna.
"Health Risk Assessment from Exposure to
Heavy Metals in Surface and Groundwater
Resources within Barkin Ladi, North Central
Nigeria", Journal of Geoscience and
Environment Protection, 2019

<1%

- Publication
- Teresa K. Pong, John Besida, Thomas A.
 O'Donnell, David G. Wood. "A Novel Fluoride
 Process for Producing TiO2 from Titaniferous
 Ore", Industrial & Engineering Chemistry
 Research, 2002

<1%

19	kundoc.com Internet Source	<1%
20	recentscientific.com Internet Source	<1%
21	shodhganga.inflibnet.ac.in Internet Source	<1%
22	"Bioaccumulation of Pb and Cd on Broiler Chicken Fed in Difference Diets", 'Faculty of Engineering Diponegoro University'	<1%
23	Fahri Karayakar. "Seasonal Variation in Copper, Zinc, Chromium, Lead and Cadmium Levels in Hepatopancreas, Gill and Muscle Tissues of the Mussel Brachidontes pharaonis Fischer, Collected along the Mersin Coast, Turkey", Bulletin of Environmental Contamination and Toxicology, 09/05/2007 Publication	<1%
24	eprints.lib.hokudai.ac.jp Internet Source	<1%
25	www.fjs.fudutsinma.edu.ng Internet Source	<1%
26	A.S. Siregar, N.A. Prayogo, T Harisam. "THE ACCUMULATION OF HEAVY METALS KADMIUM (Cd) IN WATER, SEDIMENTS AND	<1%

AQUACULTURE BIOTA WHICH CONTAMINATED BY BATIK WASTE IN MULYOREJO VILLAGE PEKALONGAN", IOP Conference Series: Earth and Environmental Science, 2019

Publication

El-Shazly, Mohamed M., Wael A. Omar, Yusuf A. Edmardash, Mona Sayed Ibrahim, Emad I. Elzayat, Iman I.A. El-Sebeay, Khaled M. Abdel Rahman, and Mustafa M. Soliman. "Area reduction and trace element pollution in Nile Delta wetland ecosystems", African Journal of Ecology, 2016.

<1%

Publication

Lulu Mao, Xitao Liu, Zongxing Wang, Baodong Wang, Chunye Lin, Ming Xin, Bo-Tao Zhang, Tingting Wu, Mengchang He, Wei Ouyang.
"Trophic transfer and dietary exposure risk of mercury in aquatic organisms from urbanized coastal ecosystems", Chemosphere, 2021

<1%

Maryam Askari, Seyyed Reza Mousavi Harami, Mohammad Reza Noora. "Environmental geochemistry of heavy metals in coral reefs and sediments of Chabahar Bay", Results in Engineering, 2022

<1%

Publication

Md. Kawser Ahmed, Mohammad Abdul Baki, Md. Saiful Islam, Goutam Kumar Kundu et al. "Human health risk assessment of heavy metals in tropical fish and shellfish collected from the river Buriganga, Bangladesh", Environmental Science and Pollution Research, 2015

<1%

Publication

Md. Wahidul Alam, Mohammad Mostafizur Rahman, Md. Simul Bhuyan, Venkatramanan Senapathi et al. "Inferences on metal pollution in the natural spawning zone of Bangladesh river and pollution management strategies", Environmental Monitoring and Assessment, 2022

<1%

Publication

Millicent Kekana, Abraham Addo-Bediako,
Antoinette Jooste, Lieven Bervoets.
"Accumulation of Trace Metal(loid)s in Fish
Muscle Tissue From the Groot Letaba River,
South Africa", Polish Journal of Environmental
Studies, 2022

<1%

Publication

Patcharin Ruchuwararak, Somsak Intamat, Bundit Tengjaroenkul, Lamyai Neeratanaphan. "Bioaccumulation of heavy metals in local edible plants near a municipal landfill and the related human health risk

<1 %

assessment", Human and Ecological Risk Assessment: An International Journal, 2018

34	article.sciencepublishinggroup.com Internet Source	<1%
35	bnrc.springeropen.com Internet Source	<1%
36	jels.ub.ac.id Internet Source	<1%
37	mzuir.inflibnet.ac.in Internet Source	<1%
38	psasir.upm.edu.my Internet Source	<1%
39	repository.unhas.ac.id Internet Source	<1%
40	www.foodandnutritionjournal.org Internet Source	<1%
41	www.saulibrary.edu.bd Internet Source	<1%
42	Li, Jing. "Risk Assessment of Heavy Metals in Surface Sediments from the Yanghe River, China", International Journal of Environmental Research and Public Health, 2014.	<1%



Shengguang Yuan, wenqiang zhang, Wenye Li, Zhenhan Li, Minshan Wu, Baoqing Shan. "Accumulation and Potential Ecological Risk of Heavy Metals in the Sediments of Rivers System in Beijing-Tianjin Area", Research Square Platform LLC, 2021

<1%

Publication

Exclude quotes Off
Exclude bibliography On

Exclude matches

Off

The Occurrences of Heavy Metals in Water, Sediment and Wild_Hayati J Biosciences 29 5 2022_Herliwati et al pdf

GRADEMARK REPORT	
FINAL GRADE	GENERAL COMMENTS
/0	Instructor
, •	
PAGE 1	
PAGE 2	
PAGE 3	
PAGE 4	
PAGE 5	