

Cytotoxic Activity of Volatile Compounds in *Cymbopogon nardus*' Essential Oils

^{1*}Rizki Fitria, ²Djarot Sasongko Hami Seno, ³Bambang Pontjo Priosoeryanto, ⁴Najmah,
^{2,5}Waras Nurcholis

¹Department of Chemistry, Lambung Mangkurat University, Indonesia

²Department of Biochemistry, IPB University, Indonesia

³Department of Veterinary Clinic Reproduction and Pathology, IPB University, Indonesia

⁴Department of Chemistry, Universitas Negeri Gorontalo, Indonesia

⁵Tropical Biopharmaca Research Center, IPB University, Indonesia

rizki.fitria@ulm.ac.id , djarotse@apps.ipb.ac.id , bpontjo4@gmail.com , naimahkim@ung.ac.id ,
wnurcholis@apps.ipb.ac.id

ARTICLE INFO

Article History:

Diterima : 29-07-2022
Disetujui : 17-09-2022

Keywords:

Cytotoxic; Volatile;
Cymbopogon nardus;
Essential Oils; MCF-7



ABSTRACT

Abstract: Plants produce a variety of secondary metabolites, one of which is essential oils that contain a variety of volatile compounds and are useful for humans. *Cymbopogon nardus* contains volatile compounds that can inhibit the proliferation of cancer cells. This research aims to explore the antiproliferation activity of *C.nardus*' essential oils with different concentrations against breast cancer Michigan Cancer Foundation-7 (MCF-7) cells. Antiproliferation test was carried out with total cells method using trypan blue and cells were calculated using a microscope. Gas Chromatography-Mass Spectrometer (GC-MS) was performed to identify the volatile compounds. The results showed that the Inhibitory Concentration 50 (IC_{50}) value was 359.6 ppm with an inhibition percent of 44.49% at 170 ppm. Meanwhile, inhibition percent against Vero normal cell was 29.04%, compared to Doxorubicin 35.23%. The dominant volatile compound in *C. nardus*' essential oil were Geraniol and Citronellol.

Abstrak Tanaman menghasilkan berbagai macam metabolit sekunder, salah satunya dalam bentuk minyak atsiri yang mengandung berbagai macam senyawa volatil yang berguna bagi manusia. *Cymbopogon nardus* mengandung senyawa volatil yang dapat menghambat proliferasi sel kanker. Penelitian ini bertujuan untuk mengetahui aktifitas antiproliferasi minyak atsiri *C. nardus* dengan beberapa konsentrasi terhadap sel kanker payudara Michigan State Foundation-7 (MCF-7). Uji antiproliferasi dilakukan dengan metode total sel menggunakan *trypan blue* dan sel dihitung menggunakan mikroskop. *Gas Chromatography-Mass Spectrometer* (GC-MS) dilakukan untuk mengidentifikasi senyawa volatil. Hasil menunjukkan bahwa konsentrasi penghambatan 50 (IC_{50}) sebesar 359.6 ppm dengan persen penghambatan sebesar 44.49% pada konsentrasi 170 ppm. Sementara itu, persen ihibisi terhadap sel normal Vero sebesar 29.04%, dibandingkan dengan Doktorubisin yang sebesar 35.23%. Senyawa volatil dominan yang terdapat pada *C. nardus* adalah Geraniol dan Citronellol.



<https://doi.org/10.31764/justek.vXiY.ZZZ>



This is an open access article under the CC-BY-SA license