

Investigation of parameters in restaurant food waste for use as poultry rations

by Mohammad Rudiansyah

Submission date: 19-May-2022 01:16PM (UTC-0400)



Submission ID: 1839973082

File name: Investigation_of_parameters.pdf (1.03M)

Word count: 4303

Character count: 22018

4 Investigation of parameters in restaurant food waste for use as poultry rations

Yudi GARNIDA¹, Mohammad RUDIANSYAH^{2*} , Ghulam YASIN³, Trias MAHMUDIONO^{4*} ,
Abed Jawad KADHIM⁵, Sandhir SHARMA⁶, Hussein Ali HUSSEIN⁷, Rustem Adamovich SHICHIYAKH⁸,
Walid Kamal ABDELBASSET^{9,10}, Acim Heri ISWANTO¹¹

Abstract

One way to use food industry wastes is to feed the livestock and poultry. Applying appropriate management methods for the optimal use of agricultural waste in the preparation of livestock and poultry feed increases productivity and reduces damage to agricultural and conversion industries and the use of new resources in animal and poultry feed. In this paper, in order to investigate the possibility of recycling food waste as poultry feed, the waste produced in a restaurant has been used. The parameters of fat percentage were measured using Soxhlet extractor, percentage of dry matter using the oven, percentage of protein in it using Kjeldahl method, and also the percentage of calcium and phosphorus with the standard beard. Comparing them with the usual diet used in poultry feed shows that the average values of calcium, protein, and fat content of food waste are 0.14%, 5.65%, 7.34% higher than the average diet of poultry, respectively. Also, the average amount of phosphorus and dry matter of food waste is 0.48% and 60% less than the average diet of poultry, respectively.

Keywords: waste management; restaurant food waste; poultry rations; recycling.

Practical Application: In the current study the parameters in restaurant food waste has been investigated in order to use as poultry feed.

1 Introduction

Food waste is produced especially in the agricultural, industrial, and urban sectors of society, including by wholesalers and consumers (Seidavi et al., 2020; Molajou et al., 2021). Utilization of such wastes requires urgent research as recycling and reduction of residues can reduce pollution and improve the current situation by preparing new feed from the residues (Chen et al., 2020; Barreira et al., 2021). The use of animal waste, summer crops, and fruits in animal and poultry feed for indirect production of feed from food waste is very valuable (Kim et al., 2020). Food leftover from restaurants, homes, or other establishments is often disposed of as waste due to its high humidity and perishable nature (Ojha et al., 2020; Burzyńska, 2019). It causes water and soil pollution in landfills (Sharma et al., 2020). They may also contaminate other wastes and disrupt the incineration of other wastes. As a result, management programs are needed to recycle waste (Olszewska et al., 2020) properly. Due to rapid population growth and increasing demand for food, the amount of food waste produced in urban communities is increasing rapidly (Chen et al., 2012; Galhardo et al., 2021).

Various management methods are being implemented for food waste recycling, including composting, anaerobic digestion, burning or feeding livestock and poultry (Ramos & Nascimento, 2020). In livestock and poultry breeding centers, most of the operating costs are related to the cost of feed supply, so using cheap feed can reduce production costs and thus the cost of poultry (Chander & Kannadhasan, 2021). Restaurant waste, as part of food waste, can have a high nutritional value for poultry and, on the other hand, can reduce environmental pollution, reduce the cost of waste disposal, as well as reduce competition between animals and humans in the use of valuable food resources (Seidavi et al., 2018).

Today, food waste is being used in various countries (Walker et al., 2002; Rabêlo et al., 2021). Recycling food waste and replacing it with poultry rations is one of the most important projects pursued by the South Korean government. Also, replacing food waste as feed-in poultry diets is known as a solution for food waste recycling in the United States (Sakaguchi et al., 2018). Japan

Received 13 Oct., 2021

Accepted 28 Dec., 2021

¹ Universitas Pasundan, Bandung, Jawa Barat, Indonesia

² Division of Nephrology & Hypertension, Department of Internal Medicine, Faculty of Medicine, Universitas Lambung Mangkurat / Ulin Hospital, Banjarmasin, Indonesia

³ Bahauddin Zakariya University, Multan, Pakistan

⁴ Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya, Indonesia

⁵ Al-Nisour University College, Baghdad, Iraq

⁶ Chitkara Business School, Chitkara University, Punjab, India

⁷ Scientific Research Center, Al-Ayen University, Thi-Qar, Iraq

⁸ Kuban State University Named After I. T. Trubilin, Krasnodar, Russia

⁹ Department of Health and Rehabilitation Sciences, College of Applied Medical Sciences, Prince Sattam bin Abdulaziz University, Al Kharj, Saudi Arabia

¹⁰ Department of Physical Therapy, Kasr Al-Aini Hospital, Cairo University, Giza, Egypt

¹¹ Health Department, Faculty of Health Science, University of Pembangunan Nasional Veteran Jakarta, Jakarta, Indonesia

*Corresponding author: ghulam.3yasin@gmail.com, rudiansyah@ulm.ac.id

also recycles a significant percentage of food waste produced for livestock and poultry feed (Westendorf et al., 1998). Recycling food waste and the production of functional products from it is very important because, in addition to reducing food imports, it can also help reduce environmental pollution (Cho et al., 2004).

It should be noted that the direct use of food waste in poultry feed without their processing can lead to the spread of pathogens and infections. Therefore, proper processing and disinfection of these materials is an important issue that requires great care (Ali et al., 2021; Arifjanov et al., 2021). Wet and unprocessed feed, due to its high moisture content, disrupts nutrition, reducing dry matter intake by reducing product shelf life. By studying various methods of conversion and processing of food waste, it is possible to effectively disinfect and clean food waste by performing heat treatment at a certain time and at the appropriate temperature. Although heat treatment causes some of the nutrients to be lost, the remaining nutrients can still meet the nutritional needs. In this paper, the nutritional value of food waste in comparison with the compounds used in the preparation of poultry feed has been investigated. Therefore, the purpose of this paper is to determine the nutritional value of restaurant waste and the possibility of replacement with poultry rations (Galal et al., 2020; Ferreira et al., 2021).

2 Material and methods

This section describes the method of sampling food waste and performing tests to measure the percentage of fat, dry matter, calcium, phosphorus, and protein.

2.1 Sampling method

In this paper, restaurant food waste has been used. Randomly sampled food waste was done for seven consecutive days, and the collected waste samples included red meat, fish, chicken, rice, and legumes. All experiments were performed in 5 replications.

2.2 Fat measurement

A Soxhlet extractor device and filter paper No. 42 were used to measure the amount of fat in the samples. First, the weight of the filter paper was measured. Three grams of samples were poured on paper, and after folding it, the samples were placed vertically inside the machine. 100 cc of hexane per sample was poured into the tank and stored at 70 °C for 4 hours to evaporate and condense the hexane, and their fat was completely washed away. After 4 hours, the samples were removed from the device and placed in a desiccator for 2 hours in order to completely evaporate the hexane. The samples were placed in an oven for 24 hours to dry. Finally, the total weight was measured, and the paper weight was subtracted from the sample weight, and the fat percentage was obtained from the following equation (Equation 1).

$$\text{fat percentage} = \frac{\text{Final weight} - \text{Initial weight}}{\text{Sample weight}} \times 100 \quad (1)$$




2.3 Dry matter measurement

The total amount of food without considering the percentage of moisture is called the percentage of dry matter. This section contains protein, fat, crude fiber, and minerals in the food. Moisture content is obtained by drying the food and calculating the weight loss. In this paper, in order to measure the amount of dry matter in food waste, food waste was first poured into a mixer and mixed at medium speed for 3 minutes. The mixed mixture was weighed and placed in glass containers in an oven at 70 °C for 48 hours. After 48 hours, remove the mixture from the oven and place it in the desiccator for 2 hours until it reaches ambient temperature. The dried sample was ground at medium speed for 5 minutes, and then the residual weight was measured (Equation 2).

$$\text{Percentage of dry matter} = \frac{\text{Dry sample weight}}{\text{Wet sample weight}} \times 100 \quad (2)$$

2.4 Measurement of protein

Protein is a major component in the structure of waste. It is found in the structure of muscles, skin, blood cells, hormones, as well as bone tissue (An et al., 2019). In this paper, the following method was used to measure the amount of protein in the waste. First, nitrogen, protein, and other acid-digesting compounds were converted to ammonium sulfate using concentrated sulfuric acid. After cooling, the acidic digestion product was diluted with distilled water, and then in the distillation step, ammonia was released by adding an alkali (sodium hydroxide) to the digested mixture. The released ammonia was adsorbed on boric acid and then released during the titration of ammonia bonded with boric acid (ammonium borate complex) using a standard acidic solution (sulfuric acid). The reactions that took place in each step are as follows (Equations 3-7).



Because the average amount of nitrogen in biological materials is about 16% based on dry matter, usually, to measure the crude protein content, the total nitrogen measured must be multiplied by a factor of 6.25. Finally, the percentage of protein was calculated using the following equation (Equation 8).

$$\text{Percentage of dry matter} = \frac{\text{The amount of protein in the sample}}{\text{sample weight}} \times 100 \quad (8)$$

2.5 Measurement of calcium

In this paper, in order to measure the percentage of calcium in food waste, the sample was first placed in an oven at 70 °C for 48 hours. After drying, 1 g of each sample was poured into a porcelain dish and burned in an electric oven at 550 °C, and

the sample ash was prepared. Calcium was measured by titration with potassium permanganate. To do this, 20 cc of ash solution was transferred to a beaker, and about 100 cc of distilled water and two drops of methyl reagent were added. Using pH meter and pH ammonium hydroxide solution, the solution was reduced to about seven and titrated with 0.1 potassium permanganate tetrazole solution until pink.

2.6 Phosphorus measurement

To measure the percentage of phosphorus in food waste, first, a dried sample of the product was prepared, and 1 g of it was burned in an electric oven at 550 °C, and the sample ash was prepared. A spectrophotometric method was used to measure phosphorus. For this purpose, 5 cc of the ash solution is transferred to a 100 cc volumetric flask, 25 mL of the vanadate-molybdate reagent is added, and the volume is increased. After complete mixing, leave for 10 minutes, then the adsorption intensity of the solution was measured at 420 nm against the empty solution. The concentration of phosphorus was obtained from the adsorption intensity, and the final result was calculated according to the amount of the sample solution.

3 Results and discussion

In this section, the values obtained for calcium (Ca), phosphorus (P), dry matter (DM), crude fat (CF), and crude protein (CP) are presented and compared with normal poultry ration values.

3.1 Examine the usual poultry rations

Poultry rations are mainly composed of corn, corn gluten, wheat, barley, wheat bran, soybean meal, and sunflower meal (Gou et al., 2018). Table 1 lists the amounts of calcium, phosphorus, dry matter, crude fat, and crude protein in each of these foods.

3.2 Investigation of measured parameters in restaurant food waste

The amounts of calcium, phosphorus, dry matter, crude fat, and crude protein of the food waste tested are presented in Table 2.

3.3 Comparison of calcium percentage

Calcium is essential for the formation of bones and eggshells. This mineral is formed in the bone as calcium phosphate and in eggshells as calcium carbonate. When egg production begins, the

calcium needed to form the eggshell comes from the calcium in the diet and the bones (Chen et al., 2015). In contrast, increasing calcium intake also reduces its absorption. Therefore, the amount of calcium consumed should be adjusted according to the table of nutritional needs. A comparison between the percentage of calcium in restaurant food waste and the percentage of calcium in products used in the preparation of poultry rations is shown in Figure 1. According to the diagram, it can be seen that the percentage of calcium in all five samples tested is almost the same, but the percentage of calcium in the products used in poultry diets is different, and this difference depends on the type of application. Considering the average percentage of calcium in food waste and comparing it with other items in the poultry diet, it can be expected that replacing food waste with soybean meal will have positive effects on growth and feed consumption. However, since excessive calcium intake in poultry has a negative effect and causes calcium excretion, complete replacement of food waste with poultry rations requires further research and study on live poultry.

3.4 Comparison of phosphorus percentage

A comparison of the percentage of phosphorus in food waste and products used in the preparation of poultry feed is given in Figure 2. According to the diagram, it can be seen that the percentage of phosphorus in food waste is very low, and these materials could not be replaced with poultry diets to meet the needs of poultry phosphorus.

3.5 Comparison of dry matter percentage

A comparison of the percentage of dry matter in food waste with the materials used in the preparation of poultry rations is given in Figure 3. According to the chart, it can be seen that the percentage of dry matter in food waste is about one-third of the dry matter of diets. This is obvious considering the nature of food waste because this waste has very high humidity. As a result, from a dry matter point of view, waste could not be used in poultry feed. However, due to the fact that no other by-products were used in the processing of waste in this paper, and since bread waste is produced in abundance in restaurants and this waste has a high percentage of dry matter, it is possible to use dry bread waste, improves dry matter content in combination with food waste.

3.6 Comparison of protein percentages

A comparison of the percentage of protein in food waste with the products used in the preparation of poultry feed is given in Figure 4. According to the chart, the percentage of protein in


Table 1. Values of basic parameters in poultry diets.

Poultry rations	Ca%	P%	DM%	CF%	CP%
Corn	0.02	0.3	88	4.3	6
corn gluten	0.11	0.84	90	3.2	22
barley	0.06	0.38	89	2.1	11.3
wheat	0.05	0.43	89	2.3	14
wheat bran	0.14	1.2	89	4.6	12.5
soybean meal	0.29	0.66	88	18.8	41
sunflower meal	0.4	1.03	91	2.4	31


Table 2. Measured parameter values in restaurant food waste.

Food waste	Ca%	P%	DM%	CF%	CP%
Red meat and rice	0.31	0.25	27.82	12.38	21.91
Fish and rice	0.28	0.25	29.12	10.1	17.95
Chicken and rice	0.24	0.15	31.58	12.25	30.42
Cereals and rice	0.33	0.11	29.27	8.2	33.27
Fried chicken	0.28	0.29	27.23	20.7	25.25


Using restaurant food waste as poultry rations



6
Figure 1. Comparison of the percentage of calcium in restaurant food waste with poultry rations.



6
Figure 2. Comparison of the percentage of phosphorus in restaurant food waste with poultry rations.



6
Figure 3. Comparison of dry matter percentage of restaurant food waste with poultry rations.




Figure 4. Comparison of the protein content of restaurant food waste with poultry rations.




Figure 5. Comparison of the fat content of restaurant food waste with poultry ration.

food waste is higher than all cereals and is only slightly different from the percentage of protein in soybeans and sunflower meals. Due to the structure of soybean meal and sunflower meal as processed food that is used as a source of protein and by comparing the means of the two groups, it can be said that the percentage of protein in food waste can be replaced with protein in poultry diets.

3.7 Comparison of fat percentage

A comparison of the percentage of fat in food waste with the percentage of fat in products used in poultry diets is shown in Figure 5. According to the chart, it can be seen that the average percentage of fat in food waste is higher than the average fat in poultry rations. The percentage of fat in soybean meal and its importance in the growth and weight of poultry, and based on the fact that the presence of fat in poultry diets is important

and necessary, it can be expected that the replacement of food waste with poultry rations along with other nutrients, need to have positive effects on poultry weight and growth. In general, experiments on live poultry can be more useful in order to examine the subject more closely and achieve more credible results.

4 Conclusion

The following can be deduced from the above points about restaurant food waste and studies on dry matter, protein, calcium, phosphorus, and fat parameters. The average dry matter of restaurant food waste is about 60% less than the average poultry ration, which indicates that in terms of dry matter, food waste is not a suitable substitute for poultry ration. The average calcium of restaurant food waste is about 0.14% higher than the average diet of poultry. The average calcium in food waste is close to the amount of calcium in soybean meal, which can be replaced with this part of the poultry

diet. The average phosphorus of restaurant food waste is about 0.48% less than the average poultry diet, indicating that food waste is not a suitable substitute for poultry rations in terms of phosphorus. The average crude protein of restaurant food waste is about 5.65% higher than the average poultry ratio. In this case, food waste can be used as a substitute for poultry rations. The average crude fat of restaurant food waste is about 7.34% higher than the average poultry ratio. In this case, food waste can be used as a substitute for poultry rations. In this paper, the numerical values of the basic parameters between poultry rations and restaurant food waste have been compared, but for a definite replacement with poultry rations, several experiments should be performed on live poultry.

References

- Ali, S., Hassan, M., Essam, T., Ibrahim, M. A., & Al-Amry, K. (2021). Biodegradation of aflatoxin by bacterial species isolated from poultry farms. *Toxicon*, 195, 7-16. <http://dx.doi.org/10.1016/j.toxicon.2021.02.005>. PMID:33610638.
- An, B.-K., Choo, W.-D., Kang, C.-W., Lee, J., & Lee, K.-W. (2019). Effects of dietary lycopene or tomato paste on laying performance and serum lipids in laying hens and on malondialdehyde content in egg yolk upon storage. *The Journal of Poultry Science*, 56(1), 52-57. <http://dx.doi.org/10.2141/jpsa.0170118>. PMID:32055196.
- Arifjanov, A. M., Akmalov, S. B., & Samiev, L. N. (2021). Extraction of urban construction development with using Landsat satellite images and geoinformation systems. *Journal of Water and Land Development*, 48, 65-69.
- Barreira, T. F., Paula, G. X. Fo., Priore, S. E., Santos, R. H. S., & Pinheiro-Sant'ana, H. M. (2021). Nutrient content in ora-pro-nóbis (*Pereskia aculeata* Mill.): unconventional vegetable of the Brazilian Atlantic Forest. *Food Science and Technology*, 41(Suppl. 1), 47-51. <http://dx.doi.org/10.1590/fst.07920>.
- Burzyńska, I. (2019). Monitoring of selected fertilizer nutrients in surface waters and soils of agricultural land in the river valley in Central Poland. *Journal of Water and Land Development*, 43(1), 41-48. <http://dx.doi.org/10.2478/jwld-2019-0061>.
- Chander, M., & Kannadhasan, M. S. (2021). Landless animal and poultry production prospects: an overview on feeding and sustainability with special reference to fruit and vegetable wastes (FVWs). *Organic Agriculture*, 11(2), 285-300. <http://dx.doi.org/10.1007/s13165-020-00292-5>.
- Chen, C., Chaudhary, A., & Mathys, A. (2020). Nutritional and environmental losses embedded in global food waste. *Resources, Conservation and Recycling*, 160, 104912. <http://dx.doi.org/10.1016/j.resconrec.2020.104912>.
- Chen, T., Jin, Y., & Shen, D. (2015). A safety analysis of food waste-derived animal feeds from three typical conversion techniques in China. *Waste Management*, 45, 42-50. <http://dx.doi.org/10.1016/j.wasman.2015.06.041>. PMID:26188613.
- Chen, T., Jin, Y., Liu, F., Meng, X., Li, H., & Nie, Y. (2012). Effect of hydrothermal treatment on the levels of selected indigenous microbes in food waste. *Journal of Environmental Management*, 106, 17-21. <http://dx.doi.org/10.1016/j.jenvman.2012.03.045>. PMID:22562007.
- Cho, Y. M., Lee, G. W., Jang, J. S., Shin, I. S., Myung, K. H., Choi, K. S., Bae, I. H., & Yang, C. J. (2004). Effects of feeding dried leftover food on growth and body composition of broiler chicks. *Asian-Australasian Journal of Animal Sciences*, 17(3), 386-393. <http://dx.doi.org/10.5713/ajas.2004.386>.
- Dou, Z., Toth, J. D., & Westendorf, M. L. (2018). Food waste for livestock feeding: feasibility, safety, and sustainability implications. *Global Food Security*, 17, 154-161. <http://dx.doi.org/10.1016/j.gfs.2017.12.003>.
- Ferreira, L. S., Brito-Oliveira, T. C., & Pinho, S. C. (2021). Brazil nut (*Bertholletia excelsa*) oil emulsions stabilized with thermally treated soy protein isolate for vitamin D₃ encapsulation. *Food Science and Technology*, 1-11. Ahead of print. <http://dx.doi.org/10.1590/fst.17521>.
- Galal, H. M. F., Abbas, H. S., & Abdou, A. A. (2020). Using of dry tomato vines hay in growing lambs ration. *Journal of Animal and Poultry Production*, 11(12), 525-532. <http://dx.doi.org/10.21608/jappmu.2020.161174>.
- Galhardo, D., Garcia, R. C., Schneider, C. R., Braga, G. C., Chambó, E. D., França, D. L. B., & Ströher, S. M. (2021). Physicochemical, bioactive properties and antioxidant of *Apis mellifera* L. honey from western Paraná, southern Brazil. *Food Science and Technology*, 41(Suppl. 1), 247-253. <http://dx.doi.org/10.1590/fst.11720>.
- Kim, S., Lee, Y., Lin, K.-Y. A., Hong, H., Kwon, E. E., & Lee, J. (2020). The valorization of food waste via pyrolysis. *Journal of Cleaner Production*, 259, 120816. <http://dx.doi.org/10.1016/j.jclepro.2020.120816>.
- Molajou, A., Pouladi, P., & Afshar, A. (2021). Incorporating social system into water-food-energy nexus. *Water Resources Management*, 35(13), 4561-4580. <https://doi.org/10.1007/s11269-021-02967-4>.
- Ojha, S., Bußler, S., & Schlüter, O. K. (2020). Food waste valorisation and circular economy concepts in insect production and processing. *Waste Management*, 118, 600-609. <http://dx.doi.org/10.1016/j.wasman.2020.09.010>. PMID:33010691.
- Olszewska, M. A., Gędas, A., & Simões, M. (2020). Antimicrobial polyphenol-rich extracts: applications and limitations in the food industry. *Food Research International*, 134, 109214. <http://dx.doi.org/10.1016/j.foodres.2020.109214>. PMID:32517896.
- Rabêlo, C. A., Ricardo, M., Porfírio, J. A., Pimentel, T. C., Nascimento, J. S., & Costa, L. E. O. (2021). Psychrotrophic bacteria in Brazilian organic dairy products: identification, production of deteriorating enzymes and biofilm formation. *Food Science and Technology*, 41(3), 799-806. <http://dx.doi.org/10.1590/fst.68420>.
- Ramos, G. L. P. A., & Nascimento, J. S. (2020). *Pseudomonas* SP. in uninspected raw goat's milk in Rio de Janeiro, Brazil. *Food Science and Technology*, 40(Suppl. 2), 605-611. <http://dx.doi.org/10.1590/fst.21719>.
- Sakaguchi, L., Pak, N., & Potts, M. D. (2018). Tackling the issue of food waste in restaurants: options for measurement method, reduction and behavioral change. *Journal of Cleaner Production*, 180, 430-436. <http://dx.doi.org/10.1016/j.jclepro.2017.12.136>.
- Seidavi, A. R., Azizi, M., Rangi, M., Laudadio, V., & Tufarelli, V. (2018). Practical applications of agricultural wastes in poultry feeding in Mediterranean and Middle East regions. Part 2: tomato, olive, date, sunflower wastes. *World's Poultry Science Journal*, 74(3), 443-452. <http://dx.doi.org/10.1017/S004393391800051X>.
- Seidavi, A., Zaker-Esteghamati, H., & Salem, A. Z. M. (2020). A review on practical applications of *Citrus sinensis* by-products and waste in poultry feeding. *Agroforestry Systems*, 94(4), 1581-1589. <http://dx.doi.org/10.1007/s10457-018-0319-2>.
- Sharma, P., Gaur, V. K., Kim, S.-H., & Pandey, A. (2020). Microbial strategies for bio-transforming food waste into resources. *Bioresource Technology*, 299, 122580. <http://dx.doi.org/10.1016/j.biortech.2019.122580>. PMID:31877479.
- Walker, P. M., Finnigan, D. M., Brown, S. A., & Dust, J. M. (2002). Evaluation of feed mixtures amended with processed food waste as feedstuffs for finishing lambs. *The Professional Animal Scientist*, 18(3), 237-246. [http://dx.doi.org/10.15232/S1080-7446\(15\)31527-8](http://dx.doi.org/10.15232/S1080-7446(15)31527-8).
- Westendorf, M. L., Dong, Z. C., & Schoknecht, P. A. (1998). Recycled cafeteria food waste as a feed for swine: nutrient content digestibility, growth, and meat quality. *Journal of Animal Science*, 76(12), 2976-2983. <http://dx.doi.org/10.2527/1998.76122976x>. PMID:9928600.

Investigation of parameters in restaurant food waste for use as poultry rations

ORIGINALITY REPORT

8%

SIMILARITY INDEX

6%

INTERNET SOURCES

4%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	www.karger.com Internet Source	1%
2	repository.embuni.ac.ke Internet Source	1%
3	Letícia DOS SANTOS FERREIRA, Thais CARVALHO BRITO-OLIVEIRA, Samantha Cristina DE PINHO. "Brazil nut (<i>Bertholletia excelsa</i>) oil emulsions stabilized with thermally treated soy protein isolate for vitamin D3 encapsulation", Food Science and Technology, 2022 Publication	1%
4	old.scielo.br Internet Source	1%
5	randd.defra.gov.uk Internet Source	1%
6	Rola Khishfe, Norman Lederman. "Relationship between Instructional Context	1%

and Views of Nature of Science", International
Journal of Science Education, 2007

Publication

7	cathepsink.com Internet Source	1 %
8	www.tandfonline.com Internet Source	1 %
9	Alexandros Georganas, Elisavet Giamouri, Athanasios C. Pappas, George Papadomichelakis et al. "Bioactive Compounds in Food Waste: A Review on the Transformation of Food Waste to Animal Feed", Foods, 2020 Publication	<1 %
10	www.readkong.com Internet Source	<1 %

Exclude quotes On

Exclude matches < 15 words

Exclude bibliography On