

REKAM JEJAK DIGITAL ARTIKEL

**FUNCTIONAL OUTCOME AND SURVIVAL RATE OF METASTATIC
BONE DISEASE ON FEMUR AND PELVIC BONE AFTER SURGICAL
TREATMENT**

HIV Nursing 2022

STATUS JURNAL SAAT INI

Scopus Preview

Author Search Sources [Create account](#) [Sign in](#)

Source details

Feedback > Compare sources >

HIV Nursing
Scopus coverage years: from 2006 to 2016, from 2018 to Present
Publisher: Mediscript Ltd.
ISSN: 1474-7359
Subject area: [Nursing: Advanced and Specialized Nursing](#)
Source type: Journal

[View all documents >](#) [Set document alert](#) [Save to source list](#)

CiteScore 2021	0.4
SJR 2021	0.110
SNIP 2021	0.036

CiteScore CiteScore rank & trend Scopus content coverage

Improved CiteScore methodology

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021	CiteScoreTracker 2022
0.4 = $\frac{18 \text{ Citations 2018 - 2021}}{47 \text{ Documents 2018 - 2021}}$	0.1 = $\frac{39 \text{ Citations to date}}{711 \text{ Documents to date}}$
<small>Calculated on 05 May, 2022</small>	<small>Last updated on 05 February, 2023 - Updated monthly</small>

HIV Nursing- Impact Score, Overall Ranking, h-index, SJR, Rating, Publisher, ISSN, and Other Important Metrics


Last Updated on May 27, 2022

Impact Score  0.07	h-Index  5	Rank  24414	SJR  0.11
--	--	---	---

UPDATE IMPACT FACTOR JOURNAL

Impact Score Trend

Year wise Impact Score (IS) of HIV Nursing. Based on Scopus data.




Year	Impact Score (IS)
2022/2023	Coming Soon
2021	0.07
2020	0.29
2019	0.10
2018	0.04


SCIMAGO RANK JURNAL

HIV Nursing

<p>COUNTRY</p> <p>United Kingdom</p> <ul style="list-style-type: none"> Universities and research institutions in United Kingdom Media Ranking in United Kingdom 	<p>SUBJECT AREA AND CATEGORY</p> <p>Nursing</p> <ul style="list-style-type: none"> Advanced and Specialized Nursing 	<p>PUBLISHER</p> <p>Mediscrypt Ltd</p>	<p>H-INDEX</p> <p>5</p>
<p>PUBLICATION TYPE</p> <p>Journals</p>	<p>ISSN</p> <p>14747359</p>	<p>COVERAGE</p> <p>2006-2008, 2011, 2015-2016, 2018-2021</p>	<p>INFORMATION</p> <p>Homepage</p> <p>How to publish in this journal</p> <p>editorial@HIVNursing.net</p>



SUBMISSION (14 OKTOBER 2022) & ACCEPTANCE ARTIKEL (16 NOVEMBER 2022)



Functional Outcome and Survival rate of Metastatic Bone Disease on femur And Pelvic Bone after Surgical Treatment

Husna Dharma Putera¹, Ahmad Fauzi Kamal², Yogi Prabowo²

¹Department of Orthopaedic and Traumatology, Faculty of Medicine, Universitas Lambung Mangkurat/Ulin General Hospital, Banjarmasin

²Department of Orthopaedic and Traumatology, Faculty of Medicine, Universitas Indonesia/Cipto Mangunkusumo Hospital, Jakarta

Corresponding E-mail: hdputera@ulm.ac.id

Abstract

Introduction. Most groups of cancers that often metastasize to bone invariably induce pathological fractures. Between 9-29% of patients with metastases will have a pathological fracture and 90% of fractures require surgery. Fractures on pathologic bone have a major impact on life quality. **Methods.** Between 2013 - 2017, 30 patients with metastatic tumors on the femur and pelvic were treated at Cipto Mangunkusumo Hospital. Basic patients' medical records were analyzed, with particular focus on the cancer type, age, gender, tumor site, mechanism of fracture, treatment type, survival rate (Kaplan Meier test), and functional outcome (MSTS score). **Results.** Eleven patients were male and nineteen patients were female. The mean age was 56 + 7 years. The MSTS score was excellent in 10 (33.3%), moderate in 4 (13.3%), and poor in 4 (13.3%). The tumor located in the shaft femur and treated by ORIF had the highest mean MSTS score. Slipped was the most common mechanism of injury due to pathological fracture. The mean survival time of metastatic bone from the four most common tumor origin were 6 months (lung), 4.33 months (multiple myeloma) 18.75 months (thyroid), and 20.33 months (breast). The data were then analyzed using the Kaplan-Meier method and Log Rank (Mantel-Cox) test. From this data, the equality of survival distribution using the Log Rank (Mantel-Cox) test was calculated, and found that the P value was 0.003. It means that there is a significant difference between survival rate and tumor origin. **Conclusions.** Pathologic fractures in patients with metastatic cancer needed a combination of surgical and oncological treatment. Patients were treated surgically, and most of them have an excellent functional outcome. The survival rate varied, based on cancer type.

Keywords: Metastatic bone disease, femur & pelvic bone, functional outcome, survival

Introduction

Bone is the most frequent site of metastatic disease. The sites most commonly affected are the spine, the pelvic bones, the proximal part of the femur, and the humerus.(1-3) Approximately, 70% of patients who die of breast or prostate cancer have also had bone metastases. Thyroid, lung, and kidney cancers also have a propensity to involve the skeleton.(1) Carcinoma penis spread to bones is uncommon.(4) Most cancer groups that often metastasize to bone such as breast, lung, and kidney cancer invariably induce osteolytic lesions.(5) Approximately, 30-65% of patients with metastatic lung cancers will develop bone metastases, as will approximately 47% of patients with advanced thyroid cancer and 30% of patients with advanced renal carcinoma. 9-29% of patients with metastases will have a pathological fracture and 90% of fractures require surgery. Fractures on pathologic bone have major impact on life quality.(1, 6, 7)

Pathological fracture of the proximal femur is the main cause of loss the mobile ability in cancer survivors.(8) Metastatic lesions located in the proximal femur are particularly frequent. About 10% of patients with

primary malignant tumors will develop metastasis of the proximal femur. Among femur metastatic tumors, 50% of the lesions occur in the femoral neck, 30% occur in the sub- trochanteric site, and 20% occur in the intertrochanteric site. This is related to the well-developed vascular system in the intertrochanteric area. Most frequently bone metastases are derived from breast, kidney, thyroid, and prostate cancer, or myeloma. Besides prostate cancer, most metastases are lytic or mixed, and thus patients are at a high risk of pathological fractures.(9-11)

Massive metastases to the pelvis, especially to the periacetabular area, are still a difficult treatment problem. They inhibit patients' walking independently, thus forcing the necessity to use crutches or a walking frame.(2) Ideally, operative treatment should allow immediate weight bearing with the least possible morbidity. The goals of surgery are to achieve local tumor control and structural stability to restore function as soon as possible.(12)

This research aims to describe metastatic bone disease on the femur and pelvic bone, including cancer type, age, gender, tumor site, mechanism induced fracture, treatment type, survival rate, and

functional outcome.


Methods

Between 2013 - 2017, 30 patients with a metastatic tumor on the femur and pelvic were treated at Cipto Mangunkusomo Hospital. Basic patients' medical records were analyzed, with particular focus on the cancer type, age, gender, tumor site, mechanism induced fracture, treatment type, survival rate, and functional outcome. To determine patients' functional outcomes, MSTS scores were used. We used The Kaplan Meier test method to determine the survival rate.

Result

Based on demographic characteristics, eleven patients were male and nineteen patients were female (Fig 1). The mean age was 56 + 7 years (Fig 2). The mean MSTS score was 24.11 + 5.54. The MSTS was further classified into several groups. Excellent was defined as 75-100% of the total score, good as 70-74% of the total score, moderate as 60-69%, fair as 50-59%, and poor as <50%. The MSTS score was excellent in 10 (33.3%), moderate in 4 (13.3%), and poor in 4 (13.3%).

Fig. 1. Gender




The MSTS score was analyzed between the male and female groups. 20 patients underwent surgical treatment, but two patients had been drop out due to death after surgery. The mean MSTS score in the male group was 25.28 whereas in the female group was 23.36. The mean difference between the two groups was 1.92, 95% CI (-3.84-7.68). There was no significant difference in MSTS scores in male and female groups (p=0.49) [Tabel 1].

Table 1. Comparison of MSTS Score Between Gender Group

Gender	Mean MSTS Score	p	95% CI
Male	25.28 + 4.95	0.49	1.92 (-3.84 – 7.68)
Female	23.36 + 5.98		


The distribution of the patient was based on the group of age as below (Fig. 2). The age group between 51 – 60 years old was most frequent.

Fig. 2. Group of Age



The MSTS score was analyzed based on age group. There was no statistical difference in MSTS scores between different age groups (p=0.25). The site of bone that metastasized by cancer is as below (Fig. 3). Most metastatic bone disease was spread to the proximal femur bone.

Fig. 3. Site of bone



The tumor site of bone metastatic disease is pelvic, proximal femur, distal femur, and shaft femur. The tumor located in the shaft femur had the highest mean MSTS score (Tabel 2). However, there was no statistical difference in MSTS scores between tumor site groups (p=0.75).

Table 2. Comparison of MSTS Score Based on Tumor Site

Tumor Site	Mean Score	p
Pelvic	21.00	0.75
Proximal femur	23.54 + 5.73	
Distal femur	25.00 + 6.16	
Shaft femur	29.00	

The distribution of patients is based on the origin of cancer as below (Fig. 4). The most tumor origin was from lung cancer. The MSTS score was analyzed based on tumor origin (Tabel 3). There was no statistical difference in MSTS score between the tumor origin group (p=0.25).

Fig. 4. Type of Cancer





Table 3. Comparison of MSTS Score based on tumor origin

Tumor Origin	Mean MSTS Score	p
Lung	23.33	0.25
Thyroid	28.00	
Breast	24.33	
Multiple Myeloma	23.00	
Prostate	21.00	
Hepatoma	27.00	
Colon	29.00	
Melanoma	29.00	
Cervix	17.00	

The distribution of patients based on mechanism


injury due to fracture as below (Fig. 5). Slipped was the most common mechanism of injury due to pathological fracture.

Fig. 5. Mechanism of Injury



The distribution of patients based on treatment is as below (Fig. 6). Most of the patients were treated by hemiarthroplasty.

Fig 6. Surgical Treatment



The surgical treatment of femur and pelvic metastatic bone disease is hemiarthroplasty, arthrodesis, or open reduction internal fixation (ORIF). The group treated with ORIF had the highest mean MSTS score. However, based on statistical analysis, there was no difference in MSTS score between surgical treatment groups ($p=0.66$) [Tabel 4].

Table 4. Comparison of MSTS Score Based on Surgical Treatment

Treatment	Mean Score	p
Hemiarthroplasty	23.54 ± 5.73	0.66
Arthrodesis	23.00	
ORIF	25.60 ± 2.3	

This study used The Kaplan Meier test method to determine the survival rate. The censored patient could be defined as the patient who underwent loss of follow-up or was still alive during the collecting data process. Time was in months, while the cumulative survival of 1 means alive and 0 means event of death. Based on the result, the mean survival rate of Femur and Pelvic Metastatic Bone Disease was 11.23 months.

Most femur and pelvic bone metastatic are derived from lung, breast, thyroid, prostate, liver, or myeloma. The survival data could be seen in the following tables 4 & 5:

Bone Disease Based on Tumor Origin

Table 4. Summary of Remission Status of Femur & Pelvic Metastatic

Tumor Origin	Total N	N of Events	Censored	
			N	Percent
Lung	9	9	0	0%
Thyroid	4	4	0	0%
Breast	3	2	0	0%
Multiple Myeloma	3	3	0	0%
Prostate	2	2	0	0%
Hepar	2	2	0	0%
Colon	1	1	0	0%
Ovarium	1	1	0	0%
Cervix	1	1	0	0%
Overall	26	26	0	0%

Table 5. Mean of survival time of Pelvic Metastatic Bone Disease Based on Tumor Origin

	Mean		95% Confidence Interval
	Estimate	Standard error	
Lung	6.00	0.60	2.68 – 20.32
Thyroid	18.75	5.04	4.82 – 7.17
Breast	20.33	2.02	8.87 – 28.62
Multiple Myeloma	4.33	1.66	0 – 23.24
Prostate	11.50	4.50	1.06 – 7.60
Hepar	10.50	6.50	26.00 – 26.00
Colon	26.00	0.00	16.35 – 24.30
Ovarium	14.00	0.00	14.00 – 14.00
Cervix	5.00	0.00	5.00 – 5.00

The four most common tumor origins of the femur and pelvic metastatic bone disease were lung, multiple myeloma, thyroid, and breast. The mean survival time for each group based on tumor origin was computed. The mean survival time of metastatic bone from the four most common tumor origin were 6 months (lung), 4.33 months (multiple myeloma) 18.75 months (thyroid), and 20.33 months (breast). The data were then analyzed with the Kaplan-Meier method and Log Rank (Mantel-Cox) test. From this data, the equality of survival distribution using the Log Rank (Mantel-Cox) test was calculated, and found that the P value was 0.003. It means that there is a significant difference between survival rate and

tumor origin.

Discussion

Pathologic fractures in patients with metastatic cancer of the breast, prostate, lung, multiple myeloma, bladder, thyroid, kidney, and other primary carcinomas with skeletal involvement are a common problem in clinical orthopedic. Metastatic lesions can be lytic, blastic, or a mixed type, whereas the majority of all metastatic lesions are lytic and these lesions have the highest impact on bone strength, which causes pathological fractures. Pathological fractures of the femur mostly occur during daily activities, such as starting to walk,

standing, raising from a chair or bed, or stair climbing.^(13, 14) In this research, slipping while walking was the most mechanism-induced pathological fracture.

Most patients with bone metastases need a combination of surgical and oncological treatment. The disease is associated with the general bad condition, pain, reduced mobility, walking and working disability, and problems with independent functioning.^(9, 15) Orthopedic treatment for the metastatic bone disease of the extremities may be prophylactic to prevent a fracture or to stabilize a fracture that has occurred. Surgical resection of primary tumors or metastatic lesions of the proximal femur can result in large bone and soft tissue resections. Surgical treatment of pelvic bone tumors continues to pose a challenge to the orthopedic oncology community. Traditionally, pelvic tumors were resected through hindquarter amputation. The surgical procedure choice is associated with detrimental cosmetic, physical, and psychological outcomes. At present, the majority of patients can be treated with limb-salvaging internal hemipelvectomy.⁽¹⁶⁻¹⁹⁾

Stabilization of a fracture or prevention of an impending fracture usually requires internal fixation devices such as plates and screws or intramedullary rods. Sometimes, partial or complete bone replacement is required to help a patient return to their pre-morbid functional state. Orthopedic treatment for metastatic bone disease aims to alleviate pain and increase mobility and functional independence in a metastatic condition.^(16-18, 20, 21)

In this research, patients were treated surgically, and most of them have an excellent functional outcomes. But, a few of them have a poor functional outcome, which may be due to bad performance before undergoing surgery or progression of cancer disease. The research by Guzik et al showed surgical treatment for proximal femur bone metastases is particularly good in patients after standard or modular endoprostheses replacement.^(9, 22)

Over the past 20 years, the combination of improved systemic therapies, high-resolution cross-sectional imaging modalities, and evolving technological advances in orthopedic implant design has afforded greater opportunity for limb salvage after the oncological resection of malignant bone neoplasms.⁽²³⁾ Cancer mortalities are determined by individual characteristics and behaviors. Several cancer incidences and mortality studies in European countries have implied that there is wide variation among geographic areas because of various exposures to risk factors such as air pollution, occupational exposures, education, and differences in lifestyle.⁽²⁴⁾ In this research, the mean survival time of metastatic bone from the four most common tumor origin were 6 months (lung), 4.33 months (multiple myeloma), 18.75 months (thyroid), and 20.33 months (breast). Lung cancer research in Korea showed the median survival time at which half the

patients were expected to be alive was 12.4 months with a 95% confidence interval (CI) (11.3–13.8 months).⁽²⁴⁾ Multicenter study in German, collecting data on thyroid cancer between 2000–2015, stage-dependent OS at 6 months was 78, 54%, and 18% for stage IVA, B, and C, respectively, 29% of patients survived >1 year.⁽²⁵⁾ Study about metastatic bone breast cancer in Japan showed that the median OS and 5-year survival rates were 60.0 months and 50.0% (95% CI: 48.8–71.3 months), respectively. In Denmark, the median OS and 5-year survival rates were 17 months and 23%, respectively.⁽²⁶⁾

Conclusion

Pathologic fractures in patients with metastatic cancer need a combination of surgical and oncological treatment. Orthopedic treatment for metastatic bone disease aims to alleviate pain and increase mobility and functional independence in a metastatic condition. Patients were treated surgically, and most of them have excellent functional outcomes. The survival rate varied, depending on the cancer type.

References

1. Agarwal MG, Nayak P. Management of skeletal metastases: An orthopaedic surgeon's guide. *Indian J Orthop.* 2015;49(1):83-100.
2. Guzik G. Treatment of metastatic lesions localized in the acetabulum. *J Orthop Surg Res.* 2016;11(1):54.
3. Yosibash Z, Plitman Mayo R, Dahan G, Trabelsi N, Amir G, Milgrom C. Predicting the stiffness and strength of human femurs with real metastatic tumors. *Bone.* 2014;69:180-90.
4. Hussain S, Solanki FS, Sharma DB, Sharma D. Pathological Fracture of the Femur by Metastatic Carcinoma Penis-a Rare Presentation. *Indian J Surg.* 2016;78(2):149-50.
5. Hirata T, Park SC, Muldong MT, Wu CN, Yamaguchi T, Strasner A, et al. Specific bone region localization of osteolytic versus osteoblastic lesions in a patient-derived xenograft model of bone metastatic prostate cancer. *Asian J Urol.* 2016;3(4):229-39.
6. Anract P, Biau D, Boudou-Rouquette P. Metastatic fractures of long limb bones. *Orthop Traumatol Surg Res.* 2017;103(1S):S41-S51.
7. Derikx LC, Verdonschot N, Tanck E. Towards clinical application of biomechanical of tools for the prediction of fracture risk in metastatic bone disease. *Journal of Biomechanics.* 2015;48:6.
8. Araki N, Chuman H, Matsunobu T, Tanaka K, Katagiri H, Kunisada T, et al. Factors associated with the decision of operative procedure for proximal femoral bone metastasis: Questionnaire survey to institutions participating the Bone and Soft Tissue Tumor Study Group of the Japan Clinical Oncology Group. *J Orthop Sci.* 2017;22(5):938-45.
9. Guzik G. Oncological and functional results after surgical treatment of bone metastases at the

- proximal femur. *BMC Surg.* 2018;18(1):5.
10. Chau S, Chandra M, Grimsrud CD, Gonzalez JR, Hui RL, Lo JC. Femur fracture classification in women with a history of breast cancer. *J Bone Oncol.* 2014;3(2):49-53.
 11. Feng H, Wang J, Xu J, Chen W, Zhang Y. The surgical management and treatment of metastatic lesions in the proximal femur: A mini review. *Medicine (Baltimore).* 2016;95(28):e3892.
 12. Malviya A, Gerrand C. Evidence for orthopaedic surgery in the treatment of metastatic bone disease of the extremities: a review article. *Palliat Med.* 2012;26(6):788-96.
 13. Benca E, Patsch JM, Mayr W, Pahr DH, Windhager R. The insufficiencies of risk analysis of impending pathological fractures in patients with femoral metastases: A literature review. *Bone Rep.* 2016;5:51-6.
 14. Zaporowska-Stachowiak I, Luczak J, Hoffmann K, Stachowiak K, Bryl W, Sopata M. Managing metastatic bone pain: New perspectives, different solutions. *Biomed Pharmacother.* 2017;93:1277-84.
 15. Guzik G. Results of the treatment of bone metastases with modular prosthetic replacement--analysis of 67 patients. *J Orthop Surg Res.* 2016;11:20.
 16. Benedetti MG, Bonatti E, Malfitano C, Donati D. Comparison of allograft-prosthetic composite reconstruction and modular prosthetic replacement in proximal femur bone tumors: functional assessment by gait analysis in 20 patients. *Acta Orthop.* 2013;84(2):218-23.
 17. Puchner SE, Funovics PT, Hipfl C, Dominkus M, Windhager R, Hofstaetter JG. Incidence and management of hip dislocation in tumour patients with a modular prosthesis of the proximal femur. *Int Orthop.* 2014;38(8):1677-84.
 18. Bus MP, Szafranski A, Sellevold S, Goryn T, Jutte PC, Bramer JA, et al. LUMiC((R)) Endoprosthetic Reconstruction After Periacetabular Tumor Resection: Short-term Results. *Clin Orthop Relat Res.* 2017;475(3):686-95.
 19. Benevenia JK, R. ;Patterson, F. ;Beebe, K. ;Wirtz, D. C. ;Rivero, S. ;Palma, M. ;Friedrich, M. J. . Outcomes of a Modular Intercalary Endoprosthesis as Treatment for Segmental Defects of the Femur, Tibia, and Humerus. *Clin Orthop Relat Res.* 2016;474(2):539-48.
 20. Kirkinis MN, Lyne CJ, Wilson MD, Choong PF. Metastatic bone disease: A review of survival, prognostic factors and outcomes following surgical treatment of the appendicular skeleton. *Eur J Surg Oncol.* 2016;42(12):1787-97.
 21. Mavrovi E, Pialat JB, Beji H, Kalenderian AC, Vaz G, Richioud B. Percutaneous osteosynthesis and cementoplasty for stabilization of malignant pathologic fractures of the proximal femur. *Diagn Interv Imaging.* 2017;98(6):483-9.
 22. Pitera T, Guzik G, Biega P. Assessment of Post-operative Physical Performance in Patients after Resection Arthroplasty of the Proximal Femur. *Ortop Traumatol Rehabil.* 2017;19(4):333-40.
 23. Monument MJB, N. M. ;Bowles, A. J. ;Jones, K. B. ;Randall, R. L. . What are the 5-year survivorship outcomes of compressive endoprosthetic osseointegration fixation of the femur? *Clin Orthop Relat Res.* 2015;473(3):883-90.
 24. Kwak M, Kim C. Disparities by Age, Sex, Tumor Stage, Diagnosis Path, and Area-level Socioeconomic Status in Survival Time for Major Cancers: Results from the Busan Cancer Registry. *J Korean Med Sci.* 2017;32(12):1974-83.
 25. Wendler J, Kroiss M, Gast K, Kreissl MC, Allelein S, Lichtenauer U, et al. Clinical presentation, treatment and outcome of anaplastic thyroid carcinoma: results of a multicenter study in Germany. *Eur J Endocrinol.* 2016;175(6):521-9.
 26. Kuranishi F, Imaoka Y, Sumi Y, Uemae Y, Yasuda-Kurihara H, Ishihara T, et al. Rate of Clinical Complete Response for 1 Year or More in Bone-Metastatic Breast Cancer after Comprehensive Treatments including Autologous Formalin-Fixed Tumor Vaccine. *Int J Breast Cancer.* 2018;2018:4879406.