

Vol. 4 No. 1. Maret 2019

# DENTINO

Jurnal Kedokteran Gigi

Terbit Setiap Maret dan September

TERAKREDITASI : No. 21/E/KPT/2018

DIREKTUR JENDERAL PENGUATAN RISET DAN PENGEMBANGAN  
KEMENTERIAN RISET, TEKNOLOGI, DAN PENDIDIKAN TINGGI



Dentino. Vol. 4 No. 1. Maret 2019. P-ISSN 2337-5310. E-ISSN 2527-4937

Indexed Directory of Open Access Journals (DOAJ),

Sosialisasi Science and Technologi Index (SINTA)

# DENTINO

## JURNAL KEDOKTERAN GIGI

---

Terbit setiap Maret dan September

Terakreditasi : No. 21/E/KPT/2018

### PENGELOLA JURNAL DENTINO

**Penanggung Jawab:**

Prof. Dr. drg. H. Rosihan Adhani, S.Sos, MS  
(Dekan Fakultas Kedokteran Gigi ULM)

**Editor:**

Dr. Maharani Laillyza Apriasari, drg., Sp.PM  
(Fakultas Kedokteran Gigi ULM)

**Sekretaris :**

drg. Amy Nindia Carabelly, M.Si  
(Fakultas Kedokteran Gigi ULM)

**Design Grafis:**

Amalia Husna, Am.Kg  
(Fakultas Kedokteran Gigi ULM)

**Foto Grafi:**

Rifky Ananda, SE  
(Fakultas Kedokteran Gigi ULM)

**Reviewer Internal :**

Dr. Maharani Laillyza Apriasari, drg. Sp.PM; drg. Dewi Puspitasari, M.Si; drg. Amy Nindia Carabelly, M.Si; drg. Selviana Rizky Pramitha.

**Reviewer Ekternal :**

**Dr. Iin Eliana Triwahyuni, drg., M.Si** (Fakultas Kedokteran Gigi Universitas Jember); **Tania Saskianti, drg, Sp.KGA (K).Phd** (Fakultas Kedokteran Gigi Universitas Airlangga); **drg. Ervina Restiwulan Winoto, M.Kes, Sp.Orto** (Fakultas Kedokteran Gigi Universitas Airlangga); **drg. Dr. Eka Fitria Agustina, drg., M.Kes, Sp.Perio** (Fakultas Kedokteran Gigi Universitas Airlangga).

# DENTINO

## JURNAL KEDOKTERAN GIGI

---

Terbit setiap Maret dan September

Terakreditasi : No. 21/E/KPT/2018

### DAFTAR ISI

THE EFFECT OF MAULI BANANA STEM EXTRACT GEL ( <i>Musa acuminata</i> ) APPLICATION IN CONCENTRATION OF 37.5% ON EPITHELIAL THICKNESS (Research report)	
<i>Monica Thiodora Limay, Maharani Laillyza Apriasari, Irham Taufiqurrahman</i>	1-8
MUSCULOSKELETAL DISORDERS (MSDs) RISK ANALYSIS IN WEST SUMATERA DENTISTS AND DENTAL STUDENTS (Research report)	
<i>Abu Bakar, Valendriyani N, Elyza DG, Nadhifa S, Chusnul</i>	9-13
THE EFFECT OF GIANT SNAKEHEAD ( <i>Channa micropeltes</i> ) EXTRACT ON THE NUMBER OF MACROPHAGE IN DIABETES MELITUS WOUND HEALING (Research report)	
<i>Khairunnida, Amy Nindia Carabelly, Maharani Laillyza Apriasari</i>	14-20
EXFOLIATIVE CHEILITIS (Case report)	
<i>I Wayan Arya K.Firdaus, Maharani Laillyza Apriasari</i>	21-24
MAJOR CONTRIBUTORY FACTORS IN THE INITIATION OF ROOT RESORPTION (Article review)	
<i>Syamsiah Syam, Aries Chandra T, Muhammad Ruslin</i>	25-30
PRIMARY MANAGEMENT OF BURNING MOUTH SYNDROME FOR GENERAL DENTIST (Article review)	
<i>Rinda Prawidiastuti, NurIsmah Gitasari, Yunita Fatmala, Wahyu Hidayat, Indah Suasani Wahyuni</i>	31-36
COMPRESSIVE STRENGTH OF TYPE III GYPSUM MIXED WITH WATER OF DIFFERENT WATER HARDNESS LEVEL (Research report)	
<i>Dewi Puspitasari, Suci Fikriyati, Debby Saputera</i>	37-40

THE DIFFERENCE OF SATISFACTION LEVEL IN JAMINAN KESEHATAN NASIONAL (JKN) AND GENERAL PATIENTS TOWARDS QUALITY OF DENTAL HEALTH SERVICES AT DENTAL CLINIC IN WEST KALIMANTAN (Research report)	
<i>Erdianto SetyaWardhana, Sandy Christiono</i>	41-45
FLAVONOID LEVEL ANALYSIS OF BINJAI LEAF EXTRACT ( <i>Mangifera caesia</i> ) IN ETHANOL, METHANOL, AND N-HEXANE SOLVENTS (Research report)	
<i>Didy Adham, Irham Taufiqurrahman, Zairin Noor Helmi</i>	46-49
FLAVONOID LEVEL TEST ON ETHANOL EXTRACT OF BINJAI LEAF ( <i>Mangifera Caesia</i> ) (Research report)	
<i>Ana Azizah Ansari, Irham Taufiqurrahman, Nurdiana Dewi</i>	50-54
ANTIOXIDANT ACTIVITY OF BINJAI LEAVES ( <i>Mangifera caesia</i> ) ETHANOL EXTRACTS (Research report)	
<i>Anita Diana Putri, Irham Taufiqurrahman, Nurdiana Dewi</i>	55-59
TOTAL FLAVONOID CONTENT ANALYSIS OF RAMANIA LEAVES' EXTRACT USING ETHANOL, METHANOL AND N-HEXANE AS SOLVENTS (Research report)	
<i>Risa Hermalinda, Irham Taufiqurrahman, Zairin Noor Helmi</i>	60-63
THE VARIANCE OF RADIOPAQUE IMAGES COMMONLY FOUND IN PERIAPICAL LESION USING PERIAPICAL RADIOGRAPHIC TECHNIQUE (Article review)	
<i>Indra G, Ria F, Lusi E, Azhari</i>	64-69
ATROPHIC GLOSSITIS AS A CLINICAL SIGN FOR ANEMIA IN THE ELDERLY (Case report)	
<i>Rahmatia Djou, Indah Suasani Wahyuni</i>	70-76
THE EFFICACY OF GREEN TEA ( <i>Camellia sinensis</i> ) LEAVES EXTRACT AS CORROSION INHIBITOR FOR ORTHODONTICS STAINLESS-STEEL WIRE (Research report)	
<i>Neny Roeswahjuni, Delvi Fitriani, Annisa Dwi Wardanianti</i>	77-82

THE EFFECT OF BIOLOGICAL MEMBRANES TO FIBROBLAST PROLIFERATION: PLATELET-RICH FIBRIN RELEASATE VS AMNIOTIC (Research report)	
<i>Rezmelia Sari, Sri Pramestri Lastianny, Alya Nur Fadhilah</i>	83-86
CORRELATIVE STUDY BETWEEN MALOCCLUSION SEVERITY LEVEL AND ORAL HYGIENE STATUS AMONG STUDENTS AT SMP LPP WAKAF FOUNDATION UNIVERSITAS MUSLIM INDONESIA IN 2017 (Research report)	
<i>Rafika COUNSUL, Yustisia Puspitasari, Sarahfin Aslan</i>	87-93
RADIOGRAPHIC FEATURES OF INTERNAL STRUCTURE IN JAW LESIONS (Review article)	
<i>Norlaila Sarifah, Ria N. Firman, Farina Pramanik, Lusi Epsilawati</i>	94-100
MANAGEMENT OF VARICELLA ZOSTER IN ADULT PATIENT (Case report)	
<i>Maharani Laillyza Apriasari, Selviana Rizky Pramitha</i>	101-105
DIFFERENCE IN SETTING TIME AND WORKING TIME INTERVAL OF CONVENTIONAL ZINC PHOSPHATE CEMENT AND NANOPARTICLE ZINC PHOSPHATE CEMENT (Research report)	
<i>Chandra Sari Kurniawati, Delvi Fitriani, Sheila Ananda Putri</i>	106-109

**DENTINO**  
**JURNAL KEDOKTERAN GIGI**  
 Vol IV. No 1. Maret 2019

**FLAVONOID LEVEL ANALYSIS OF BINJAI LEAF EXTRACT (*Mangifera caesia*) IN ETHANOL, METHANOL, AND N-HEXANE SOLVENTS**  
 (Research report)

**Didy Adham<sup>1</sup>, Irham Taufiqurrahman<sup>2</sup>, Zairin Noor Helmi<sup>3</sup>**

<sup>1</sup>Student of the Faculty of Dentistry, Lambung Mangkurat University, Banjarmasin

<sup>2</sup>Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Lambung Mangkurat University, Banjarmasin

<sup>3</sup>Department of Surgery, Orthopedic Division, Department of Medical Education, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin

**ABSTRACT**

**Background:** Many wounds can manifest in various complaints such as pain, swelling and bruising. However, almost all of these wounds can be healed faster by utilizing herbal medicine. Binjai is one of herbal medicine originated from *Mangifera* genus and *Anarcadiaceae* family which roots and stems has been proven to contain secondary metabolites. Flavonoids are one of the metabolites which have the antioxidant, antibacterial and anti-inflammatory effect. This effect may contribute in the acceleration of wound healing process especially through flavanoid compound mechanism. Flavonoids themselves are found in the form of glycosides that are more soluble in polar solvents. Thus, they can be obtained by dissolving binjai leaves in methanol and ethanol solvents.

**Purpose:** To determine differences of total flavonoid in binjai leaf extract using ethanol and methanol solvent.

**Methods:** This type of research is a true experimental study with post-test only control group design. The samples was comprised of 27 repetitions consisting three groups, namely 70% ethanol treatment groups and 70% methanol treatment groups. 70% n-hexane group was used as control. **Results:** There were significant differences between ethanol to methanol group at  $p = 0.000$ , ethanol with n-hexane group at  $p = 0.000$  and between methanol with n-hexane group at  $p = 0.000$ . **Conclusions:** There were differences of total flavonoid in binjai leaf extract using ethanol and methanol solvent.

**Keywords:** binjai leaf, ethanol solvent, methanol solvent, total flavonoids

**Correspondence:** Didy Adham, Faculty of Dentistry, Universitas Lambung Mangkurat, Jl. Veteran N0.128 B, Banjarmasin, Kalimantan Selatan, email: [didyadham21@gmail.com](mailto:didyadham21@gmail.com)

**INTRODUCTION**

Wounds are injuries of body tissues due to physical force, causing disruption in normal tissue continuity thus supposed to be treated. One example of the most frequent injury to be complained in the field of dentistry is a post-extraction wound. Friedman (2007) declared that there are 11 million patients per day who experience complaints such as pain, swelling and postoperative bruising after third molar extraction. Wound care aims to stop bleeding and clean the wound area from foreign objects, dead cells and bacteria for the healing process. In general, wound healing process will occur physiologically.

However, problem may arise along the process which will cause pain and discomfort.<sup>1</sup>

Several methods can promote the acceleration of wound healing process, one of them is by the use of herbal medicines.<sup>1</sup> Herbal remedies are traditional medicines derived from natural ingredients which are provided by nature in the form of plants and have been known and used by Indonesian people. Generally, traditional medicine is more easily accepted by the community because it has been familiarized among them. In addition, this medicine is cheaper and easier to obtain.<sup>2</sup>

Binjai (*Mangifera caesia*) is categorized in *Mangifera* genus and *Anarcadiaceae* family. This

plant can be discovered in South Kalimantan and consumed by people who live in the region. Some experts believe that Kalimantan is the the origin place of this plant. Afterwards, it was brought and cultivated in Bali, Java, Philippines and Thailand. The society of Hulu Tabalong, South Kalimantan and surrounding areas use binjai plant roots as a diabetes medicine. A study by Mustikasari and Ariyani (2008) mentioned that binjai plants are proven to contain secondary metabolites, one of them is flavonoid.<sup>3,4</sup>

Flavonoids are the largest natural phenol compounds found in all green plants. Antioxidant activity from plants is caused by the presence of polyphenol compounds which are known having capability as free radical scavenger, hydrolysis and oxidative enzyme inhibitors and also work as anti-inflammatory agent.<sup>5</sup> Kurniasari (2006) stated that a number of medicinal plants contains flavonoids which possess antioxidant, antibacterial, antiviral, anti-inflammatory, anti-allergic and anticancer effects.<sup>5</sup> The process of wound healing can be accelerated by the antioxidant, antibacterial and anti-inflammatory effect of flavonoids compound by inhibiting the production of sodium oxide which has a role in causing tissue damage, preventing secondary infections in wound area, reducing acute inflammatory exudates and stimulating T lymphocytes and macrophages activation.<sup>1</sup>

Flavonoids act as antioxidants by donating hydrogen atoms to bind metals not only in the form of glucosides, but also the free forms called aglycones.<sup>6</sup> Antioxidants demonstrate an effect as osteoblast stimulator and osteoclasts inhibitor.<sup>6</sup> They can affect bone formation by stimulating alkaline phosphatase activity. With the result that, the osteoblast activity increases.<sup>5</sup> Flavonoids are found in the form of glycosides which are more soluble in polar solvents.<sup>6</sup>

Polar solvents are able to extract compounds such as quaternary alkaloids, phenolic components, carotenoids, tannins, sugars, amino acids and glycosides.<sup>7</sup> Active polar substances such as flavonoids can be dissolved by methanol and ethanol solvents which later affect the type of medicine preparations.<sup>7,8</sup>

According to the explanation above, the researcher is interested in conducting a study to compare the total flavonoid content of binjai leaf extract in ethanol solvents and methanol solvents as a preliminary study for medicine preparations of wound healing process.

## MATERIALS AND METHODS

This study is a pure experimental study with post-test only control group design conducted in August 2015 at Biochemistry Laboratory of Medical

Faculty, Lambung Mangkurat University, Banjarbaru. This study used simple random sampling, consisting of 3 treatments. The number of repetitions for each treatment was determined using Federer formula. Each group required 9 times of repetition, with minimal total of sample number required was 27 samples. The sample in this study was binjai leaves taken from Desa Mandiangin, Kecamatan Karang Intan, Kabupaten Banjar, South Kalimantan.

The data obtained from this study was measured by observing the total number of flavonoids through UV-Vis spectrophotometer method to analyze ethanol, methanol and n-hexane groups using the following formula:

$$y = ax + b$$


Description:

$y$  = absorbance value  
 $x$  = flavonoid levels  
 $a, b$  = constant

Data analysis was performed by testing data normality using Shapiro-Wilk<sup>5</sup> test and its homogeneity using Levene's test. The data was evaluated based on the type of variable measurement scale. It was obtained that variable data was categorized as numerical comparative unpaired more than 2 groups and then analyzed using Oneway ANOVA to test hypothesis with 95% confidence level ( $\alpha = 0.05$ ). The significant differences between each of the test groups can be attained. The analysis continued with Post Hoc LSD (Least Significant Difference) test to compare the significant difference among test groups.


## RESULT

According to the study result of total flavonoid content comparison of binjai leaf extract in ethanol and methanol solvents as a preliminary study, the preparation of medicine preparations for wound healing process revealed that the total flavonoid content of ethanol, methanol and n-hexane groups have different values.



**Figure 1.**Diagram of Average Total Flavonoid Content in Binjai Leaf Extract

The results of data analysis in this study indicate that there are significant differences among the solvents. According to the diagram (figure 1), highest total flavonoids is denoted in methanol group with an average of 0.0522 mgQE / mg, it is followed by ethanol with an average of 0.0382 mgQE / mg and the lowest are in n-hexane group with an average of 0.0121 mgQE / mg.



**Figure 2.**Wavelength Graph of Binjai Leaf Extract

The results also reveal that flavonoids found in binjai leaf were classified in chalcone and auron. This is based on the maximum wavelength result (figure 2) which is at 250 nm (band II) and 390 nm (band I).

## DISCUSSION

According to the results of this study, methanol solvent has higher effectivity in binding flavonoids compared to ethanol and n-hexane solvents. Markom (2007) declared that this is caused by different polarity of each solvents, that is 5.2 for ethanol solvent, 5.6 for methanol solvent and 0.1 for n-hexane solvent.<sup>7</sup>

It is also mentioned that total flavanoid content is directly proportional to the polarity levels of solvents where the highest to the lowest values are methanol, ethanol and n-hexane respectively. According to the Rijkstatement (2005), the higher the level of polarity of a solvent, the higher the flavonoids that can be extracted. It is related with the number of unsubstituted hydroxyl groups. Hence, flavonoids can be extracted using polar solvents.

The results also claimed that the types of flavonoids contained in binjai leaf were chalcone and auron. This can be observed through the maximum wavelength which is 250 nm (band II) and 390 nm (band I). According to an article written by Neldawati (2013), it is known that flavonoids in the form of chalcone have a maximum absorption band area in the range of 230-270 nm (band II) and 340-390 nm (band I). Whereas, flavonoids in the form of auron have a maximum absorption band area in the range of 230-270 nm (band II) and 380-430 nm (band I).<sup>8</sup>

Chalcone-type flavanoids are secondary metabolites of several flavonoids found, especially in plants whereas auron is a derivate of chalcone that has cyclization. In spite of the spread in a variety of plant families, chalcone and auron are among the type of flavonoids which are rarely found because of their limited number compared to other flavonoid compounds. This resulted the compound to be classified into minor category because the small percentage in plants and structure variety.<sup>9,10,11</sup> A number of chalcone compounds and their derivatives have been known to have biological activities such as anticancer, anti-inflammatory, antioxidant, antitumor, antimicrobial, antifungal, antibacterial, antimutagenic and hypo-allergenic.<sup>9,11</sup>

Based on the conducted research, it can be concluded that there are differences in the total flavanoid content of binjai leaf dissolved in ethanol solvent and methanol solvent.

## REFERENCES

1. Susilawati, Khafid M, Tiarisna HN, Narendra KW, Chotimah C. Potensi Kulit dan Biji Kelengkeng (*Euphoria longan*) sebagai Gel Topikal untuk Mempercepat Penyembuhan Luka pasca Ekstraksi Gigi. BIMKGI 2013; 1(2): 1-3.
2. Lumbessy M, Abidjulu J, Paendong JJE. Uji Total Flavonoid Pada Beberapa Tanaman Obat Tradisional Di Desa Waitina Kecamatan Mangoli Timur Kabupaten Kepulauan Sula Provinsi Maluku Utara. JURNAL MIPA UNSRAT ONLINE 2013; 1: 50-55.
3. Mustikasari K, Ariyani D. Studi Potensi Binjai (*Mangifera caesia*) dan Kasturi (*Mangifera casturi*) Sebagai Antidiabetes Melalui Skrining Fitokimia pada Akar dan Batang. Sains dan Terapan Kimia 2008; 2: 64-73.



4. Rosyidah K, Siska, Astuti MD. Isolasi senyawa antioksidan dari kulit batang tumbuhan binjai (*Mangifera caesia*). *Jurnal Sains dan Terapan Kimia*, 2011; 5(1): 8.
5. Hermizi H, Hasmah A, Ima N S. The Potential Role of *Quercus infectoria* Gall Extract on Osteoblast Function and Bone Metabolism. *Open Journal of Endocrine and Metabolic Diseases*, 2012; 2: 82-88.
6. Donia. Phytochemical screening and insecticidal activity of three plants from chenopodiaceae family. *Journal of Medicinal Plants Research* 2012; 6(48): 5863-5867.
7. Markom M, Hasan M, Daud WRW, Singh H, Jaim JM. Extraction of hydrolysable tannins from *Phyllanthus niruri* Linn: Effects of solvents and extraction methods. *Separation and Purification Technology* 2007; 52: 487-496.
8. Neldawati, Ratnawulan, Gusnedi. Analisis Nilai Absorbansi dalam Penentuan Kadar Flavonoid untuk Berbagai Jenis Daun Tanaman Obat. *PILLAR OF PHYSICS* 2013; 2: 76-83.
9. Jasril, Teruna HY, Zamri A, Alfatos D, Yuslinda E, Nurulita Y. Sintesis dan Uji Antibakteri Senyawa Bromo Kalkon Piridin. *Jurnal Natur Indonesia* 2012; 14: 172-175.
10. Eryanti Y, Zamri A, Jasril, Rahmita. Sintesis Turunan 2'-hidroksi Kalkon melalui Kondensasi *Claisen-Schmidt* dan Uji Aktivasnya sebagai Antimikroba. *Jurnal Natur Indonesia* 2010; 12 (2): 223-227.
11. Andarwulan N, Faradilla RHF. Senyawa Fenolik pada Beberapa Sayuran Indigenus dari Indonesia. SEAFast center IPB: Bogor. 2012. Hal: 33-48.