

2021-8 Syarifuddin K Jurnal Internasional _ARInt Land- CLASSIFIERS FOR LAND USE Juni 2021

by Syarifuddin Kadir

Submission date: 12-May-2023 06:44PM (UTC-0400)

Submission ID: 2091731447

File name: Internasional_ARInt_Land-CLASSIFIERS_FOR_LAND_USE_Juni_2021.pdf (1.32M)

Word count: 6293

Character count: 37162

COMPARISON OF MAXIMUM LIKELIHOOD AND SUPPORT VECTOR MACHINE CLASSIFIERS FOR LAND USE/LAND COVER MAPPING USING MULTITEMPORAL IMAGERY

Nurlina¹, Syarifuddin Kadir², Ahmad Kurnain³, Wahyuni Ilham²

¹ Doctoral Program of Agriculture Science, Postgraduate Lambung Mangkurat University, Banjarbaru, South Kalimantan,

² Faculty of Forestry, Lambung Mangkurat University, Banjarbaru 70714, South Kalimantan,

³ Faculty of Agriculture, Lambung Mangkurat University, Banjarbaru 70714, South Kalimantan, INDONESIA

Address

Physics Study Program, Mathematic and Natural Science Faculty of Lambung Mangkurat University

Jl. Ahmad Yani km. 36 Banjarbaru City

South Kalimantan Province

Indonesia

E-mail : nurlina_abdullah@ulm.ac.id

COMPARISON OF MAXIMUM LIKELIHOOD AND SUPPORT VECTOR MACHINE CLASSIFIERS FOR LAND USE/LAND COVER MAPPING USING MULTITEMPORAL IMAGERY

Nurlina¹, Syarifuddin Kadir², Ahmad Kurnain³, Wahyuni Ilham²

¹ Doctoral Program of Agriculture Science, Postgraduate Lambung Mangkurat University, Banjarbaru, South Kalimantan,

² Faculty of Forestry, Lambung Mangkurat University, Banjarbaru 70714, South Kalimantan,

³ Faculty of Agriculture, Lambung Mangkurat University, Banjarbaru 70714, South Kalimantan, INDONESIA

³²
E-mail : nurlina_abdullah@ulm.ac.id

ABSTRACT

⁴ Most of the previous research was related to non-parametric classification methods, states that the Maximum Likelihood (ML) and Support Vector Machine (SVM) methods are the leading classification methods in producing high accuracy. However, only a small proportion of studies have compared the performance of these two methods using multitemporal remote sensing imagery, particularly on Sentinel-2 and Landsat data. This study tries to evaluate and compare the performance of ML and SVM classifiers to mapping land use/cover using Sentinel-2 and Landsat multitemporal imagery data. The Tabunio watershed with an area of 62.586 ha has been mapped with ten types of land use/cover are water bodies, forest, bare land, residential, plantations, agriculture, swamps, shrubs, pond, and mining. The confusion matrix and the Kappa coefficient was used to assess classification accuracy. All classification results show a high overall accuracy (OA) ranging from 86% to 95%. Among the two classifiers, four data series with different images and sample sizes, SVM produced the highest OA than ML.

⁴²
Keywords: Sentinel-2; Landsat, Maximum Likelihood (ML); Support Vector Machine (SVM);

INTRODUCTION

Land use/cover maps are currently very important and indispensable in various fields, especially for monitoring and management of natural resources, development strategies, and land change studies (Auliana et al., 2018; S. Kadir et al., 2013, 2016; Z. Abidin, 2019). The land use/cover map is one of the most important documents providing information for various applications, such as monitoring land use, environmental services, urban planning, natural resource conservation, agricultural monitoring, and land use/cover change dynamics (Abbas & Jaber, 2020; Gebhardt et al., 2014; Gómez et al., 2016; Guidici & Clark, 2017; Nurlina et al., 2020; Yunus et al., 2018).

⁷
Multiresolution remote sensing satellite imagery data as one of the most important data sources for land cover mapping (Topaloğlu et al., 2016) apart from its wide geographical coverage and efficient cost, it also provides information from semi-detailed to very detailed scale which makes Remote Sensing data irreplaceable (Khatam et al., 2016). Land use/cover maps are usually made based on the approach of several remote sensing image classification methods (Chen et al., 2017; Duro et al., 2012; Imran & He, 2015). However, accuracy and processing time are still challenges for the remote sensing community (Gómez et al., 2016).

¹⁵
Sentinel-2 with the latest generation of earth observation missions from ESA (European

Space Agency) designed for land and coastal applications, including the Sentinel-2 A and Sentinel-2 B satellites launched in June 2015 and March 2017, respectively (Thanh Noi & Kappas, 2017). Sentinel-2 remains active and enhances the mission of Landsat and SPOT (Système Probatoire Observation de la Terre) (Wang et al., 2017). Sentinel-2 has wide coverage capability, high spatial resolution (10–60 m), and temporal resolution (ten days for Sentinel-2 A, B / five days Sentinel-2 A + B), and includes multispectral satellite imagery (13 spectral channels). Sentinel-2 has also received major research attention because of its free access to global coverage. Various applications have been applied with Sentinel-2 A, particularly in land use and land cover mapping, the practicality and effectiveness of Sentinel-2 have been tested and show high application potential (Gao et al., 2017; Yan et al., 2017). However, because this is a new type of satellite image, so there is still little research using Sentinel-2 for land use/cover mapping, more research is needed to conduct and evaluate the usefulness of this satellite imagery. Before Sentinel-2 One of the most important and most widely used digital data for remote sensing work is the Landsat satellite. Landsat satellite missions have continued to collect global imagery since 1972 and monitor the Earth biweekly with a resolution of 30 m x 30 m. Landsat's open access policy in 2008 allowed researchers to access this data freely so that monitoring of previously impossible land cover changes was very easy to implement (Eitel et al., 2011).

Landsat data is also not static either, these satellite data products have evolved rapidly over time, providing more data for researchers, and allowing more accurate classification of different and more advanced processes. Landsat 5-TM (8 bits radiometric) has seven channels, and Landsat 7- ETM+ (9 bits radiometric) has eight spectral bands with a resolution of 30 m. whereas, the latest generation Landsat 8-OLI has 11 bands (12 bit radiometric), and this technology is considered the best choice for environmental analysis (Clevers et al., 2017; Sibanda et al., 2015). Classification using Landsat imagery is not only cost-effective but also accurate for making land cover maps that can be used for environmental management, urban planning, forestry, agriculture, and many other sectors.

According to Lu and Weng (Lu & Weng, 2007), not only image suitability but also the determination of the appropriate classification method can affect the accuracy of land use/cover mapping. Several kinds of literatures with various classification methods have been developed and tested for land use/cover mapping from remote sensing data (Brodley & Friedl, 1997; Li et al., 2014; Wang & Braun, 2009). This method uses a supervised parametric algorithm classifier, namely Maximum Likelihood (ML) and the Support Vector Machine (SVM) algorithm. In recent years, nonparametric methods (machine learning-based algorithms) have become a major concern of remote sensing based applications. The use of the SVM and ML classification algorithms has significantly increased. Most of the studies used the ML method as one of the criteria for comparison with other machine learning algorithms (Basukala et al., 2017; Jhonnerie et al., 2015; Khatami et al., 2016).

Several studies have been conducted to determine the best classification method for land use/cover mapping trying to compare the accuracy of these classifiers with either the same method or with different classification methods. However, the conclusions are quite mixed. Additionally, to the best of our knowledge, only a small of published studies have compared and evaluated the time series performance SVM and ML against different satellite imagery, particularly in Indonesia. Therefore, the purpose of this study was to compare and evaluates the performance of the two best classifier ML and SVM for land use/cover mapping in the Tenujo watershed of South Kalimantan using multi-spatial-temporal satellite data from Sentinel-2 and Landsat data series.

48 MATERIALS AND METHODS

Materials used in this study include a 1: 50,000 scale digital map of the Indonesian Earth Administration

(RBI), as a reference for administrative boundaries at the research location. Obtained from the Geospatial Information Agency (BIG). Landsat TM, ETM+ /OLI 8 OLI TIRS Satellite Imagery 2005, 2010, 2015, 2020 which is multispectral data with a spatial resolution of 30 m downloaded from <http://earthexplorer.usgs.gov/> in 117 row 62. Sentinel-2 the year 2020 Satellite Imagery which can be accessed through the website <https://sentinel.esa.int/web/sentinel/sentinel-data-access>. The details of the Landsat images used for this study are provided in Table 1.

Table 1. Satellite images used in this study

Year	2005	2010	2015	2020
Months	15-October	20-November	18-November	12- September
Sensor	Landsat ETM+	Landsat ETM+	Landsat OLI	Sentinel 2

Study Area

The research location is in the Tabunio Watershed (DAS) which is located in Tanah Laut Regency with an area of 62,558.56 ha which is geographically located at 3 ° 37'2.72 " -3 ° 51'51.43" LS and 114 ° 36'12.02 " -114 ° 57'47.62" East Longitude. The Tabunio watershed from upstream to downstream is a rural, urban, and coastal area with a distinctive heterogeneous land cover, covering ten land cover classes, namely: settlements, plantations, rice fields, bare land, mining, forest, swamps, shrubs, ponds, and water bodies. The Tabunio watershed consists of 44 villages administratively, 4 sub-districts, and 10 sub-watersheds (ecologically). The map of the Tabunio watershed research location is shown in Figure 1.




Figure 1. Location of the study area in Tabunio Watershed

TM is Thematic Mapper, ETM+ is Enhanced Thematic Mapper, and OLI is Operational Land Imager. The maximum likelihood classification is calculated using the following discriminant functions for each pixel.

$$g_i(x) = \ln p(\omega_i) - \frac{1}{2} \ln |\Sigma_i| - \frac{1}{2} (x - m_i)^t \Sigma_i^{-1} (x - m_i)$$

where i = class, x = n -dimensional data (where n is the number of bands), $p(\omega_i)$ = probability that class ω_i occurs in the image and is assumed same for all classes, $|\Sigma_i|$ = determinant of the covariance matrix of the data in a class, Σ_i^{-1} = its inverse matrix, m_i = mean vector of a class.

SVM algorithm finds a hyperplane to separate the database based on a pre-defined number of categories (Mountrakis et al., 2011). SVMs approach is generally organized into four Kernel functions: linear, polynomial, radial basis function (RBF), and sigmoid (Kavzoglu & Colkesen, 2009; Lee et al., 2017). RBFs are more powerful kernels than others (linear, polynomial, radial) and are used to achieve better results (Rimal et al., 2020). The following are the equation of each Kernel functions used in SVM:

- (i) Linear: $K(x_i, y_i) = x_i^T \cdot x_j$
- (ii) Polynomial: $K(x_i, y_i) = (g \cdot x_i^T \cdot x_j + r)^d, g > 0$
- (iii) Radial basis function: $K(x_i, y_i) = e^{-g(x_i - x_j)^2}, g > 0$
- (iv) Sigmoid: $K(x_i, y_i) = \tanh(g \cdot x_i^T \cdot x_j + r)$

where g , d and, r are user-controlled parameters of kernel function

Training and Testing Sample Datasets

Training data (training and test samples) is drawn based on manual interpretation of the Sentinel-2 and Landsat data series and available high-resolution imagery from Google Earth. To collect training sample data, the polygon generates tool in the ArcGIS 10.5 toolbox is used to create polygons for each land cover class. Due to the different polygon sizes, the number of pixels for each land cover class is also different (Table 2).

Table 2. Training and testing sample sizes were used in this study.

Land Cover	Training sample (pixels)				Testing sample (pixels)			
	2005	2010	2015	2020	2005	2010	2015	2020
Water body	354	450	348	403.335	406	124	86	159.505
Forest	259	56.082	17.044	8.996.502	314	38.261	5.121	2.455.313
Bare land	21.804	19.716	4.643	6.723.879	22.193	10.246	890	4.258.246
Residential	1.142	4.882	3.152	2.136.693	1.188	1.903	351	1.083.735
Plantation	893	31.736	37.881	19.184.554	937	14.380	7.776	9.444.056
Agriculture	1.351	1.314	876	4.154.898	1.596	713	200	1.308.608
Swamp	16.741	334	370	148.135	17.183	145	66	16.405
Shrubs	204	251	532	133.906	251	88	134	50.072
Pond	166	330	73	43.405	322	194	18	45.559
Mining	724	3.159	2.182	330.744	738	2.565	279	172.107

We took ten land cover categories, namely: water body, forest, bare land, residential, plantations, agriculture, swamps, shrubs, pond, mining. A total of 86,345 sample points were used for training and then tested for accuracy assessment. Training samples are often used for accuracy assessments (Jensen, 1996; Sexton et al., 2013; Sloan & Pelletier, 2012). The accuracy classifications were observed based on field survey data and high-resolution satellite imagery from Google Earth that was taken randomly for each land cover class. Furthermore, overall accuracy (OA), user accuracy (UA), and manufacturer accuracy (PA) are calculated and tested using the confusion matrix and Kappa coefficients.

RESULT

Major transformations include an increase in housing and plantations and a sharp decline in forests and shrubs. residential coverage in the study area increased by 1,382.18 hectares, (from 619.07 hectares in 2005 to 2,001.24 hectares in 2020) with an increase of 223.27%, while plantations increased by 23,811.15 hectares, (from 502, 16 hectares) in 2005 to 24,313.31 hectares in 2020) with an increase of 474.2%, while forests decreased by 3,056.79 hectares (from 16,223.67 hectares to 13,166.88 hectares) while shrubs decreased by 9,630, 20 hectares (from 10,846.53 acres to 1,216.33 acres) from 2005 to 2020. The historical changes in land cover in the study area during 2005-2020 are shown in Figures 4a-d, Table 3 and Table 4.




Figure 2. Land cover map classified based on SVM approach a 2005; b 2010; c 2015; d 2020

Table 3. Land Use/Land Cover Data in Tabunio Watershed 2005 – 2020

Land Use/Land Cover	Tahun			
	2005 (acres)	2010 (acres)	2015 (acres)	2020 (acres)
Water body	592,64	386,64	368,50	406,48
Forest	16.223,67	14.004,85	14.699,89	13.166,88
Bare land	3.712,99	4.945,80	13.247,55	7.906,35
Residential	619,07	991,83	1.451,73	2.001,24
Plantation	502,16	7.710,81	20.866,44	24.313,31
Agriculture	21.021,27	10.313,42	8.366,95	12.917,27
Swamp	6.759,52	3.818,56	161,37	181,88
Shrubs	10.846,53	17.042,34	1.695,94	1.216,33
Pond	45,88	126,24	47,96	36,14
Mining	2.172,66	3.155,88	1.590,04	350,50

Table 4. Land use/land cover change between 2005 and 2020

Land Use/Land Cover	2005–2010 (acres)	2010–2015 (acres)	2015–2020 (acres)	2005–2020 (acres)
Water body	-206,00	-18,14	37,98	-186,16
Forest	-2.218,82	695,04	-1.533,01	-3.056,79
Bare land	1.232,82	8.301,75	-5.341,20	4.193,36
Residential	372,77	459,90	549,51	1.382,18
Plantation	7.208,65	13.155,63	3.446,87	23.811,15
Agriculture	-10.707,86	-1.946,46	4.550,32	-8.104,00
Swamp	-2.940,96	-3.657,19	20,50	-6.577,65
Shrubs	6.195,82	-15.346,40	-479,62	-9.630,20
Pond	80,37	-78,28	-11,82	-9,73
Mining	983,22	-1.565,85	-1.239,54	-1.822,17

Accuracy Assessment and Comparisons

In order to assess classification accuracy, there are many methods available in the literature. The confusion matrix and the Kappa coefficient are among the most popular. For several decades, the Kappa coefficient has been rarely used in assessing the classification accuracy of remote sensing data (Heydari & Mountrakis, 2018). One of the drawbacks of OA metrics is the lack of performance classes that are specific to them. He and Garcia (Gautheron et al., 2019) stated that if the input data (training sample) is not balanced, then the OA value could be wrong because the last class will be classified very poorly. He and Garcia (Gautheron et al., 2019) also suggest that when choosing OA as the criterion metric, the class distribution should be followed by those that occur naturally.

In this study, we used a stratified sampling approach; This approach fits well with OA metrics. In addition, to compare the accuracy of each classification method, we used the same training (input) and testing (validation) dataset; thus, the effect of individual class distributions on OA does not bias the results. We also calculated the 95% confidence interval (error tolerance) δ of the probability estimates (Baraldi et al., 2006) for each OA. Since we used the same test dataset for a classification accuracy assessments, δ did not differ significantly. Therefore, to assess and compare the performance of different classifiers and data sets, we used overall accuracy (OA) as the criterion.

In this study, the overall accuracy of the LULC classification achieved by using the SVM classifier is 96.79% (2005), 92.7458% (2010), 90.93% (2015), and 86.20% (2020). The overall classification accuracy of the alternative ML classifier is 94.79% (2005), 88.64% (2010), 85.38% (2015), and 64.20% (2020). The SVM classifier received a higher OA than the ML classifier across all classification years. SVM obtains a maximum accuracy of 96.79% and a minimum of 86.20%, while the ML classification ranges from a minimum of 64.20% in 2020 to a maximum of 94.79% in 2005. The average overall accuracy of SVM is 91.66% and ML 83.25%. The difference in OA between the two classifications shows that SVM has a better accuracy of 10.8% compared to ML in determining the type of land cover.

The SVM classifier identifies all classes more accurately than the ML classifier (Figures 4, 5). For example, during 2005, the highest UA SVM in terms of Forest (99.56%) was seen, while the ML classification for that year was relatively lower (60.5%). Likewise, the highest SVM related to mining, swamps, deforested land, settlement were 98%, 97%, 96.5%, 95%, and 94.5% respectively during 2005, 2010, 2015, and 2020. In contrast, ML classifier for each class in the same year is as follows: 60.5%, 98%, 81%, and 87.56%.




Figure 3. User's accuracy assessment




Figure 4. Producer's accuracy assessment

Producer accuracy (PA) of the SVM classifier is also relatively higher than the highest ML.PA classifier for SVM it was 99.98% for swamps in 2020, while ML was 92.45% for agriculture. PA ponds were found 83.0% in 2015, and forests remained the highest (96.56%) in 2016. Again, 2020 was found to be important for Bare land (98.1%), whereas PA SVM was found to be consistently dominant in 2005, 2015, and 2020. On the other hand, ML was found to be 99.64% in 2005 and bare land was found to be 98.87% in 2005. The ML classification of sand areas for 2013 was 86.67%, and that the water body for 2015 was 94.18 %. The PA yields for ML shrubs observed in 2005, 2010, and 2015 were 81.27%, 97.77%, and 94.65%. The highest UA and PA from SVM classifiers were most seen in the bare land (Figs. 4 and 5), and the lowest UA from SVM was observed in housing (84% during 2005) and the lowest PA SVM in Pond (75.56%) during 2005) in Table 5.

Table 5. Producer's Accuracy dan User's Accuracy between SVM and ML

Land Use/Land cover	2005		2010		2015		2020	
	SVM	ML	SVM	ML	SVM	ML	SVM	ML
<i>(a) User's Accuracy</i>								
Water body	87,19	99,64	55,46	0,92	49,77	81,03	79,24	21,44
Forest	98,25		99,56	91,84	87,17	88,9	60,39	0
Bare land	96,13	98,87	88,57	81,08	98,1	83,97	96,15	85,02
Residential	95,3	87,24	79,78	87,17	84,6	79,65	85,37	40,39
Plantation		56,86	72,68	86,68	92,45	89,32	90,96	0
Agriculture	84,65	98,1	8,75	57,14	24,55	69,21	53,06	44,16
Swamp	97,43	81,29	86,08	71,34	73,19	40,22	99,77	3,61
Shrubs	81,27	29,72	58,27	54,28	43,82	37,4	50,9	28,3
Pond	51,55	94,3	72,77	54,59	8,55	14,68	31,12	31,72
Mining	98,10	60	98,63	73,15	78	27,63	67,17	42,85
<i>(b) Producer's Accuracy</i>								
Water body	87,84	94,52	91,08	16,49	86,36	42,18	74,95	53,73
Forest	96,72	95,25	93,67	86,34	96,56	86,16	83,41	0
Bare land	98,28	97,76	93,51	9,05	95,92	78,16	66,19	72,43
Residential	92,92	94	94,46	62,01	73,15	55,2	85,17	66,31
Plantation		87,09	94,74	85,21	96,01	83,83	74,56	0
Agriculture	79,8	93,67	1,38	86,08	46,01	54,32	87,54	92,45
Swamp	99,36	91,7	59,02	25	81,58	50,71	99,98	44,44
Shrubs	70,59	91,36	52,59	74,54	46,85	77,75	53,72	86,44
Pond	83	98,51	79,89	79,2	11,16	86,43	23,33	23,1
Mining	98,5	94,52	60,83	69,06	29,17	69,86	80,9	71,41

SVM = Support Vecor Machine, ML = Maximum Likelihood

DISCUSSION

SVM and ML are well-known methods for assessing the accuracy of land cover classification in any area (Bray & Han, 2004; Srivastava et al., 2012). ML is a classical parametric classification method used assuming multivariate normally distributed data (Kavzoglu & Colkesen, 2009). In particular, SVM yields better accurate land cover classification due to its nonparametric nature (Rimal et al., 2020; Thanh Noi & Kappas, 2017; Vapnik & Chervonenkis, 2015). SVM reduces the misclassification of land cover from hidden information or controls the level of certain misclassifications. SVM and ML are very popular in land cover classification because they have higher accuracy compared to other algorithms in identifying land cover classes in watersheds and others (Bray & Han, 2004; Huang et al., 2002; Kavzoglu & Colkesen, 2009; Schneider, 2012). However, (Campbell, 1981; Higin et al., 1980; Scholz et al., 1979) argued that the selection of sample data (training data) is more important than the selection of a classification algorithm to achieve a higher classification accuracy of the classified images. Accuracy assessment is an important stage and must be carried out in the classification and mapping of land cover (Lin, 2017). Accuracy assessment refers to the analysis process carried out to show the level of truth of a map or classification results (Foody, 2002). Accuracy assessments are carried out to assess map quality, evaluate various

classification algorithms and identify errors. The assessment and validation of land cover maps provide a measure of data quality including overall accuracy, user accuracy, and producer accuracy. In an assessment, high accuracy means that misclassification of land cover is low. Producer accuracy states how well a certain area can be classified, and user accuracy ensures that the pixels classified in the image exactly match the category in the field (Congalton, 1991). Accuracy assessment is fundamental but challenging in the thematic mapping (Foody, 2002).

CONCLUSIONS

Higher user and producer accuracy are obtained from the SVM classification method compared to the ML classifier. SVM has proven to be effective in determining land cover classification, especially on bare land. This is associated with higher accuracy ratings due to different signatures. However, the different signatures of bare land also result in a higher accuracy of the ML classification method. Of the ten total land cover classes, the highest accuracy of users and producers is seen in bare land, while the accuracy of users and producers in pond classes is lower. Based on the evidence obtained from our study, we recommend SVM as a suitable option for proper land use/land cover classification, particularly in heterogeneous areas such as watershed area.

REFERENCES

- [1]. Abbas, Z., & Jaber, H. S. (2020). Accuracy assessment of supervised classification methods for extraction land use maps using remote sensing and GIS techniques. *IOP Conference Series: Materials Science and Engineering*, 745(1). <https://doi.org/10.1088/1757-899X/745/1/012166>
- [2]. Auliana, A., Ridwan, I., & Nurlina, N. (2018). Analisis Tingkat Kekritisan Lahan di DAS Tabunio Kabupaten Tanah Laut. *Positron*, 7(2), 54. <https://doi.org/10.26418/positron.v7i2.18671>
- [3]. Baraldi, A., Puzzolo, V., Blonda, P., Bruzzone, L., & Tarantino, C. (2006). Automatic spectral rule-based preliminary mapping of calibrated landsat TM and ETM+ images. *IEEE Transactions on Geoscience and Remote Sensing*, 44(9), 2563–2585. <https://doi.org/10.1109/TGRS.2006.874140>
- [4]. Basukala, A. K., Oldenburg, C., Schellberg, J., Sultanov, M., & Dubovyk, O. (2017). Towards improved land use mapping of irrigated croplands: performance assessment of different image classification algorithms and approaches. *European Journal of Remote Sensing*, 50(1), 187–201. <https://doi.org/10.1080/22797254.2017.1308235>
- [5]. Bray, M., & Han, D. (2004). *modelling*. 265–280.
- [6]. Brodley, C. E., & Friedl, M. A. (1997). Decision tree classification of land cover from remotely sensed data. *Remote Sensing of Environment*, 61(3), 399–409. [https://doi.org/10.1016/S0034-4257\(97\)00049-7](https://doi.org/10.1016/S0034-4257(97)00049-7)
- [7]. Campbell, J. B. (1981). Spatial correlation effects upon accuracy of supervised classification of land cover. *Photogrammetric Engineering and Remote Sensing*, 47(3), 355–363.
- [8]. Chen, Y., Dou, P., & Yang, X. (2017). Improving land use/cover classification with a multiple classifier system using AdaBoost integration technique. *Remote Sensing*, 9(10). <https://doi.org/10.3390/rs9101055>
- [9]. Clevers, J. G. P. W., Kooistra, L., & van den Brande, M. M. M. (2017). Using Sentinel-

- 2 data for retrieving LAI and leaf and canopy chlorophyll content of a potato crop. *Remote Sensing*, 9(5), 1–15. <https://doi.org/10.3390/rs9050405>
- [10]. Congalton, R. G. (1991). A review of assessing the accuracy of classifications of remotely sensed data. *Remote Sensing of Environment*, 37(1), 35–46. [https://doi.org/10.1016/0034-4257\(91\)90048-B](https://doi.org/10.1016/0034-4257(91)90048-B)
- [11]. Duro, D. C., Franklin, S. E., & Dubé, M. G. (2012). A comparison of pixel-based and object-based image analysis with selected machine learning algorithms for the classification of agricultural landscapes using SPOT-5 HRG imagery. *Remote Sensing of Environment*, 118, 259–272. <https://doi.org/10.1016/j.rse.2011.11.020>
- [12]. Eitel, J. U. H., Vierling, L. A., Litvak, M. E., Long, D. S., Schulthess, U., Ager, A. A., Krofcheck, D. J., & Stoscheck, L. (2011). Broadband, red-edge information from satellites improves early stress detection in a New Mexico conifer woodland. *Remote Sensing of Environment*, 115(12), 3640–3646. <https://doi.org/10.1016/j.rse.2011.09.002>
- [13]. Foody, G. M. (2002). Status of land cover classification accuracy assessment. In *Remote Sensing of Environment* (Vol. 80, Issue 1, pp. 185–201). Elsevier Inc. [https://doi.org/10.1016/S0034-4257\(01\)00295-4](https://doi.org/10.1016/S0034-4257(01)00295-4)
- [14]. Gao, Q., Zribi, M., Escorihuela, M. J., & Baghdadi, N. (2017). Synergetic use of sentinel-1 and sentinel-2 data for soil moisture mapping at 100 m resolution. *Sensors (Switzerland)*, 17(9). <https://doi.org/10.3390/s17091966>
- [15]. Gautheron, L., Habrard, A., Morvant, E., & Sebban, M. (2019). Metric learning from imbalanced data. *Proceedings - International Conference on Tools with Artificial Intelligence, ICTAI, 2019-Novem(9)*, 923–930. <https://doi.org/10.1109/ICTAI.2019.00131>
- [16]. Gebhardt, S., Wehrmann, T., Ruiz, M. A. M., Maeda, P., Bishop, J., Schramm, M., Kopeinig, R., Cartus, O., Kelldorfer, J., Ressler, R., Santos, L. A., & Schmidt, M. (2014). MAD-MEX: Automatic wall-to-wall land cover monitoring for the Mexican REDD-MRV program using all Landsat data. *Remote Sensing*, 6(5), 3923–3943. <https://doi.org/10.3390/rs6053923>
- [17]. Gómez, C., White, J. C., & Wulder, M. A. (2016). Optical remotely sensed time series data for land cover classification: A review. *ISPRS Journal of Photogrammetry and Remote Sensing*, 116, 55–72. <https://doi.org/10.1016/j.isprsjprs.2016.03.008>
- [18]. Guidici, D., & Clark, M. L. (2017). One-dimensional convolutional neural network land-cover classification of multi-seasonal hyperspectral imagery in the San Francisco Bay Area, California. *Remote Sensing*, 9(6). <https://doi.org/10.3390/rs9060629>
- [19]. Heydari, S. S., & Mountrakis, G. (2018). Effect of classifier selection, reference sample size, reference class distribution and scene heterogeneity in per-pixel classification accuracy using 26 Landsat sites. *Remote Sensing of Environment*, 204(September), 648–658. <https://doi.org/10.1016/j.rse.2017.09.035>
- [20]. Hixson, M., Schulz, D., Fuh, N., & Akiyama, T. (1980). Evaluation of several schemes for classification of remotely sensed data. *Photogrammetric Engineering & Remote Sensing*, 46(12), 1547–1553.
- [21]. Huang, C., Davis, L. S., & Townshend, J. R. G. (2002). International Journal of Remote Sensing An assessment of support vector machines for land cover classification An assessment of support vector machines for land cover classification. *International Journal of Remote Sensing*, 23(4), 725–749.

- <http://www.tandfonline.com/loi/tres20%5Cnhttp://dx.doi.org/10.1080/01431160110040323%5Cnhttp://www.tandfonline.com/page/terms-and-conditions>
- [22]. Imran, F. M., & He, M. (2015). Extended morphological profiles with duality for hyperspectral image classification. *Workshop on Hyperspectral Image and Signal Processing, Evolution in Remote Sensing, 2015-June(9)*, 4768–4786. <https://doi.org/10.1109/WHISPERS.2015.8075493>
- [23]. Jensen, J. R. (1996). *Introductory digital processing: a remote sensing perspective*. Upper Saddle River, NJ: Prentice-Hall.
- [24]. Jhonnerie, R., Siregar, V. P., Nababan, B., Prasetyo, L. B., & Wouthuyzen, S. (2015). Random Forest Classification for Mangrove Land Cover Mapping Using Landsat 5 TM and Alos Palsar Imageries. *Procedia Environmental Sciences*, 24, 215–221. <https://doi.org/10.1016/j.proenv.2015.03.028>
- [25]. Kadir, S., Rayes, M. L., Ruslan, M., & Kusuma, Z. (2013). Infiltration to control flood vulnerability: a case study of rubber plantation of Dayak Deah community in Negara sub-watershed, South Kalimantan Province. *Academic Research International*; 2013, 4(5), 1–13. <http://www.savap.org.pk/journals...>
- [26]. Kadir, Syarifuddin, Badaruddin, Nurlina, Ridwan, I., & Rianawaty, F. (2016). The recovery of tabunio watershed through enrichment planting using ecologically and economically valuable species in south Kalimantan, Indonesia. *Biodiversitas*, 17(1), 140–147. <https://doi.org/10.13057/biodiv/d170121>
- [27]. Kavzoglu, T., & Colkesen, I. (2009). A kernel functions analysis for support vector machines for land cover classification. *International Journal of Applied Earth Observation and Geoinformation*, 11(5), 352–359. <https://doi.org/10.1016/j.jag.2009.06.002>
- [28]. Khatami, R., Mountrakis, G., & Stehman, S. V. (2016). A meta-analysis of remote sensing research on supervised pixel-based land-cover image classification processes: General guidelines for practitioners and future research. *Remote Sensing of Environment*, 177, 89–100. <https://doi.org/10.1016/j.rse.2016.02.028>
- [29]. Lee, S., Hong, S. M., & Jung, H. S. (2017). A support vector machine for landslide susceptibility mapping in Gangwon Province, Korea. *Sustainability (Switzerland)*, 9(1), 15–19. <https://doi.org/10.3390/su9010048>
- [30]. Li, C., Wang, J., Wang, L., Hu, L., & Gong, P. (2014). Comparison of classification algorithms and training sample sizes in urban land classification with landsat thematic mapper imagery. *Remote Sensing*, 6(2), 964–983. <https://doi.org/10.3390/rs6020964>
- [31]. Lin, S.-K. (2013). *Introduction to Remote Sensing*. Fifth Edition. By James B. Campbell and Randolph H. Wynne, The Guilford Press, 2011; 662 pages. Price: £80.75, ISBN 978-1-60918-176-5. *Remote Sensing*, 5(1), 282–283. <https://doi.org/10.3390/rs5010282>
- [32]. Lu, D., & Weng, Q. (2007). A survey of image classification methods and techniques for improving classification performance. *International Journal of Remote Sensing*, 28(5), 823–870. <https://doi.org/10.1080/01431160600746456>
- [33]. Mountrakis, G., Im, J., & Ogole, C. (2011). Support vector machines in remote sensing: A review. *ISPRS Journal of Photogrammetry and Remote Sensing*, 66(3), 247–259. <https://doi.org/10.1016/j.isprsjprs.2010.11.001>
- [34]. Nurlina, Ridwan, I., Hadi Haryanti, N., & Hanik, U. (2020). Analysis of Environmental

- Vulnerability Using Satellite Imagery and Geographic Information System in Coal Mining Area Planning at Banjar Regency. *IOP Conference Series: Earth and Environmental Science*, 499, 012014. <https://doi.org/10.1088/1755-1315/499/1/012014>
- [35]. Rimal, B., Rijal, S., & Kunwar, R. (2020). Comparing Support Vector Machines and Maximum Likelihood Classifiers for Mapping of Urbanization. *Journal of the Indian Society of Remote Sensing*, 48(1), 71–79. <https://doi.org/10.1007/s12524-019-01056-9>
- [36]. Saputra, A. E., Ridwan, I., & Nurlina, N. (2019). *Analisis Tingkat Resapan Air Menggunakan Sistem Informasi Geografis di Das Tabunio. 1*, 149–158.
- [37]. Schneider, A. (2012). Monitoring land cover change in urban and peri-urban areas using dense time stacks of Landsat satellite data and a data mining approach. *Remote Sensing of Environment*, 124, 689–704. <https://doi.org/10.1016/j.rse.2012.06.006>
- [38]. Scholz, D., Fuhs, N., & Hixson, M. (1979). *An evaluation of several different classification schemes - Their parameters and performance. [maximum likelihood decision for crop identification]*.
- [39]. Sexton, J. O., Song, X. P., Huang, C., Channan, S., Baker, M. E., & Townshend, J. R. (2013). Urban growth of the Washington, D.C.-Baltimore, MD metropolitan region from 1984 to 2010 by annual, Landsat-based estimates of impervious cover. *Remote Sensing of Environment*, 129, 42–53. <https://doi.org/10.1016/j.rse.2012.10.025>
- [40]. Sibanda, M., Mutanga, O., & Rouget, M. (2015). Examining the potential of Sentinel-2 MSI spectral resolution in quantifying above ground biomass across different fertilizer treatments. *ISPRS Journal of Photogrammetry and Remote Sensing*, 110, 55–65. <https://doi.org/10.1016/j.isprsjprs.2015.10.005>
- [41]. Sloan, S., & Pelletier, J. (2012). How accurately may we project tropical forest-cover change? A validation of a forward-looking baseline for REDD. *Global Environmental Change*, 22(2), 440–453. <https://doi.org/10.1016/j.gloenvcha.2012.02.001>
- [42]. Srivastava, P. K., Han, D., Rico-Ramirez, M. A., Bray, M., & Islam, T. (2012). Selection of classification techniques for land use/land cover change investigation. *Advances in Space Research*, 50(9), 1250–1265. <https://doi.org/10.1016/j.asr.2012.06.032>
- [43]. Thanh Noi, P., & Kappas, M. (2017). Comparison of Random Forest, k-Nearest Neighbor, and Support Vector Machine Classifiers for Land Cover Classification Using Sentinel-2 Imagery. *Sensors (Basel, Switzerland)*, 18(1). <https://doi.org/10.3390/s18010018>
- [44]. Topaloğlu, R. H., Sertel, E., & Musaoğlu, N. (2016). Assessment of classification accuracies of Sentinel-2 and Landsat-8 data for land cover/use mapping. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences - ISPRS Archives*, 41(June), 1055–1059. <https://doi.org/10.5194/isprsarchives-XLI-B8-1055-2016>
- [45]. Vapnik, V. N., & Chervonenkis, A. Y. (2015). On the uniform convergence of relative frequencies of events to their probabilities. *Measures of Complexity: Festschrift for Alexey Chervonenkis, XVI(2)*, 11–30. https://doi.org/10.1007/978-3-319-21852-6_3
- [46]. Wang, Q., Blackburn, G. A., Onojeghuo, A. O., Dash, J., Zhou, L., Zhang, Y., & Atkinson, P. M. (2017). Fusion of Landsat 8 OLI and Sentinel-2 MSI Data. *IEEE Transactions on Geoscience and Remote Sensing*, 55(7), 3885–3899. <https://doi.org/10.1109/TGRS.2017.2683444>

- [47]. Waske, B., & Braun, M. (2009). Classifier ensembles for land cover mapping using multitemporal SAR imagery. *ISPRS Journal of Photogrammetry and Remote Sensing*, 64(5), 450–457. <https://doi.org/10.1016/j.isprsjprs.2009.01.003>
- [48]. Yang, X., Zhao, S., Qin, X., Zhao, N., & Liang, L. (2017). Mapping of urban surface water bodies from sentinel-2 MSI imagery at 10 m resolution via NDWI-based image sharpening. *Remote Sensing*, 9(6), 1–19. <https://doi.org/10.3390/rs9060596>
- [49]. Yunus, M. Z. M., Ahmad, F. S., & Ibrahim, N. (2018). Mangrove vulnerability index using GIS. *AIP Conference Proceedings*, 1930. <https://doi.org/10.1063/1.5022901>
- [50]. Z. Abidin, Nurlina, I. R. (2019). Hubungan Geomorfometri dengan Bentuk dan Penggunaan Lahan Sub-Sub DAS Riam Kanan Kabupaten Banjar Kalimantan Selatan. *Jurnal Fisika FLUX*, 16(2), 131–140. <https://doi.org/https://doi.org/10.20527/flux.v16i2.5935>

2021-8 Syarifuddin K Jurnal Internasional _ARInt Land-CLASSIFIERS FOR LAND USE Juni 2021

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

17%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Lamyaa Gamal El-deen Taha, Rania Elsayed Ibrahim. "Assessment of Approaches for the Extraction of Building Footprints from Pléiades Images", Geomatics and Environmental Engineering, 2021
Publication 1%
- 2 ediss.uni-goettingen.de
Internet Source 1%
- 3 Kuncahyo Setyo Nugroho, Anantha Yullian Sukmadewa, Angga Vidianto, Wayan Firdaus Mahmudy. "Effective predictive modelling for coronary artery diseases using support vector machine", IAES International Journal of Artificial Intelligence (IJ-AI), 2022
Publication 1%
- 4 www.mdpi.com
Internet Source 1%
- 5 pdfs.semanticscholar.org
Internet Source 1%

6

Marina-Ramona Rujoiu-Mare, Bogdan Olariu, Bogdan-Andrei Mihai, Constantin Nistor, Ionuț Săvulescu. "Land cover classification in Romanian Carpathians and Subcarpathians using multi-date Sentinel-2 remote sensing imagery", European Journal of Remote Sensing, 2017

Publication

<1 %

7

Duinkherjav Bayarmaa, Gankhuyag Gantulga, Chuluunbaatar Gantigmaa, Dashzeveg Ganbat. "Classification of Pest Damaged Trees with Multi-Temporal Sentinel-2 Data", Proceedings of the Institute of Biology, 2022

Publication

<1 %

8

www.int-arch-photogramm-remote-sens-spatial-inf-sci.net

Internet Source

<1 %

9

journal.ugm.ac.id

Internet Source

<1 %

10

S. Thakur, I. Mondal, P. B. Ghosh, P. Das, T. K. De. "A review of the application of multispectral remote sensing in the study of mangrove ecosystems with special emphasis on image processing techniques", Spatial Information Research, 2019

Publication

<1 %

11

Saidy, A.R., R.J. Smernik, J.A. Baldock, K. Kaiser, and J. Sanderman. "The sorption of organic carbon onto differing clay minerals in the presence and absence of hydrous iron oxide", *Geoderma*, 2013.

Publication

<1 %

12

sutir.sut.ac.th:8080

Internet Source

<1 %

13

Hu, Qiong, Wenbin Wu, Tian Xia, Qiangyi Yu, Peng Yang, Zhengguo Li, and Qian Song. "Exploring the Use of Google Earth Imagery and Object-Based Methods in Land Use/Cover Mapping", *Remote Sensing*, 2013.

Publication

<1 %

14

Israa Kadhim, Fanar Abed. "The Potential of LiDAR and UAV-Photogrammetric Data Analysis to Interpret Archaeological Sites: A Case Study of Chun Castle in South-West England", *ISPRS International Journal of Geo-Information*, 2021

Publication

<1 %

15

Xiaoyu Song, Chenghai Yang, Mingquan Wu, Chunjiang Zhao, Guijun Yang, Wesley Hoffmann, Wenjiang Huang. "Evaluation of Sentinel-2A Satellite Imagery for Mapping Cotton Root Rot", *Remote Sensing*, 2017

Publication

<1 %

16

Jiyao Zhao, Guangzhao Chen, Le Yu, Chao Ren, Jing Xie, Lamuel Chung, Hao Ni, Peng Gong. "Mapping urban morphology changes in the last two decades based on local climate zone scheme: A case study of three major urban agglomerations in China", *Urban Climate*, 2023

Publication

<1 %

17

dspace.ut.ee

Internet Source

<1 %

18

scirp.org

Internet Source

<1 %

19

www.eemj.icpm.tuiasi.ro

Internet Source

<1 %

20

www.iirs-nrsa.gov.in

Internet Source

<1 %

21

Bhagawat Rimal, Roshan Sharma, Ripu Kunwar, Hamidreza Keshtkar, Nigel E. Stork, Sushila Rijal, Syed Ajijur Rahman, Himlal Baral. "Effects of land use and land cover change on ecosystem services in the Koshi River Basin, Eastern Nepal", *Ecosystem Services*, 2019

Publication

<1 %

22

George P. Petropoulos, Dionissios P. Kalivas, Iro A. Georgopoulou, Prashant K. Srivastava. "Urban vegetation cover extraction from

<1 %

hyperspectral imagery and geographic information system spatial analysis techniques: case of Athens, Greece", Journal of Applied Remote Sensing, 2015

Publication

23

Matthew L. Clark. "Comparison of multi-seasonal Landsat 8, Sentinel-2 and hyperspectral images for mapping forest alliances in Northern California", ISPRS Journal of Photogrammetry and Remote Sensing, 2020

Publication

<1 %

24

psasir.upm.edu.my

Internet Source

<1 %

25

tud.qucosa.de

Internet Source

<1 %

26

Abebe Mohammed Ali, Roshanak Darvishzadeh, Andrew Skidmore, Tawanda W. Gara et al. "Comparing methods for mapping canopy chlorophyll content in a mixed mountain forest using Sentinel-2 data", International Journal of Applied Earth Observation and Geoinformation, 2020

Publication

<1 %

27

Caroline M. Gevaert. "Explainable AI for earth observation: A review including societal and regulatory perspectives", International Journal of Applied Earth Observation and

<1 %

28

George P. Petropoulos, Charalambos C. Kontoes, Iphigenia Keramitsoglou. "Land cover mapping with emphasis to burnt area delineation using co-orbital ALI and Landsat TM imagery", International Journal of Applied Earth Observation and Geoinformation, 2012

Publication

<1 %

29

Isaac Larbi. "Land use-land cover change in the Tano basin, Ghana and the implications on sustainable development goals", Heliyon, 2023

Publication

<1 %

30

Springer Remote Sensing/Photogrammetry, 2015.

Publication

<1 %

31

Terry Idol, Barry Haack, Ron Mahabir. "An evaluation of Radarsat-2 individual and combined image dates for land use/cover mapping", Geocarto International, 2015

Publication

<1 %

32

ijeab.com

Internet Source

<1 %

33

research.edgehill.ac.uk

Internet Source

<1 %

34

scholars.unh.edu

<1 %

35

"Proceedings of International Conference on Remote Sensing for Disaster Management", Springer Science and Business Media LLC, 2019

Publication

<1 %

36

Alejandra Aurelia López-Caloca, Amilcar Morales Gamas, María Gabriela López Aguilar. "Wetland landscape based on Sentinel-2 images and geo-tagged photographs in Centla, Tabasco", Terra Digitalis, 2021

Publication

<1 %

37

Bhagawat Rimal, Lifu Zhang, Hamidreza Keshtkar, Barry Haack, Sushila Rijal, Peng Zhang. "Land Use/Land Cover Dynamics and Modeling of Urban Land Expansion by the Integration of Cellular Automata and Markov Chain", ISPRS International Journal of Geo-Information, 2018

Publication

<1 %

38

Linking People Place and Policy, 2002.

Publication

<1 %

39

Vikas Kumar Rana, Tallavajhala Maruthi Venkata Suryanarayana. "Performance evaluation of MLE, RF and SVM classification algorithms for watershed scale land use/land

<1 %

cover mapping using sentinel 2 bands",
Remote Sensing Applications: Society and
Environment, 2020

Publication

40 africanremotesensing.org <1 %
Internet Source

41 opencommons.uconn.edu <1 %
Internet Source

42 perakgateway.utp.edu.my <1 %
Internet Source

43 repository.up.ac.za <1 %
Internet Source

44 research.library.mun.ca <1 %
Internet Source

45 www.mcser.org <1 %
Internet Source

46 www.suaire.sua.ac.tz <1 %
Internet Source

47 Comber, Alexis, Peter Fisher, Chris Brunsdon,
and Abdulhakim Khmag. "Spatial analysis of
remote sensing image classification
accuracy", Remote Sensing of Environment,
2012.
Publication

48 Doni Rahmat Wicakso, Sutijan, Rochmadi,
Arief Budiman. "Study of catalytic upgrading
<1 %

of biomass tars using Indonesian iron ore",
AIP Publishing, 2017

Publication

49

Foody, G.M.. "Toward intelligent training of supervised image classifications: directing training data acquisition for SVM classification", Remote Sensing of Environment, 20041030

Publication

<1 %

50

Nguyen-Thanh Son, Bui-Xuan Thanh. "Decadal assessment of urban sprawl and its effects on local temperature using Landsat data in Cantho city, Vietnam", Sustainable Cities and Society, 2018

Publication

<1 %

51

Nubia da Silva, Manuel Eduardo Ferreira, Cátia Nunes da Cunha, Gustavo Manzon Nunes. "Object-Based Classification of the veredas wetland macrohabitat using multispectral imagery from a Remotely Piloted Aircraft System.", Research Square Platform LLC, 2022

Publication

<1 %

52

agritrop.cirad.fr

Internet Source

<1 %

53

citeseerx.ist.psu.edu

Internet Source

<1 %

54	core.ac.uk Internet Source	<1 %
55	dergipark.org.tr Internet Source	<1 %
56	eprints.utas.edu.au Internet Source	<1 %
57	ir.canterbury.ac.nz Internet Source	<1 %
58	lup.lub.lu.se Internet Source	<1 %
59	research.vu.nl Internet Source	<1 %
60	theses.ncl.ac.uk Internet Source	<1 %
61	ufdcimages.uflib.ufl.edu Internet Source	<1 %
62	waushara.extension.wisc.edu Internet Source	<1 %
63	www.scirp.org Internet Source	<1 %
64	Subhankar Das. "A New approach to assessing the accuracy of the LULC classification using Remote Sensing & GIS in	<1 %

65

Özlem Akar. "Mapping land use with using Rotation Forest algorithm from UAV images", European Journal of Remote Sensing, 2017

Publication

<1 %

66

Du, Peijun, Alim Samat, Björn Waske, Sicong Liu, and Zhenhong Li. "Random Forest and Rotation Forest for fully polarized SAR image classification using polarimetric and spatial features", ISPRS Journal of Photogrammetry and Remote Sensing, 2015.

Publication

<1 %

67

Ihsan Noor, Yudi Firmanul Arifin, Bambang Joko Priatmadi, Akhmad Rizalli Saidy. "Laboratory simulation of the swampy forest system for the passive treatment of acid mine drainage in coal mine reclamation areas", Scientific Reports, 2023

Publication

<1 %

68

Ilina Kamenova, Petar Dimitrov. "Evaluation of Sentinel-2 vegetation indices for prediction of LAI, fAPAR and fCover of winter wheat in Bulgaria", European Journal of Remote Sensing, 2020

Publication

<1 %

69

Laura Piedadlobo, David Hernández-López, Rocío Ballesteros, Amal Chakhar et al.

"Scalable pixel-based crop classification combining Sentinel-2 and Landsat-8 data time series: Case study of the Duero river basin", *Agricultural Systems*, 2019

Publication

<1 %

70

Martin Schmieder, Felix Holl, Marina L. Fotteler, Michael Ortl, Elmar Buchner, Walter Swoboda. "Remote sensing and on-site characterization of wetlands as potential habitats for malaria vectors – A pilot study in southern Germany", 2020 IEEE Global Humanitarian Technology Conference (GHTC), 2020

Publication

<1 %

71

Robert Chastain, Ian Housman, Joshua Goldstein, Mark Finco. "Empirical cross sensor comparison of Sentinel-2A and 2B MSI, Landsat-8 OLI, and Landsat-7 ETM+ top of atmosphere spectral characteristics over the conterminous United States", *Remote Sensing of Environment*, 2019

Publication

<1 %

72

Yisa Ginath Yuh, Wiktor Tracz, H. Damon Matthews, Sarah E. Turner. "Application of machine learning approaches for land cover

<1 %

monitoring in northern Cameroon", Ecological Informatics, 2022

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

2021-8 Syarifuddin K Jurnal Internasional _ARInt Land-CLASSIFIERS FOR LAND USE Juni 2021

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14
