



Re: Manuscript Submission

4 messages

IJDDT Journal <ijddtjournal@gmail.com>
To: "esuhartono@ulm.ac.id" <esuhartono@ulm.ac.id>

Sun, Mar 6, 2022 at 1:54 PM

The manuscript is provided with number 03062022DDTN

On Fri, Jan 28, 2022 at 5:02 AM esuhartono@ulm.ac.id <esuhartono@ulm.ac.id> wrote:**1. Name of Corresponding Author**

Eko Suhartono

2. Email of Corresponding Authoresuhartono@ulm.ac.id**Institute of Affiliation of Corresponding Author**1) Department of Medical Chemistry/Biochemistry, Faculty of Medicine, Lambung Mangkurat University,
Banjarmasin, Indonesia**5. Title of Manuscript**PROTECTIVE ROLE OF KELAKAI (*Stenochlaena palustris*) EXTRACT ON MALATHION-INDUCED GENOTOXIC:
FTIR SPECTROSCOPY STUDY**6. Abstract**

Malathion is a genotoxic pesticide that destructs DNA. Many researches have proven it, yet the mechanism has not been clear. That is the reason why this research needs to be conducted. This research used two groups, control, and two treatment group. The control group (P0) was 100 μ L of Human Genomic DNA Female solution, while the treatment group (P1) was 100 μ L of Human Genomic DNA Female solution which was added with 100 μ L of 6 mM malathion solution. The other treatment group (P2) was 100 μ L of Human Genomic DNA Female solution which was added with 100 μ L of 6 mM malathion solution. Each group of solutions was incubated at 37oC for 48 hours. Then the absorbance was determined by FTIR. The content of the kelakai extract was determined by GC-MS. The results of the GC-MS analysis showed that the ethanolic extract of kelakai contained 6.1% hexadecanoic acid and 6.54% Neophytadiene. the percentage changes in the absorbance of guanine, thymine, cytosin, and adenine were significantly different in each group. So as deformation of the NH-groups in DNA bases. This means that kelakai extract can inhibit DNA damage.

7. Keywords (2-10)

Malathion, Genotoxic, Kelakai, hexadecenoic acid, neophytadiene

8. Complete manuscript in one file (word)

- [Publikasi_IJDDT_Fix.docx](#)

--
Editor
International Journal of Drug Delivery Technology
ISSN: 0975-4415
website: www.ijddt.com

IJDDT Journal <ijddtjournal@gmail.com>
To: "esuhartono@ulm.ac.id" <esuhartono@ulm.ac.id>

Tue, Mar 15, 2022 at 8:14 PM

Dear author

Your manuscript number " 03062022DDTN " bearing with title " PROTECTIVE ROLE OF KELAKAI (*Stenochlaena palustris*) EXTRACT ON MALATHION-INDUCED GENOTOXIC: FTIR SPECTROSCOPY STUDY " has been

provisionally accepted after minor modifications. It may be published in the upcoming issue of the journal subject to deposition of Open Access Fee USD200 to ijddtjournal@gmail.com
Refer this link to pay <https://ijddt.com/fee/>

Note: The authors may **reply this mail** with modified manuscript containing minor typography errors if left coincidentally during submission. The errors must be highlighted clearly in **red color font**. The modifications are subject to approval of editorial board. Any typological/coincidental error left after this stage will not be considered. In case the modifications are not minor, then authors may be asked to withdraw and resubmit manuscript.

Important: After making a transaction, please confirm transaction by providing transaction details as a reply to this mail only.

--
Publisher

[Quoted text hidden]

Eko Suhartono <esuhartono@ulm.ac.id>
To: IJDDT Journal <ijddtjournal@gmail.com>


Wed, Mar 16, 2022 at 3:52 PM

Dear editorial team

We send proof of payment and articles that have been revised.
Thank You

Best Regards
Eko Suhartono
[Quoted text hidden]

2 attachments



WhatsApp Image 2022-03-16 at 15.38.45.jpeg
82K

 **article_Eko Suhartono_IJDDT.docx**
181K

IJDDT Journal <ijddtjournal@gmail.com>
To: Eko Suhartono <esuhartono@ulm.ac.id>

Thu, Mar 17, 2022 at 2:12 PM

publication charges received
[Quoted text hidden]

Protective Role of Kelakai (*Stenochlaena Palustris*) Extract on Malathion-induced Genotoxic: FTIR Spectroscopy Study

Q2 Eko Suhartono^{1,2*}, Noor Muthmainah³, Dona Marisa⁴, Salmon Charles P T Siahaan⁵,
Noer Komari⁶

Q3 ¹Department of Medical Chemistry, Biochemistry, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

²Wetland Disease Center, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

³Department of Microbiology and Parasitology, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

⁴Department of Biomedic, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

⁵Faculty of Medicine, University of Ciputra, Surabaya, Indonesia

⁶Department of Chemistry, Faculty of Mathematics and Natural Sciences, Lambung Mangkurat University, Banjarmasin, Indonesia

Received: 28th January, 2022; Revised: 01st March, 2021; Accepted: 17th March, 2022; Available Online: 25th March, 2022

ABSTRACT

Malathion is a genotoxic pesticide that destructs DNA. Many research have proven it, yet the mechanism has not been clear. That is the reason why this research needs to be conducted. This research used two groups, control, and two treatment group. The control group (P0) was 100 µL of Human Genomic DNA Female solution, while the treatment group (P1) was 100 µL of Human Genomic DNA Female solution which was added with 100 µL of 6 mM malathion solution. The other treatment group (P2) was 100 µL of Human Genomic DNA Female solution which was added with 100 µL of 6 mM malathion solution. Each group of solutions was incubated at 37°C for 48 hours. Then the absorbance was determined by FTIR. The content of the kelakai extract was determined by GC-MS. The results of the GC-MS analysis showed that the ethanolic extract of kelakai contained 6.1% hexadecanoic acid and 6.54% Neophytadiene. the percentage changes in the absorbance of guanine, thymine, cytosin, and adenine were significantly different in each group. So as deformation of the NH-groups in DNA bases. This means that kelakai extract can inhibit DNA damage.

Keywords: Genotoxic, Hexadecenoic acid, Kelakai, Neophytadiene, Malathion.

International Journal of Drug Delivery Technology (2022); DOI: 10.25258/ijddt.12.1.00

How to cite this article: Suhartono E, Muthmainah N, Marisa D, Siahaan SCPT, Komari N. Protective Role of Kelakai (*Stenochlaena Palustris*) Extract on Malathion-induced Genotoxic: FTIR Spectroscopy Study. International Journal of Drug Delivery Technology. 2022;12(1):1-4.

Source of support: Nil.

Conflict of interest: None

INTRODUCTION

Malathion is an organophosphate pesticide that is widely used by farmers as pest control. On the other hand, malathion is also a cancer-causing toxic substance. Previous research stated that malathion is a type of pesticide that triggers ovarian cancer and breast cancer.¹ Research by El Baz *et al.*² also revealed that malathion can trigger leukemia in the children of farmers who use the malathion pesticide.

The mechanism of malathion as a carcinogen is not fully known, but many studies have linked it to the DNA methylation process. DNA methylation can interfere with the transcription process so that gene expression is disrupted.³ In addition, the

nucleophilic group in malathion can cause the breaking of phosphodiester bonds in DNA.^{4,5}

Several studies have revealed that hexadecanoic acid compounds can prevent proliferation by inhibiting DNA-Topoisemerase-1.⁶ Research by Bharath *et al*⁷ also stated that hexadecanoic acid derived from the algae *Turbinaria ornata* has the potential as an anticancer of the colon in vitro. Nurmalatina's research⁸ states that hexadecanoic acid compounds can also be found in kelakai (*Stenochlaena palustris*).

Kelakai is a type of ferns, belonging to the pteridaceae family that grows in areas of high humidity such as peatlands in

*Author for Correspondence: esuhartono@ulm.ac.id

Kalimantan. Kelakai is consumed by the people in Kalimantan as a traditional food and is empirically useful in preventing anemia, stimulating breast milk production, and treating wounds^{9,10}. In addition, at a dose of 100 mg/kg, Kelakai acts as immunomodulators, which modulate the production of IL-10 cytokines in *P. berghei* infection and suppress the production of IFN- γ , IL-6, TNF- α by T cells.¹¹ Kelakai also has a cytotoxic effect on T47D breast cancer cells by increasing the expression of the p53 gene, but the mechanism is still unclear.¹² Therefore, in this study, it is necessary to study the Malathion-DNA interaction and the role of malathion extract in inhibiting DNA damage due to malathion.

MATERIAL AND METHODS

Material

The Kelakai were taken from a peat swamp in Banjarbaru, South Kalimantan, and the young leaves were selected. The leaves are dried for maceration. Calf Thymus DNA was obtained from SigmaAldrich, St. Louis, MO, USA to examine its interaction with malathion (SigmaAldrich, St. Louis, MO, USA) and Kelakai extract.

Measurement by GC-MS

The leaves are washed and then dried under the sun to air dry for 2–3 days. After drying, 250 g were ground and then extracted by maceration with ethanol as a solvent. Maceration was carried out for 3 days. The extraction results were filtered and evaporated using a rotary evaporator and then measured by GC-MS at the Banjarbaru Industrial and Trade Research Institute, Indonesia.

Experiment

This research used two groups, control, and two treatment group. The control group (P0) was 100 μ L of Human Genomic DNA Female solution, while the treatment group (P1) was 100 μ L of Human Genomic DNA Female solution which was added with 100 μ L of 6 mM malathion solution. The other treatment group (P2) was 100 μ L of Human Genomic DNA Female solution which was added with 100 μ L of 6 mM malathion solution. Each group of solutions was incubated at 37°C for 48 hours.

Measurement Absorbance by ATR-FTIR

The solutions contained in the control and treatment groups were each taken by 20 μ L and their absorbance measured by ATR-FTIR spectroscopy using ATR crystalline silicon (top plate 45°C) from BioATR Cell II.^{13,14}

Data Analysis

Observation, that was focussed to DNA vibration ribbon at 1710, 1662, 1613, and 1492 cm^{-1} , showed the existence of guanine (G), thymine (T), adenine (A), dan cytosine nitrogen (C) for each. In addition, in the ribbon 1228 and 1087 cm^{-1} each showed asymmetrical and symmetrical phosphate vibrations.¹⁵ In addition, DNA oxidation was observed at wave number 1050 cm^{-1} , while DNA methylation was observed at 1596 cm^{-1} . NH deformation was observed at wave numbers 1288, 1294 and 1684 cm^{-1} .¹⁵ Data

obtained were then compared between the control and treatment groups using the Kruskal Wallis test at $\alpha = 0.05$.

RESULT

The results of the GC-MS analysis showed that the ethanolic extract of kelakai contained 6.1% hexadecanoic acid and 6.54% Neophytadiene (Figure 1).

Spectroscopy is the study of the interaction of light with atoms and molecules. This interaction can be used to assess the strength of the bonds between molecules by calculating the value of the binding constant. The results of the interaction between DNA-Malathion and DNA-Malathion-extract kelakai are presented in Figure 2.

Based on Figure 1, there is a change in the absorbance spectra between DNA and DNA+Malathion. The presence of an extract of malachite showed that there was DNA protection from Malathion. The percentage change in absorbance of each group can be seen in Table 1.

DISCUSSION

Based on data from the World Health Organization (WHO) in 2014, pesticide toxication reached one million cases in the world, with 20,000 death per year. Malathion toxication as pesticide started when malathion enters the body and inhibits the metabolism of acetylcholinesterase (AchE), cytochrome





Figure 1: Hexadecanoic Acid (a) dan Neophytadiene (b)



①②③④⑤⑥⑦⑧⑨⑩

Figure 2: Spectra of FTIR DNA

Table 1: The percentage change in DNA base

Base	Wavenumbers	Absorbance change in FTIR (%)		
		P0	P1	P2
Guanine	1710	100	61,75 ± 1,21	79,99 ± 2,39*
Thymine	1992	100	56,96 ± 1,19	79,99 ± 1,97*
Adenine	1613	100	58,71 ± 1,44	79,99 ± 2,04*
Cytocin,	1492	100	60,74 ± 1,38	80,00 ± 2,72*

Note: *) significantly different ($p < 0,05$)

Changes in DNA bases can be caused by deformation of the NH-groups in DNA. This can be seen in table 2.

Table 2: Deformation of the NH-groups in DNA bases

Base	Wavenumbers	Absorbance change in FTIR (%)		
		P0	P1	P2
Guanine	1684	100	59,43 ± 3,02	80,00 ± 2,87*
Thymine	1288	100	62,31 ± 2,66	80,00 ± 2,91*
Adenine	-	-	-	-
Cytosine	1294	100	62,04 ± 2,98	80,00 ± 2,07*

Note: *) significantly different ($p < 0,05$)

P450, and glutathione S-transferases (GSTs).^{16,17} Next, the inhibition caused the increase of reactive oxygen species (ROS) production, the decrease of enzymatic antioxidant activity, induce apoptosis (cytotoxicity), and DNA damage.^{18,19}

FT-IR is one of the many spectroscopic methods used in the study of DNA with pollutants, drugs, and other molecules in solution. This method is ideal for systematic DNA studies because it requires small samples. The region of 1800–800 cm^{-1} FT-IR spectroscopy is a characteristic peak of infrared absorption from the free located B-form DNA. The ribbon at 1710 cm^{-1} shows stretching carbonyl (C=O). This is associated with guanine (G) C7N in vibration stretching fields. The tape observed at 1658 cm^{-1} shows a stretched carbonyl. This is associated with stretching thymine (T) C2O. The ribbon 1613 cm^{-1} occurred due to scissoring NH2. This is associated with adenine (A). Bands at 1492 cm^{-1} correspond to cytosine (C) base vibrations¹⁵ [26]. The bands at 1228 and 1087 cm^{-1} show asymmetric and symmetric vibrations of phosphate, respectively¹⁵ [26]. The absorption band in the spectral region of 1750–1450 cm^{-1} is assigned to stretch the vibrations of nucleotide base bonds in the field and is generally used for evaluation of pair-base interactions.²⁰

DNA damage because of malathion has been studied through FTIR spectroscopy. In table 1, there is an absorbance decrease for DNA significantly between the Calf-Thymus DNA group and the Calf-Thymus DNA+Malathion group. This shows that malathion interacts with DNA through thymine, adenine, guanine, and cytosine. In addition, the absorbance value for phosphorus in Calf-Thymus DNA significantly increased. The phosphorus group in methyl paration is a good substrate for nucleophilic attack. This attack causes an increase in phosphorylation, which can cause DNA damage.¹⁶

This interaction of malathion and DNA can also be caused by nitrogen bases which have free electron pairs. This electron pair allows the rearrangement of nitrogen by the presence of

metilparation. This causes an increase in NH-deformation in nitrogen bases DNA base. This study is in accordance with the research of Yu Li *et al.*,²¹ who explained that the chlorpropham herbicide can bind specifically to -NH in DNA so that the deformation –NH increases.

The research found that the absorbance of DNA + malathion oxidation increased by almost 250x. This oxidation begins with the reaction between the nitrogenous base of DNA and methyl parathion, which causes changes in hydrogen bonds in DNA bases. These changes result in deformation of the NH-groups and other groups, so that oxo compounds can be formed from DNA bases.

Kelakai in a wetlands endemic plant that can detain DNA damage. It contains neophytadiene which has the potential to prevent DNA damage due to malathion. Neophytadiene works by binding to DNA, between C-17 in hexadecanoic acid and dG-12A in DNA, so malathion cannot interact with DNA. In addition, the atomic contact energy of DNA-Malathion-hexadecanoic acid is greater than that of DNA-Malathion so that the DNA-Malathion bonds are more easily broken. The area of contact between DNA-Malathion-hexadecanoic acid atoms is larger than the others. The larger the contact area, the faster the reaction will be than the others.

Besides the physical aspects, there is also chemical aspect, which is the presence of hydrogen bonding. Hydrogen bonding occurs due to the intermolecular attractive forces that occur between hydrogen atoms with highly electronegative atoms (N, O, and F) and lone pairs of electrons from other highly electronegative atoms. This bonding can occur in DNA, through N-H and O-H bonds with a partial positive charge on the H and a partial negative charge on the N and O atoms.

CONCLUSION

FTIR spectroscopy can determine the interaction of DNA-Malathion and DNA-Malathion-Kelakai Extract. Kelakai

Q8

Q6

Q6

Q7

extract contains hexadecanoic acid and neophytadiene which can inhibit DNA damage due to malathion induced.

ACKNOWLEDGEMENTS

The authors thankfully acknowledge the Rector of Lambung Mangkurat University for providing research funding, through the PDWM program, in2021.

REFERENCES

- Lerro CC, *et al*. Organophosphate insecticide use and cancer incidence among spouses of pesticide applicators in the Agricultural Health Study. *Occup Environ Med* 2015;72:736–744
- El Baz MA, El Deek SE, Sayed AA, Amin AF. In Utero Pesticides Exposure and Generation of Acute Myeloid Leukemia Associated Translocation (8; 21). *MOJ Toxicol* 2016; 2(3): 00037.
- Zhang X, Wallace AD, Du P, Kibbe WA, Jafari N, Xie H, *et al*. DNA methylation and alterations in response to pesticide exposure in vitro. *Environmental and Molecular Mutagenesis* 2012; 53:542-9.
- Mathew J, dan Thoppil JE. Genotoxicity of methyl parathion and antimutagenic activity of salvia officinalis L. (sage) extracts in swiss albino mice. *Asian Journal of Pharmaceutical and Clinical Research* 2012; 5(2):164-70.
- Camden AJ, Walsh SM, Suk SH, Silverman SK. DNA oligonucleotide 3'-phosphorilation by a DNA enzyme. *Biochemistry* 2016; 55:2671-6.
- Ravi L and Krishnan K. Cytotoxic Potential of N-hexadecanoic Acid Extracted from *Kigelia pinnata* Leaves. *Asian J. Cell Biol* 2017; 12 (1): 20-27
- Bharath B, Perinbam K, Devanesan B, AlSalhi M, Saravanan M. Evaluation of the anticancer potential of Hexadecanoic acid from brown algae *Turbinaria ornata* on HT–29 colon cancer cells. *Journal of Molecular Structure* 2021; 1235: 130229
- Nurmilatina, N. Analisis Komposisi Kimia Daun Kelakai (*Stenochlaena palustris* Bedd.) dengan Berbagai Pelarut menggunakan GCMS (Chemical Composition Analysis of *Stenochlaena palustris* Bedd. Leaves using Various Solvents on GCMS). *Jurnal Riset Industri Hasil Hutan* 2017; 9(1): 9 – 16
- Suhartono E, Viani E, Rahmadhan MA, Gultom IS, Rakhman MF, and Indrawardhana D. Screening of Medicinal Plant for Total Flavonoid and Antioxidant Activity in South Kalimantan of Indonesian. *International Journal of Chemical Engineering and Applications* 2012. 3(4): 297-299
- Suhartono E, Bahriansyah M, Triawanti. The Inhibition Effect of Kelakai (*Stenochlaena palustris*) Extract on Cadmium-Induced Glycation and Fructation *In-vitro*. *International Journal of Pharmaceutical and Clinical Research* 2016; 8(4): 248-253
- Margono DPNH, Suhartono E, Arwati H. Potensi Ekstrak Kelakai (*Stenochlaena Palustris* (Burm.F) Bedd) Terhadap Kadar Tumor Necrosis Factor-Alfa (Tnf-A) Pada Mencit Balb/C Yang Diinfeksi Plasmodium berghei ANKA. *Berkala Kedokteran* 2016; 12(1): 77-85
- Safitri RA, Saptarini, Sunarni T. Cytotoxic Activity, Expression of p53, and Bcl-2 Extract Fraction of Kelakai Herbs (*Stenochlaena palustris* (Burm.F.) Bedd.) to Breast Cancer T47D Cells. *Jurnal Biotek Medisiana Indonesia* 2020; 9(2): 113-127
- Suhartono E. Noor Z, Edyson; Budianto WY, Idroes R. Effect of chronic lead exposure on bone using ATR-FTIR spectroscopy. In *Proceedings of the AIP Conference Proceedings* 2019; p. 020025.
- Al-Jorani K. R  ther A, Martin M, Haputhanthri R, Deacon G, Li H, Wood B. The Application of ATR-FTIR Spectroscopy and the Reversible DNA Conformation as a Sensor to Test the Effectiveness of Platinum (II) Anticancer Drugs. *Sensors* 2018; 18: 4297.
- Movasaghi Z, Rehman S, Ihtehsham, Fourier Transform (FTIR) Spectroscopy of Biological Tissues. *Applied Spectroscopy Review* 2008, 43: 134-179
- Bianchi J, Mantovani MS, Morales MAM. Analysis of the genotoxic potential of low concentrations of Malathion on the *Allium Cepapa* cells and rat hepatoma tissue culture. *Journal of Environmental Sciences*. 2015; 36:102-11.
- Chear, N. J. Y., Khaw, K. Y., Murugaiyah, V., & Lai, C. S. Cholinesterase inhibitory activity and chemical constituents of *Stenochlaena palustris* fronds at two different stages of maturity. *Journal of Food and Drug Analysis* 2016; 24: 358-366
- Garcia FP, Ascencio SYC, Oyarzun JCG, Hernandez AC, Alavarado PV. Pesticides : classification, uses, and toxicity. Measures of exposure and genotoxic risks. *J Res. Environ. Sci. Toxicol* 2012; 1(11):279-93.
- Ojha A, Yaduvanshi SK, Pant SC, Lomash V, Srivastava N. Evaluation of DNA damage and cytotoxicity induced by three commonly used organophosphate pesticides individually and in mixture, in rat tissues *Environ. Toxicol* 2013; 28: 543–552.
- Colovic, MB. Krstic DZ, Lazarevic-Pasti, T.D, Bondzic, A.M, Vasic, VM. Acetylcholinesterase inhibitors: pharmacology and toxicology. *Curr. Neuropharmacol* 2013; 11: 315–335.
- Yu Li, Guowen Zhang, Mo Tao. 2014. Binding properties herbicide chlorpropham to DNA: Spectroscopy, Chemometrics, and modelling investigations, *Journal Photochemistry and Photobiology B: Biology* 2014; 138:109-117

Query Report

Q1 Article Type is missing

Q2 Only 1 affiliation will be considered. Plz confirm which one must be written here

Q3 Plz share the department name again

Q4 Short title of the Article is missing

Q5 Provide full name

Q6 Reference 15 or 26?

Q7 Plz correct the word

Q8 Cross-check for accuracy/data/format, correct, and provide all references in consistent Vancouver format.

Sample reference formatted to act as an example and to maintain consistency of format (Vancouver) required

Kumar RA, Sridevi K, Kumar NV, Nanduri S, Rajagopal S. Anticancer and immunostimulatory compounds from *Andrographis paniculata*. *Journal of Ethnopharmacology*. 2004;92:291-295. Available from: doi.org/10.1016/j.jep.2004.03.004

Protective Role of Kelakai (*Stenochlaena Palustris*) Extract on Malathion-induced Genotoxic: FTIR Spectroscopy Study

Q2 Eko Suhartono¹, Noor Muthmainah³, Dona Marisa⁴, Salmon Charles P T Siahaan⁵,
Noer Komari⁶

Q3 ¹Department of Medical Chemistry, Biochemistry, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

²Wetland Disease Center, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

³Department of Microbiology and Parasitology, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

⁴Department of Biomedic, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia

⁵Faculty of Medicine, University of Ciputra, Surabaya, Indonesia

⁶Department of Chemistry, Faculty of Mathematics and Natural Sciences, Lambung Mangkurat University, Banjarmasin, Indonesia

Received: 28th January, 2022; Revised: 01st March, 2021; Accepted: 17th March, 2022; Available Online: 25th March, 2022

ABSTRACT

Malathion is a genotoxic pesticide that destructs DNA. Many research have proven it, yet the mechanism has not been clear. That is the reason why this research needs to be conducted. This research used two groups, control, and two treatment group. The control group (P0) was 100 μ L of Human Genomic DNA Female solution, while the treatment group (P1) was 100 μ L of Human Genomic DNA Female solution which was added with 100 μ L of 6 mM malathion solution. The other treatment group (P2) was 100 μ L of Human Genomic DNA Female solution which was added with 100 μ L of 6 mM malathion solution. Each group of solutions was incubated at 37°C for 48 hours. Then the absorbance was determined by FTIR. The content of the kelakai extract was determined by GC-MS. The results of the GC-MS analysis showed that the ethanolic extract of kelakai contained 6.1% hexadecanoic acid and 6.54% Neophytadiene. the percentage changes in the absorbance of guanine, thymine, cytosin, and adenine were significantly different in each group. So as deformation of the NH-groups in DNA bases. This means that kelakai extract can inhibit DNA damage.

Keywords: Genotoxic, Hexadecenoic acid, Kelakai, Neophytadiene, Malathion.

International Journal of Drug Delivery Technology (2022); DOI: 10.25258/ijddt.12.1.00

How to cite this article: Suhartono E, Muthmainah N, Marisa D, Siahaan SCPT, Komari N. Protective Role of Kelakai (*Stenochlaena Palustris*) Extract on Malathion-induced Genotoxic: FTIR Spectroscopy Study. International Journal of Drug Delivery Technology. 2022;12(1):1-4.

Source of support: Nil.

Conflict of interest: None

INTRODUCTION

Malathion is an organophosphate pesticide that is widely used by farmers as pest control. On the other hand, malathion is also a cancer-causing toxic substance. Previous research stated that malathion is a type of pesticide that triggers ovarian cancer and breast cancer.¹ Research by El Baz *et al.*² also revealed that malathion can trigger leukemia in the children of farmers who use the malathion pesticide.

The mechanism of malathion as a carcinogen is not fully known, but many studies have linked it to the DNA methylation process. DNA methylation can interfere with the transcription process so that gene expression is disrupted.³ In addition, the

nucleophilic group in malathion can cause the breaking of phosphodiester bonds in DNA.^{4,5}

Several studies have revealed that hexadecanoic acid compounds can prevent proliferation by inhibiting DNA-Topoisemerase-1.⁶ Research by Bharath *et al*⁷ also stated that hexadecanoic acid derived from the algae *Turbinaria ornata* has the potential as an anticancer of the colon in vitro. Nurmalatina's research⁸ states that hexadecanoic acid compounds can also be found in kelakai (*Stenochlaena palustris*).

Kelakai is a type of ferns, belonging to the pteridaceae family that grows in areas of high humidity such as peatlands in

Kalimantan. Kelakai is consumed by the people in Kalimantan as a traditional food and is empirically useful in preventing anemia, stimulating breast milk production, and treating wounds^{9,10}. In addition, at a dose of 100 mg/kg, Kelakai acts as immunomodulators, which modulate the production of IL-10 cytokines in *P. berghei* infection and suppress the production of IFN- γ , IL-6, and TNF- α by T cells.¹¹ Kelakai also has a cytotoxic effect on T47D breast cancer cells by increasing the expression of the p53 gene, but the mechanism is still unclear.¹² Therefore, in this study, it is necessary to study the Malathion-DNA interaction and the role of malathion extract in inhibiting DNA damage due to malathion.

MATERIAL AND METHODS

Material

The Kelakai were taken from a peat swamp in Banjarbaru, South Kalimantan, and the young leaves were selected. The leaves are dried for maceration. Calf Thymus DNA was obtained from SigmaAldrich, St. Louis, MO, USA to examine its interaction with malathion (SigmaAldrich, St. Louis, MO, USA) and Kelakai extract.

Measurement by GC-MS

The leaves are washed and then dried under the sun to air dry for 2–3 days. After drying, 250 g were ground and then extracted by maceration with ethanol as a solvent. Maceration was carried out for 3 days. The extraction results were filtered and evaporated using a rotary evaporator and then measured by GC-MS at the Banjarbaru Industrial and Trade Research Institute, Indonesia.

Experiment

This research used two groups, control, and two treatment group. The control group (P0) was 100 μ L of Human Genomic DNA Female solution, while the treatment group (P1) was 100 μ L of Human Genomic DNA Female solution which was added with 100 μ L of 6 mM malathion solution. The other treatment group (P2) was 100 μ L of Human Genomic DNA Female solution which was added with 100 μ L of 6 mM malathion solution. Each group of solutions was incubated at 37°C for 48 hours.

Measurement Absorbance by ATR-FTIR

The solutions contained in the control and treatment groups were each taken by 20 μ L and their absorbance measured by ATR-FTIR spectroscopy using ATR crystalline silicon (top plate 45°C) from BioATR Cell II.^{13,14}

Data Analysis

Observation, that was focussed to DNA vibration ribbon at 1710, 1662, 1613, and 1492 cm^{-1} , showed the existence of guanine (G), thymine (T), adenine (A), dan cytosine nitrogen (C) for each. In addition, in the ribbon 1228 and 1087 cm^{-1} each showed asymmetrical and symmetrical phosphate vibrations.¹⁵ In addition, DNA oxidation was observed at wave number 1050 cm^{-1} , while DNA methylation was observed at 1596 cm^{-1} . NH deformation was observed at wave numbers 1288, 1294 and 1684 cm^{-1} .¹⁵ Data

obtained were then compared between the control and treatment groups using the Kruskal Wallis test at $\alpha = 0.05$.

RESULT

The results of the GC-MS analysis showed that the ethanolic extract of kelakai contained 6.1% hexadecanoic acid and 6.54% Neophytadiene (Figure 1).

Spectroscopy is the study of the interaction of light with atoms and molecules. This interaction can be used to assess the strength of the bonds between molecules by calculating the value of the binding constant. The results of the interaction between DNA-Malathion and DNA-Malathion-extract kelakai are presented in Figure 2.

Based on Figure 1, there is a change in the absorbance spectra between DNA and DNA+Malathion. The presence of an extract of malachite showed that there was DNA protection from Malathion. The percentage change in absorbance of each group can be seen in Table 1.

DISCUSSION

Based on data from the World Health Organization (WHO) in 2014, pesticide toxication reached one million cases in the world, with 20,000 death per year. Malathion toxication as pesticide started when malathion enters the body and inhibits the metabolism of acetylcholinesterase (AchE), cytochrome




Figure 1: Hexadecanoic Acid (a) dan Neophytadiene (b)




Figure 2: Spectra of FTIR DNA

Table 1: The percentage change in DNA base

Base	Wavenumbers	Absorbance change in FTIR (%)		
		P0	P1	P2
Guanine	1710	100	61,75 ± 1,21	79,99 ± 2,39*
Thymine	1992	100	56,96 ± 1,19	79,99 ± 1,97*
Adenine	1613	100	58,71 ± 1,44	79,99 ± 2,04*
Cytocin,	1492	100	60,74 ± 1,38	80,00 ± 2,72*

Note: *) significantly different ($p < 0,05$)

Changes in DNA bases can be caused by deformation of the NH-groups in DNA. This can be seen in table 2.

Table 2: Deformation of the NH-groups in DNA bases

Base	Wavenumbers	Absorbance change in FTIR (%)		
		P0	P1	P2
Guanine	1684	100	59,43 ± 3,02	80,00 ± 2,87*
Thymine	1288	100	62,31 ± 2,66	80,00 ± 2,91*
Adenine	-	-	-	-
Cytosine	1294	100	62,04 ± 2,98	80,00 ± 2,07*

Note: *) significantly different ($p < 0,05$)

P450, and glutathione S-transferases (GSTs).^{16,17} Next, the inhibition caused the increase of reactive oxygen species (ROS) production, the decrease of enzymatic antioxidant activity, induce apoptosis (cytotoxicity), and DNA damage.^{18,19}

FT-IR is one of the many spectroscopic methods used in the study of DNA with pollutants, drugs, and other molecules in solution. This method is ideal for systematic DNA studies because it requires small samples. The region of 1800–800 cm^{-1} FT-IR spectroscopy is a characteristic peak of infrared absorption from the free located B-form DNA. The ribbon at 1710 cm^{-1} shows stretching carbonyl (C=O). This is associated with guanine (G) C7N in vibration stretching fields. The tape observed at 1658 cm^{-1} shows a stretched carbonyl. This is associated with stretching thymine (T) C2O. The ribbon 1613 cm^{-1} occurred due to scissoring NH2. This is associated with adenine (A). Bands at 1492 cm^{-1} correspond to cytosine (C) base vibrations.¹⁵ The bands at 1228 and 1087 cm^{-1} show asymmetric and symmetric vibrations of phosphate, respectively.¹⁵ The absorption band in the spectral region of 1750–1450 cm^{-1} is assigned to stretch the vibrations of nucleotide base bonds in the field and is generally used for evaluation of pair-base interactions.²⁰

DNA damage because of malathion has been studied through FTIR spectroscopy. In table 1, there is an absorbance decrease for DNA significantly between the Calf-Thymus DNA group and the Calf-Thymus DNA+Malathion group. This shows that malathion interacts with DNA through thymine, adenine, guanine, and cytosine. In addition, the absorbance value for phosphorus in Calf-Thymus DNA significantly increased. The phosphorus group in methyl parathion is a good substrate for nucleophilic attack. This attack causes an increase in phosphorylation, which can cause DNA damage.¹⁶

This interaction of malathion and DNA can also be caused by nitrogen bases which have free electron pairs. This electron pair allows the rearrangement of nitrogen by the presence of

maliparation. This causes an increase in NH-deformation in nitrogen bases DNA base. This study is in accordance with the research of Yu Li *et al.*,²¹ who explained that the chlorpropham herbicide can bind specifically to -NH in DNA so that the deformation –NH increases.

The research found that the absorbance of DNA + malathion oxidation increased by almost 250x. This oxidation begins with the reaction between the nitrogenous base of DNA and methyl parathion, which causes changes in hydrogen bonds in DNA bases. These changes result in deformation of the NH-groups and other groups, so that oxo compounds can be formed from DNA bases.

Kelakai in a wetlands endemic plant that can detain DNA damage. It contains neophytadiene which has the potential to prevent DNA damage due to malathion. Neophytadiene works by binding to DNA, between C-17 in hexadecanoic acid and dG-12A in DNA, so malathion cannot interact with DNA. In addition, the atomic contact energy of DNA-Malathion-hexadecanoic acid is greater than that of DNA-Malathion so that the DNA-Malathion bonds are more easily broken. The area of contact between DNA-Malathion-hexadecanoic acid atoms is larger than the others. The larger the contact area, the faster the reaction will be than the others.

Besides the physical aspects, there is also chemical aspect, which is the presence of hydrogen bonding. Hydrogen bonding occurs due to the intermolecular attractive forces that occur between hydrogen atoms with highly electronegative atoms (N, O, and F) and lone pairs of electrons from other highly electronegative atoms. This bonding can occur in DNA, through N-H and O-H bonds with a partial positive charge on the H and a partial negative charge on the N and O atoms.

CONCLUSION

FTIR spectroscopy can determine the interaction of DNA-Malathion and DNA-Malathion-Kelakai Extract. Kelakai

extract contains hexadecanoic acid and neophytadiene which can inhibit DNA damage due to malathion induced.

ACKNOWLEDGEMENTS

The authors thankfully acknowledge the Rector of Lambung Mangkurat University for providing research funding, through the PDWM program, in2021.

REFERENCES

1. Lerro CC, et al. Organophosphate insecticide use and cancer incidence among spouses of pesticide applicators in the Agricultural Health Study. *Occup Environ Med* 2015;72:736–744
2. El-Daz MA, El Deek SE, Sayed AA, Amin AF. In Utero Pesticides Exposure and Generation of Acute Myeloid Leukemia Associated Translocation (8; 21). *MOJ Toxicol* 2016; 2(3): 00037.
3. Zhang X, Wallace AD, Du P, Kibbe WA, Jafari N, Xie H, et al. DNA methylation and alterations in response to pesticide exposure in vitro. *Environmental and Molecular Mutagenesis* 2012; 53:542-9.
4. Mathew J, dan Thoppil JE. Genotoxicity of methyl parathion and antimutagenic activity of salvia officinalis L. (sage) extracts in swiss albino mice. *Asian Journal of Pharmaceutical and Clinical Research* 2012; 5(2):164-70.
5. Cunden AJ, Walsh SM, Suk SH, Silverman SK. DNA dependent nucleotide 3'-phosphorilation by a DNA enzyme. *Biochemistry* 2016; 55:2671-6.
6. L and Krishnan K. Cytotoxic Potential of N-hexadecanoic Acid Extracted from *Kigelia pinnata* Leaves. *Asian J. Cell Biol* 2017; 12 (1): 20-27
7. Rath B, Perinbam K, Devanesan B, AlSalhi M, Saravanan M. Evaluation of the anticancer potential of Hexadecanoic acid from brown algae *Turbinaria ornata* on HT–29 colon cancer cells. *Journal of Molecular Structure* 2021; 1235: 130229
8. Milatina, N. Analisis Komposisi Kimia Daun Kelakai (*Stenochlaena palustris* Bedd.) dengan Berbagai Pelarut menggunakan GCMS (Chemical Composition Analysis of *Stenochlaena palustris* Bedd. Leaves using Various Solvents on GCMS). *Jurnal Riset Industri Hasil Hutan* 2017; 9(1): 9 – 16
9. Suhartono E, Viani E, Rahmadhan MA, Gultom IS, Rakhman M, and Indrawardhana D. Screening of Medicinal Plant for Total Flavonoid and Antioxidant Activity in South Kalimantan of Indonesian. *International Journal of Chemical Engineering and Applications* 2012. 3(4): 297-299
10. Suhartono E, Bahriansyah M, Triawanti. The Inhibition Effect of Kelakai (*Stenochlaena palustris*) Extract on Cadmium-Induced Glycation and Fructation *In-vitro*. *International Journal of Pharmaceutical and Clinical Research* 2016; 8(4): 248-253
11. Suhartono E, DPNH, Suhartono E, Arwati H. Potensi Ekstrak Kelakai (*Stenochlaena Palustris* (Burm.F) Bedd) Terhadap Kadar Tumor Necrosis Factor-Alfa (Tnf-A) Pada Mencit Balb/C Yang Diinfeksi Plasmodium berghei ANKA. *Berkala Kedokteran* 2016; 12(1): 77-85
12. Sari RA, Saptarini, Sunarni T. Cytotoxic Activity, Expression of p53, and Bcl-2 Extract Fraction of Kelakai Herbs (*Stenochlaena palustris* (Burm.F.) Bedd.) to Breast Cancer T47D Cells. *Jurnal Biotek Medisiana Indonesia* 2020; 9(2): 113-127
13. Suhartono E, Noor Z, Edyson; Budianto WY, Idroes R. Effect of chronic lead exposure on bone using ATR-FTIR spectroscopy. In *Proceedings of the AIP Conference Proceedings* 2019; p. 020025.
14. Sari K. Rüter A, Martin M, Haputhanthri R, Deacon G, Li H, Wood B. The Application of ATR-FTIR Spectroscopy and the Reversible DNA Conformation as a Sensor to Test the Effectiveness of Platinum (II) Anticancer Drugs. *Sensors* 2018; 18: 4297.
15. Niswaghi Z, Rehman S, Ihtehsham, Fourier Transform (FTIR) Spectroscopy of Biological Tissues. *Applied Spectroscopy Review* 2008, 43: 134-179
16. Achy J, Mantovani MS, Morales MAM. Analysis of the genotoxic potential of low concentrations of Malathion on the *Allium Ceppa* cells and rat hepatoma tissue culture. *Journal of Environmental Sciences*. 2015; 36:102-11.
17. Ar, N. J. Y., Khaw, K. Y., Murugaiyah, V., & Lai, C. S. Cholinesterase inhibitory activity and chemical constituents of *Stenochlaena palustris* fronds at two different stages of maturity. *Journal of Food and Drug Analysis* 2016; 24: 358-366
18. Garcia FP, Ascencio SYC, Oyarzun JCG, Hernandez AC, Alavarado PV. Pesticides : classification, uses, and toxicity. Measures of exposure and genotoxic risks. *J Res. Environ. Sci. Toxicol* 2012; 1(11):279-93.
19. Sa A, Yaduvanshi SK, Pant SC, Lomash V, Srivastava N. Evaluation of DNA damage and cytotoxicity induced by three commonly used organophosphate pesticides individually and in mixture, in rat tissues *Environ. Toxicol* 2013; 28: 543–552.
20. Djovic, MB. Krstic DZ, Lazarevic-Pasti, T.D, Bondzic, A.M, Vasic, VM. Acetylcholinesterase inhibitors: pharmacology and toxicology. *Curr. Neuropharmacol* 2013; 11: 315–335.
21. Li, Guowen Zhang, Mo Tao. 2014. Binding properties herbicide chlorpropham to DNA: Spectroscopy, Chemometrics, and modelling investigations, *Journal Photochemistry and Photobiology B: Biology* 2014; 138:109-117

Query Report

Q1 Article Type is missing

Q2 Only 1 affiliation will be considered. Plz confirm which one must be written here

Q3 Plz share the department name again

Q4 Short title of the Article is missing

Q5 Provide full name

Q6 Reference 15 or 26?

Q7 Plz correct the word

Q8 Cross-check for accuracy/data/format, correct, and provide all references in consistent Vancouver format.

Sample reference formatted to act as an example and to maintain consistency of format (Vancouver) required

Kumar RA, Sridevi K, Kumar NV, Nanduri S, Rajagopal S. Anticancer and immunostimulatory compounds from *Andrographis paniculata*. *Journal of Ethnopharmacology*. 2004;92:291-295. Available from: doi.org/10.1016/j.jep.2004.03.004

UNIVERSITAS
LAMBUNG MANGKURAT

Eko Suhartono <esuhartono@ulm.ac.id>

IJDDT - Vol 12 Issue 01 - Eko Suhartono - IJDDT,Vol12,Issue1,Article3 v1 - First author proof with queries

2 messages

Rohit Jaiswal <process@mripub.com>
To: esuhartono@ulm.ac.id
Cc: pradeep@mripub.com, ashwani@mripub.com

Thu, Mar 31, 2022 at 3:17 PM

Dear Author(s),

Please find attached the first author proof with queries, for your kind perusal.

Your valuable feedback is awaited.

DOs:

1. Mention your comments/responses to our specific queries (if any), in the attached **PDF file**.
2. While responding to our queries, either add **Sticky Notes** to the attached first author proof (PDF) file point-wise, **or** respond specifically to our queries in a separate **word file**.
3. We shall do further processing once we receive your valuable responses to **all** our queries.
4. Select "**Reply All**" while sending your response(s).
5. Please send your response(s) within **24 hours** of the receipt of this email.

DON'Ts:

1. **Do not modify the subject-line of this email.**
2. Do not draft a new email to send your response; only respond to this email to keep track.

Please feel free to mail us for any further assistance.

Regards,

Rohit

**Eko Suhartono - IJDDT,Vol12,Issue1,Article3 v1.pdf**
371K

Eko Suhartono <esuhartono@ulm.ac.id>
To: Rohit Jaiswal <process@mripub.com>

Thu, Mar 31, 2022 at 8:48 PM

Dear Rohit

We resubmit articles that have been revised according to the instructions.

Best regard

Eko Suhartono

[Quoted text hidden]



Eko Suhartono - IJDDT,Vol12,Issue1,Article3 v1.pdf

444K