

# Short Communication: Bird species in the cement factory complex of Tarjun, South Kalimantan, Indonesia

*by Mochamad Arief Soendjoto*

---

**Submission date:** 25-May-2023 11:01AM (UTC+0700)

**Submission ID:** 2101344731

**File name:** 2019\_Biodiversitas\_20\_1\_218-225\_Riefani\_et\_al.\_Lengkap.pdf (1.19M)

**Word count:** 7315

**Character count:** 43309

## Short Communication: Bird species in the cement factory complex of Tarjun, South Kalimantan, Indonesia

MAULANA KHALID RIEFANI<sup>1\*</sup>, MOCHAMAD ARIEF SOENDJOTO<sup>2</sup>, ANDY M. MUNIR<sup>3</sup>

<sup>1</sup>Faculty of Teaching and Educational Sciences, Universitas Lambung Mangkurat, Jl. Hasan Basry, Kayutangi, Banjarmasin 70123, South Kalimantan, Indonesia. \* email: maulanakriefani@ulm.ac.id

<sup>2</sup>Forestry Faculty, Universitas Lambung Mangkurat, Jl. Ahmad Yani Km 36, Banjarbaru 70714, South Kalimantan, Indonesia

<sup>3</sup>PT Indocement Tunggul Prakarsa Tbk., Tarjun, Kotabaru, Indonesia

Manuscript received: 29 September 2018. Revision accepted: 21 December 2018.

**Abstract.** Riefani MK, Soendjoto MA, Munir AM. 2019. Bird species in the cement factory complex of Tarjun, South Kalimantan, Indonesia. *Biodiversitas* 20: 218-225. The factory complex of PT Indocement Tunggul Prakarsa Tbk., Tarjun, South Kalimantan, Indonesia consists of cement manufacturing plants and employee settlements. Supporting facilities for the two main areas are open spaces in the form of waters (lakes, small rivers), grasslands (golf courses, grassy fields with several woody individuals growing on it), and forests (mangrove forests, secondary lowland forests, including shrubs). However, the data related to bird species that live in complex are not available yet. The purpose of this study was to inventory bird species in the ITP complex and to analyze qualitatively the presence of birds. These data will be used as a baseline for the development of the complex, comparative data for birds in the operational area of mining located 27 km outside the complex, and the completeness of the requirements for submitting a company performance rating (Proper). Birds were inventoried four periods between 2015-2016 at 06:00 - 08:00 and 16:30 - 18:00. The period here was treated as repetition. Observers equipped with binoculars and telephoto cameras walked around through footpaths and roads exploring the whole complex. In water and grassland areas, birds recorded can be at an observation distance of more than 50 m, whereas in forest areas, birds recorded were at a maximum distance of 50 m. A total of eighty-seven species and 41 families of birds were recorded. Of them, eighteen species and 8 families are waterbirds. The most recorded birds were classified as residents. All of the birds were recorded or observed easily every day or almost every day, at least at the research sites. Others (8 species) were categorized as migratory birds. Fourteen species are protected by the Regulation of the Minister of Environment and Forestry, the Republic of Indonesia No. P.92/MENLHK/SETJEN/KUM.1/8/2018, 30 August 2018.

**Keywords:** Birds inventory, PT Indocement Tunggul Prakarsa, open space, settlement, South Kalimantan.

### INTRODUCTION

PT Indocement Tunggul Prakarsa Tbk. (ITP) is the second largest cement producer in Indonesia (2018b) and operates three cement-producing factories (Heidelberg Cement Group 2018). One factory (with 10 plants) is located at Citeureup, Bogor, West Java Province, 1 factory (with 2 plants) at Palimanan, Cirebon, West Java Province, and the other factory (1 plants) at Tarjun Village, Kelumpang Hilir District, Kotabaru Regency South Kalimantan Province.

The Tarjun Factory Complex lies on the shore of Laut Strait, a narrow sea that extends from north to south and separates Laut Island in a total of  $\pm 1.873,36$  km<sup>2</sup> (east) from Borneo which covers a total of 743.330 km<sup>2</sup> (west). Borneo is the third-largest island in the world and controlled by Indonesia, Malaysia, and Brunei Darussalam. The ITP factory location is strategic. From a plant with a production capacity of 2.6 million tons per year (ITP 2018c), cement both in packaging and in bulk form can be loaded directly to the ship to be transported to distributors and retailers in Java, Sumatra, Kalimantan, Sulawesi and other islands in Indonesia.

The ITP complex covering an area of about 586 ha (ITP 2016, 2017) consists of two main areas, namely cement manufacturing plants and employee settlements. Supporting facilities for two main areas are open spaces in the form of waters (lakes, ponds, creeks), grassland areas (golf courses, soccer fields, grass fields with some woody individuals growing on them), forests (mangroves, dry-land forests, including shrubs and nursery gardens). The function of these supporting facilities certainly makes this area safe and ecologically safe, especially for factory employees and buyers from the surrounding areas (in South Kalimantan and East Kalimantan Province) who come directly to drive cement transport vehicles (trucks), and anyone who gets permission to enter the ITP complex. Employees can work according to or even exceed their performance targets and find their nuclear families healthy and not anxious as long as they are present or even absent from their daily lives. Buyers can also use their free time in stress-free conditions while waiting to get calls from the factory authority or waiting for their turn to load cement into their transport vehicles.

On the other hand, the supporting facilities become a safe and comfortable place for plants and animals. The birds currently discussed can do all kinds of behavior

without fear of being hunted or consumed by people. Birds can play, search food, and breed to keep their population preserved or even increase. Moreover, the company has committed itself to conserving biodiversity and natural resources whose strategies include preventing extinction of endemic birds at the plant site and planting trees to develop bird habitats (ITP 2015).

Unfortunately, data related to bird species that live in the Tarjun Factory Complex are not sufficient. We did not get environmental documents (such as Environmental Impact Analysis) or official reports containing fauna data (especially in this case, birds) in the Tarjun Factory Complex. Data about birds were actually obtained from two other factory complexes. In the Citeureup Factory Complex 41 species of birds were recorded in 2015 (ITP 2016) and then 83 species in 2016 (ITP 2017) and also in 2017 (ITP 2018d). One of them was Javan Kingfisher (*Halcyon cyanoventris*), endemic species on the Java Island (ITP 2018d). In the Palimanan Factory Complex 66 species of birds were found in 2016 (ITP 2017) and also in 2017 (ITP 2018d). One of them was Java Sparrow (*Padda oryzivora*), an endemic species of Java Island and vulnerable based on the International Union for the Conservation of Nature and Natural Resources (IUCN) (ITP 2018d).

The purpose of this study was to inventory bird species in the Tarjun Factory Complex and to qualitatively analyze the factors that caused the existence of birds. Data obtained from this bird inventory activity has multiple benefits. The data will be used as a baseline for the continuous development of the Tarjun Factory Complex, comparative data with bird data that are presented or found in the operational area of cement raw material mining located 27 km outside the complex, and the completeness of important requirements submitted for the assessment program for company performance (Program Penilaian Peringkat Kinerja Perusahaan, PROPER). The annual assessment program is launched by the Ministry of Environment and Forestry, the Republic of Indonesia to assess the company's compliance with environmental management (related to water management, air management, and management of waste from hazardous and toxic materials), development of community welfare, and clean production. Companies involved in PROPER are those whose industries have an important impact on the environment and their companies care about reputation.

## MATERIALS AND METHODS


Through the opportunistic observation method, birds found at 06:00 - 08:00 and 16:30 - 18:00 in the ITP factory complex of Tarjun in the four periods of 2015-2016 were recorded. This observation hour was actually the free time before and after we observed diurnal birds in the quarry area of cement raw-materials which is 27 km from the factory complex. The periods, namely P1 (8 - 15 April 2015), P2 (29 July - 4 August 2015), P3 (5 - 12 November

2015), and P4 (28 February - 8 March 2016) were treated as repetitions, even though observation routes in each period were not exactly the same. Habitat types were not differentiated, because we classified the entire factory complex as a modified habitat that was relatively highly influenced by human activity.

Combined transect and concentration count methods were used to be able to explore birds throughout the Tarjun Factory Complex (Figure 1). With the first method, observers recorded birds through paths (in forest areas, areas of tree vegetation, areas where people rarely pass) or paved / rocky roads (infrastructure for general use in the complex). Occasionally the observers stopped for 30 minutes. With the concentration count method, observers not only recorded, but also photographed birds that were within the range of sight. In waters and grassland areas, birds that were recorded can be at an observation distance of more than 50 m, whereas in forest areas, birds recorded at a maximum distance of 50 m. Bird species whose morphology or sounds were known directly by observers were recorded in the databook. However, birds whose morphology and sound were unknown or doubtful were further observed with the help of binoculars or cameras equipped with telelens (80 x 400) or (70 x 300). The field guides used to identify birds were MacKinnon et al. (2010) and Soendjoto et al. (2015b). The comparison field guides were Smythies (1981) and Strange (2002). Then the taxonomic names are referred to IUCN (2018).

The data is tabulated in a 14 column table. A column after the serial number column (first column) contains the family and scientific names of the bird species. Both the family name and the species name in this column are arranged in alphabetical A-Z.

Furthermore, birds are classified (i) whether they are waterbirds, (ii) according to their occupancy status, and (iii) according to their protection status. Birds are classified as waterbirds if their morphology refers to the criteria of waterbirds, such as both long feet and long beak (for example Anhingidae, Ardeidae, Charadriidae, Ciconiidae, Rallidae, and Scolopacidae) or have membranes between their fingers (for example Anatidae and Laridae). The bird family taxa (32 families) included in waterbirds are listed by Wetlands International (2012). Bird occupancy status consists of two groups, namely resident and migrant. Bird species is a resident if this bird is recorded or observed easily every day or almost every day, at least in the ITP complex and its surroundings. Migratory bird status refers to MacKinnon et al. (2010). The status of bird protection is determined according to Regulation of the Minister of Environment and Forestry, the Republic of Indonesia No. P.92/MENLHK/SETJEN/KUM.1/8/2018, 30 August 2018, concerning Amendment to Regulation of the Minister of Environment and Forestry No. P.20/MENLHK/SETJEN/KUM.1/6/2018 concerning the Protected Species of Plants and Animals. This is a substitute for Government Regulation No. 7, 27 January 1999 concerning Preservation of Species of Plants and Animals.



**Figure 1.** Geographical position of PT. Indocement Tungal Prakarsa Tbk., Tarjun

## RESULTS AND DISCUSSION

A total of 87 species belongs to 41 families of birds were recorded (Table 1). Of these species, 14 are protected birds and eighteen species (8 families) are waterbirds. Most birds are residents (79 of 87 bird species) and others are categorized as migrants. Five of the 8 species of migratory birds are waterbirds, two species are predators, and the rest is Java Sparrow, species specifically in this study are grouped in migrants. Java Sparrow is an endemic species of Java, Bali, and Kangean which was then widely introduced from Southeast Asia to Australia (Smythies 1981; MacKinnon et al. 2010).

These waterbirds can be recorded in the waters of the complex. One of them is Lake Tiro (the acronym of Tiga Roda, Three Wheels in English), which is about 5 ha in size. The widest lake in the factory complex is formed after the flow of water in the river which empties into the Terusan River is dammed. The edge of the lake bordering

the settlement, in general, is relatively shallow. Lake-forming mud appears, especially if the volume of lake water decreases or subsidizes. In muddy locations that are not overgrown with grass or overgrown with grass, found waterbirds that have long legs and neck and large body sizes such as Great Egret, and Little Egret. In addition, there are also feet and long necks, but small birds, like the Common Sandpiper. At the edge of the lake overgrown with Poaceae (grass family) and Cyperaceae (sedge family) can be found Javan Pond-heron, Striated Heron, White-breasted Waterhen, Common Moorhen, and Wood Sandpiper. On cables stretching over the surface of the lake or at the end of wooden pillars that stick to the bottom of the lake, birds like Little Tern and Common Tern often sit. At the edge of the lake near the dam wall which limits lake water with relatively deep rivers, some waterbirds are found swimming. Birds include Oriental Darter, Wandering Whistling-duck, and Sunda Teal.

**Table 1.** Bird species recorded in the factory complex of PT Indocement Tunggal Prakarsa Tbk., Tarjun

Family and scientific name	Indonesian name	Common name	Obs. period				IUCN status	Waterbirds		Prot.	Occupancy	
			P1	P2	P3	P4		Fam.	Sp.		Res.	Mig.
<b>Acanthizidae</b>												
<i>Gerygone sulphurea</i>	Remetek laut	Golden-bellied Gerygone	●	●	●	●	LC-d	-	-	-	●	-
<b>Accipitridae</b>												
<i>Haliaeetus leucogaster</i>	Elang-laut perut-putih	White-bellied Fish-eagle	-	-	●	●	LC-d	-	-	●	●	-
<i>Haliastur indus</i>	Elang bondol	Brahminy Kite	●	●	●	●	LC-d	-	-	●	●	-
<b>Aegithinidae</b>												
<i>Aegithina tiphia</i>	Cipoh kacat	Common Iora	-	-	-	●	LC-u	-	-	-	●	-
<i>Aegithina viridissima</i>	Cipoh jantung	Green Iora	●	●	●	●	NT-d	-	-	-	●	-
<b>Alcedinidae</b>												
<i>Alcedo meninting</i>	Raja-udang meninting	Blue-eared Kingfisher	●	●	●	●	LC-d	-	-	-	●	-
<i>Todiramphus chloris</i>	Cekakak sungai	Collared Kingfisher	●	●	●	●	LC-d	-	-	-	●	-
<i>Todiramphus sanctus</i>	Cekakak suci	Sacred Kingfisher	●	●	●	●	LC-i	-	-	-	●	-
<i>Pelargopsis capensis</i>	Pekaka emas	Stork-billed Kingfisher	●	●	●	●	LC-d	-	-	-	●	-
<b>Anatidae</b>												
<i>Anas gibberifrons</i>	Itik benjut	Sunda Teal	-	-	●	●	NT-s	●	●	-	●	-
<i>Dendrocygna arcuata</i>	Belibis kembang	Wandering Whistling-duck	●	●	●	●	LC-d	-	●	-	●	-
<b>Anhingidae</b>												
<i>Anhinga melanogaster</i>	Pecuk-ular asia	Oriental Darter	-	-	●	●	NT-d	●	●	●	●	-
<b>Ardeidae</b>												
<i>Ardeola speciosa</i>	Blekok sawah	Javan Pond-heron	-	-	●	●	LC-u	-	●	-	●	-
<i>Butorides striata</i>	Kokokan laut	Striated Heron	●	●	●	●	LC-d	-	●	-	●	-
<i>Ardea alba</i>	Kuntul besar	Great Egret	-	-	●	●	LC-u	-	●	-	●	-
<i>Egretta garzetta</i>	Kuntul kecil	Little Egret	●	●	●	●	LC-i	-	●	-	●	-
<i>Egretta sacra</i>	Kuntul karang	Pacific Reef-egret	-	-	●	●	LC-s	-	●	-	●	-
<i>Ixobrychus cinnamomeus</i>	Bambangan merah	Cinnamon Bittern	●	●	●	●	LC-s	-	●	-	●	-
<i>Ixobrychus sinensis</i>	Bambangan kuning	Yellow Bittern	●	●	●	●	LC-u	●	●	-	●	-
<b>Artamidae</b>												
<i>Artamus leucoryn</i>	Kekep babi	White-breasted Woodswallow	●	●	●	●	LC-s	-	-	-	●	-
<b>Bucerotidae</b>												
<i>Anthracoceros albirostris</i>	Kangkareng perut putih	Oriented Pied Hornbill	-	●	-	-	LC-s	-	-	●	●	-
<b>Campephagidae</b>												
<i>Lalage nigra</i>	Kapasan kemiri	Pied Triller	●	●	●	●	LC-s	-	-	-	●	-
<i>Pericrocotus flammeus</i>	Sepah hutan	Scarlet Minivet	-	-	●	●	LC-s	-	-	-	●	-
<b>Caprimulgidae</b>												
<i>Caprimulgus affinis</i>	Cabak kota	Savannah Nightjar	-	●	●	●	LC-s	-	-	-	●	-
<b>Charadriidae</b>												
<i>Charadrius dubius</i>	Cerek-kalung kecil	Little Ringed Plover	-	-	●	●	LC-s	●	●	-	-	●
<b>Ciconiidae</b>												
<i>Leptoptilos javanicus</i>	Bangau tongtong	Lesser Adjutant	●	●	●	●	Vu-d	●	●	●	●	-
<b>Cisticolidae</b>												
<i>Orthotomus ruficeps</i>	Cinene kelabu	Ashy Tailorbird	●	●	●	●	LC-s	-	-	-	●	-
<i>Orthotomus sericeus</i>	Cinene merah	Rufous-tailed Tailorbird	●	●	●	●	LC-s	-	-	-	●	-
<i>Prinia flaviventris</i>	Prenjak rawa	Yellow-bellied Prinia	●	●	●	●	LC-d	-	-	-	●	-
<b>Columbidae</b>												
<i>Geopelia striata</i>	Perkutut	Zebra Dove	●	●	●	●	LC-s	-	-	-	●	-
<i>Spilopelia chinensis</i>	Tekukur	Eastern Spotted Dove	●	●	●	●	LC-i	-	-	-	●	-
<i>Treron vernans</i>	Punai gading	Pink-necked Green-pigeon	-	-	●	●	LC-s	-	-	-	●	-
<b>Corvidae</b>												
<i>Corvus macrorhynchos</i>	Gagak kampung	Large-billed Crow	-	-	●	-	LC-s	-	-	-	●	-
<b>Cuculidae</b>												
<i>Cacomantis merulinus</i>	Wiwik kelabu	Plantive Cuckoo	-	-	●	●	LC-s	-	-	-	●	-
<i>Centropus bengalensis</i>	Bubut kecil	Lesser Coucal	-	●	-	●	LC-i	-	-	-	●	-
<i>Centropus sinensis</i>	Bubut besar	Greater Coucal	-	-	●	●	LC-s	-	-	-	●	-
<i>Rhinortha chlorophaea</i>	Kadalan selaya	Raffles's Malkoha	-	-	●	●	LC-s	-	-	-	●	-
<i>Phaenicophaeus curvirostris</i>	Kadalan birah	Chestnut-breasted Malkoha	-	-	●	-	LC-s	-	-	-	●	-
<b>Dicaeidae</b>												
<i>Dicaeum cruentatum</i>	Cabai merah	Scarlet-backed Flowerpecker	-	-	-	●	LC-s	-	-	-	●	-
<i>Dicaeum trochileum</i>	Cabai jawa	Scarlet-headed Flowerpecker	●	●	●	●	LC-s	-	-	-	●	-



<i>Dicaeum trigonostigma</i>	Cabai bunga api	Orange-bellied Flowerpecker	- ● - ●	LC-s	-	-	-	●	-
<b>Estrildidae</b>									
<i>Lonchura fuscans</i>	Bondol kalimantan	Dusky Munia	● ● ● ●	LC-s	-	-	-	●	-
<i>Lonchura malacca</i>	Bondol rawa	Black-headed Munia	- - ● ●	LC-s	-	-	-	●	-
<i>Lonchura oryzivora</i>	Gelatik jawa	Java Sparrow	- - - ●	Vu-d	-	-	-	-	●
<i>Lonchura punctulata</i>	Bondol peking	Scaly-breasted Munia	● ● ● ●	LC-s	-	-	-	●	-
<b>Falconidae</b>									
<i>Falco moluccensis</i>	Alap-alap sapi	Spotted Kestrel	- - ● ●	LC-i	-	-	●	-	●
<b>Hemiprocnidae</b>									
<i>Hemiprocne longipennis</i>	Tepekong jambul	Grey-rumped Treeswift	- - ● ●	LC-u	-	-	-	●	-
<b>Hirundinidae</b>									
<i>Hirundo tahitica</i>	Layang-layang batu	Pacific Swallow	● ● ● ●	LC-u	-	-	-	●	-
<b>Laniidae</b>									
<i>Lanius schach</i>	Bentet kelabu	Long-tailed Shrike	● ● ● ●	LC-u	-	-	-	●	-
<b>Laridae</b>									
<i>Sterna hirundo</i>	Dara-laut biasa	Common Tern	- - ● ●	LC-u	●	●	●	-	●
<i>Sternula albifrons</i>	Dara-laut kecil	Little Tern	- - ● ●	LC-d	-	●	●	-	●
<b>Megalaimidae</b>									
<i>Psilopogon rafflesii</i>	Takur tutut	Red-crowned Barbet	- - - ●	NT-d	-	-	-	●	-
<i>Caloramphus fuliginosus</i>	Takur ampis	Brown Barbet	- - ● -	LC-d	-	-	-	●	-
<b>Meropidae</b>									
<i>Merops philippinus</i>	Kirik-kirik laut	Blue-tailed Bee-eater	- - ● ●	LC-s	-	-	-	●	-
<b>Motacillidae</b>									
<i>Motacilla flava</i>	Kicuit kerbau	Western Yellow Wagtail	- - ● -	LC-d	-	-	-	●	-
<i>Anthus novaeseelandiae</i>	Apung tanah	Common Pipit	- - ● ●	LC-s	-	-	-	●	-
<b>Muscicapidae</b>									
<i>Cyornis rufigaster</i>	Sikatan bakau	Mangrove Blue-flycatcher	- ● - ●	LC-d	-	-	-	●	-
<i>Muscicapa dauurica</i>	Sikatan bubuk	Asian Brown Flycatcher	- - ● ●	LC-s	-	-	-	●	-
<b>Nectariniidae</b>									
<i>Aethopyga siparaja</i>	Burung-madu sepah raja	Crimson Sunbird	● ● ● ●	LC-s	-	-	●	●	-
<i>Anthreptes malacensis</i>	Burung-madu kelapa	Plain-throated Sunbird	● ● ● ●	LC-s	-	-	-	●	-
<i>Anthreptes simplex</i>	Burung-madu polos	Plain Sunbird	- - ● ●	LC-d	-	-	-	●	-
<i>Arachnothera longirostra</i>	Pijantung kecil	Little Spiderhunter	- - - ●	LC-s	-	-	-	●	-
<i>Chalcoparia singalensis</i>	Burung-madu belukar	Ruby-cheeked Sunbird	- - - ●	LC-s	-	-	-	●	-
<i>Cinnyris jugularis</i>	Burung-madu sriganti	Olive-backed Sunbird	● ● ● ●	LC-s	-	-	-	●	-
<i>Leptocoma calcostetha</i>	Burung-madu bakau	Copper-throated Sunbird	● ● ● ●	LC-s	-	-	-	●	-
<b>Pandionidae</b>									
<i>Pandion haliaetus</i>	Elang tiram	Osprey	- - - ●	LC-i	-	-	●	-	●
<b>Paridae</b>									
<i>Parus major</i>	Gelatik batu kelabu	Great Tit	- - ● -	LC-i	-	-	-	●	-
<b>Passeridae</b>									
<i>Passer montanus</i>	Burung gereja	Tree Sparrow	● ● ● ●	LC-d	-	-	-	●	-
<b>Picidae</b>									
<i>Dinopium javanense</i>	Pelatuk besi	Common Goldenback	● ● ● ●	LC-d	-	-	-	●	-
<i>Meiglyptes tukki</i>	Caladi badok	Buff-necked Woodpecker	- - ● ●	NT-d	-	-	-	●	-
<i>Picoides moluccensis</i>	Caladi tilik	Sunda Woodpecker	● ● ● ●	LC-i	-	-	-	●	-
<i>Chrysophlegma miniaceum</i>	Pelatuk merah	Banded Woodpecker	● ● ● -	LC-s	-	-	-	●	-
<b>Psittacidae</b>									
<i>Loriculus galgulus</i>	Serindit melayu	Blue-crowned Hanging-Parrot	- - - ●	LC-s	-	-	●	●	-
<i>Psittacula longicauda</i>	Betet ekor-panjang	Long-tailed Parakeet	● ● ● ●	NT-d	-	-	●	●	-
<b>Pycnonotidae</b>									
<i>Pycnonotus aurigaster</i>	Cucak kutilang	Sooty-headed Bulbul	● ● ● ●	LC-d	-	-	-	●	-
<i>Pycnonotus brunneus</i>	Merbah mata-merah	Red-eyed Bulbul	- - - ●	LC-d	-	-	-	●	-
<i>Pycnonotus goiavier</i>	Merbah cerucuk	Yellow-vented Bulbul	● ● ● ●	LC-i	-	-	-	●	-
<i>Pycnonotus plumosus</i>	Merbah belukar	Olive-winged Bulbul	● ● ● ●	LC-s	-	-	-	●	-
<b>Rallidae</b>									
<i>Amaurornis phoenicurus</i>	Kareo padi	White-breasted Waterhen	● ● ● ●	LC-u	-	●	-	●	-
<i>Gallinula chloropus</i>	Mandar batu	Common Moorhen	● ● ● ●	LC-s	●	●	-	●	-
<b>Rhipiduridae</b>									
<i>Rhipidura javanica</i>	Kipasan belang	Pied Fantail	● ● ● ●	LC-s	-	-	●	●	-
<b>Scolopacidae</b>									
<i>Actitis hypoleucos</i>	Trinil pantai	Common Sandpiper	● ● ● ●	LC-d	-	●	-	-	●
<i>Tringa glareola</i>	Trinil semak	Wood Sandpiper	- ● ● ●	LC-s	●	●	-	-	●
<b>Sturnidae</b>									
<i>Acridotheres javanicus</i>	Kerak kerbau	Javan Myna	- - ● ●	Vu-d	-	-	-	●	-

<i>Aplonis panayensis</i>	Perling kumbang	Asian Glossy Starling	- - ● ●	LC-u	-	-	-	●	-
<b>Timaliidae</b>									
<i>Mixornis gularis</i>	Ciung-air coreng	Striped Tit-babbler	● ● ● ●	LC-s	-	-	-	●	-
<b>Zosteropidae</b>									
<i>Zosterops flavus</i>	Kacamata jawa	Javan White-eye	- - - ●	Vu-d	-	-	●	●	-
		<b>Σ species/period</b>	<b>42 48 73 80</b>	-	-	-	-	-	-
		<b>Σ all species</b>	<b>- - - 87 87</b>	-	-	<b>18 14</b>	<b>79</b>	<b>8</b>	-
		<b>Σ all families</b>	<b>- - - 41</b>	-	<b>8</b>	-	-	-	-

Note: Observation period: P1 (8 - 15 April 2015), P2 (29 July - 4 August 2015), P3 (5 - 12 November 2015), and P4 (28 February - 8 March 2016). Waterbirds: Fam. = family; Sp. = species; LC = least concern; NT = near threaten; Vu = vulnerable. i = increasing; d = decreasing; s = stable; u = unknown. Occupancy: Res. = resident; Mig. = migrant. Prot. = protected by the Regulation of the Minister of Environment and Forestry, the Republic of Indonesia No. P.92/MENLHK/SETJEN/KUM.1/8/2018, 30 August 2018

Waterbirds can also be recorded in small ponds scattered in the complex. The size of the pond varies from around 20 m<sup>2</sup> to 500 m<sup>2</sup> with a maximum depth of around 3 m. Some ponds are usually visited by fish anglers. Waterbirds commonly recorded in ponds like this include Striated Heron and Little Egret. This bird can be found in the pond if there are no humans (such as anglers, lawn mowers) near the pond. If there is, the position of humans is relatively far from birds. The activity is not excessive or does not disturb the birds.

The number of birds in the Tarjun Factory Complex should be compared to that in the same complex according to the observation period to analyze the development of birds in the complex. However, this cannot be done because there were no previous reports. However, when compared to data from other ITP factory complexes, namely Citeurep and Palimanan and without differentiating their habitat types, the number of birds in Tarjun is slightly larger.

The number of species that are classified as numerous and varied in the Tarjun Factory Complex may be due to three factors. First, the ITP complex has a diversity of landscapes (waters, grasslands, vegetation areas) and also plant diversity growing on it. The quality of landscapes, as well as, plants gradually increases in accordance with age and the positive treatments given to it. In the beginning, the landscape can be categorized as artificial and some plants that grow and develop on it are intentionally planted. Habitat conditions with various species of life or life interactions between physical components make a safe and comfortable habitat for birds. Different habitat from birds caused the feed resources (quantity and quality), competition between fauna in the resources (space and time).

The complex was built by another company before according to ITP (2018a), was acquired by ITP on December 29, 2000. Indeed there is no data related to flora fauna species in the landscape published to the public. However some plants that were visually identified intentionally planted were forestry plants, such as *Acacia mangium*, *A. auriculiformis*, mahoni (*Swietenia macrophylla*), sengon (*Paraserianthes falcataria*), trembesi (*Samanea saman*), horticultural crops, such as patai (*Parkia speciosa*), mango (*Mangifera* spp.), watery rose-apple (*Syzygium aqueum*), apple guava (*Psidium guajava*), and banana (*Musa* spp.), and decorative plants, such as

peacock flower (*Caesalpinia pulcherrima*), asoka (*Ixora* spp.), and *Hippobroma longiflora*. In addition, some plants grow spontaneously, such as bamboo, various species of mahang (*Macaranga* spp.), karamunting (*Rhodomyrtus tomentosa*), ketapang (*Terminalia catappa*), alaban (*Vitex pubescens*), kirinyuh (*Chromolaena odorata*), tembelekan (*Lantana camara*), Poaceae, and Cyperaceae.

Changes or differences in habitats affect animals; in this discussion birds. The landscape gradients affect the diversity and distribution of avifauna (Altaf et al. 2018). A gradient from arable fields to forest affects farmland and forest avifauna species (Kosickia and Chylarecki 2012). The diversity of habitat types which are reflected by physical conditions (watery or not, there are buildings or not, there are human activities or not) and biological conditions (plant species that form habitats and their vegetation strata) triggers bird species diversity (Soendjoto et al. 2014). The habitat changes from natural forest to secondary forest, land with agroforestry systems, and land with annual cultures gradually alter the composition of bird species (Waltert et al. 2004). The number of bird species in the reclaimed and revegetated ex-mining sites in South Kalimantan Province was increased as long as the vegetation in the site was preserved (Soendjoto et al. 2018). The old age stands of large-scale oil palm plantations have more diverse birds than the young one (Erniwati et al. 2016). The diversity or abundance of bird communities on oil palm plantations is worse than in natural forests (Teuscher et al. 2015); likewise in rice fields or roads when compared to agroforestry (Tanalgo et al. 2015). Microclimate, as well as both comfortable and safe conditions on vegetation and vegetated land, are needed animals, including birds need for resting, playing, hiding from predators, voicing, and even breeding (Soendjoto et al. 2015a).

Second, the conditions inside the ITP complex are balanced by the conditions of the surrounding environment which in this case also consist of landscape diversity and plant diversity. The latter landscape consists of the Terusan River, which is more than 30 m wide and more than 5 m deep, mangrove forests, lowland secondary forests and oil palm plantations. The plants are diverse. Bakau (*Rhizophora apiculata*, *R. mucronata*), rambai/mangrove apple (*Sonneratia caseolaris*), rambai bogam (*S. alba*), and langgadai/large-leafed mangrove (*Bruguiera gymnorrhiza*) are some main plants of the mangrove ecosystem. Alaban,

ketapang, piai (*Acrostichum aureum*), bintaro (*Cerbera manghas*), and rattan spontaneously develop in lowland secondary forests. Oil palm is the main crop in the large-scale plantations. Woody plants are not allowed to live on oil palm plantations. Plants commonly found in oil palm plantations are wild plants, such as ferns and grasses. The others are plants that are intentionally planted to divert pest and disease vectors to attack oil palm. The latter plant is *Turnera subulata*, a flowering plant of the Turneraceae family.

Landscape and plants outside the ITP complex are very useful for birds. This area can be used by birds at least to save themselves. On the other side, although the movement of birds or animals from one place or another does not recognize administrative boundaries, this limit can be used as an imaginary line that distinguishes habitat types or human treatments in habitat (anthropogenic factors). When the birds inside the complex are disturbed, they fly or move to adjacent landscapes outside the complex. Factors that may play an important role in the similarity of bird communities are the short distance between two adjacent forest types, the abundance of insect food sources, and the succession stages similar to forest types (Kaban et al. 2017). Hunting can make a bird behavior change, in the form of increased chances of flying, more time wasted on alert, and less rest or bird move from hunting areas to areas where there is no hunting (Casas et al. 2009).

Third, the imposition of a hunting ban on ITP complex facilitates birds to be safe and comfortable from interference. Hunting ban boards that also contain sanctions for violations of these regulations are installed in strategic places (such as public transport roads and lakeside) that are easily seen by employees or the public entering the complex. The ban is obeyed by them. In turn, it has a positive impact. Under certain conditions, birds can adapt physiologically and behaviorally (Sokos et al. 2013). Birds are accustomed to these conditions so that population and composition or in general bird diversity increases gradually.

The causes of the reduction in bird populations are not only deforestation (Symes et al. 2018), but also bird hunting for food, used as pets, traded, or other purposes (Fernandes-Ferreira et al. 2012, Iskandar et al. 2016). Hunting can change animal community composition and may affect a large number of ecological interactions (Barbosa et al. 2014) and jeopardizes the survival of birds, especially waterbirds (Wang et al. 2018) and migratory bird species (Arizaga and Laso 2015). Although not a few researchers argue that hunting endangers the survival of bird species, some researchers who argue otherwise. Hunting has no significant effect on bird abundance (Sulaiman et al. 2014).

In addition to the total number of bird species found (including waterbirds), bird protection status, and bird occupancy status, there are results of research in the ITP factory complex that needs attention. The number of bird species found in periods 1 and 2 is smaller than in periods 3 and 4.

The main objective of this study is not to distinguish birds in the dry and rainy seasons. Thus, a mere

coincidence, if there are differences in the number of bird species in periods 1 and 2 compared to periods 3 and 4. In the case of climate in Indonesia, April - October is called the dry season. Conversely, October - April is the rainy season. In the dry season, grass generally dries. The leaves of some trees even fall out. This also happened in the Tarjun Factory Complex. Exceptions only occur in the grass on the golf course. The grass here remains green because it is treated (watered and cut) every day.

In the dry season, plants as food sources for birds in the factory complex are not available or not enough to meet the nutritional needs of birds. Conditions like this force the birds to be more active in spreading outside the complex which has sufficient food sources. Spreading to wider or farther areas can be done by birds in a short time because these animals can fly to move from unsafe and uncomfortable places to other places that are safer and more comfortable. Not surprisingly, if later within a relatively short duration of observation, i.e., 2 hours in the morning and 1.5 hours in the afternoon the number of bird species observed were 42 and 48 respectively.

The opposite happened in the rainy season. Plants are fresh and green. Most plants even produce fruit and stimulate insects to come, so that the plant environment provides enough food for birds. Birds also do not need to spread to find food to another place farther away. In this condition, 73 species of birds in period 3 and 80 species in period 4 were recorded. This size includes migratory birds with relatively few species (only 8 species or even 7 species, if Java Sparrow is removed from the category of migratory birds).

#### ACKNOWLEDGEMENTS


We would like to thank the ITP Manager at Tarjun, South Kalimantan who financed and facilitated this research. Further our sincere gratitude also goes to Wisnu Adi Patria for extensive support given during field surveys.

#### REFERENCES

- Altaf M, Javid A, Khan AM, Khan MSH, Umair M, Ali Z. 2018. Anthropogenic impact on the distribution of the birds in the tropical thorn forest, Punjab, Pakistan. *Journal of Asia-Pacific Biodiversity* 11(1): 229-236. DOI 10.1016/j.japb.2018.03.001.
- Arizaga J, Laso M. 2015. A quantification of illegal hunting of birds in Gipuzkoa (North of Spain). *Eur J Wildl Res* 61:795-799. DOI 10.1007/s10344-015-0940-6.
- Barbosa EDO, da Silva MGB, de Medeiros RO, Chaves MF. 2014. Hunting activities impact avifauna in rural areas of the Jaçanã municipality, Rio Grande do Norte State, Brazil. *Biotemas* 27(3): 175-190.
- Casas F, Mougeot F, Vinueza J, Bretagnolle V. 2009. Effects of hunting on the behaviour and spatial distribution of farmland birds: importance of hunting-free refuges in agricultural areas. *Anim Conserv* 2009: 1-9.
- Erniwati, Zuhud EAM, Santosa Y, Anas I. 2016. The value of secondary forest patches for bird conservation in palm oil landscapes of Riau, Sumatra. *Biodiversitas* 17(2): 791-798. DOI: 10.13057/biodiv/d170256.
- Fernandes-Ferreira H, Mendonc a SV, Albano C, Ferreira FS, Alves RRN. 2011. Hunting, use and conservation of birds in Northeast Brazil. *Biodivers Conserv* 21(1): 221-244. DOI 10.1007/s10531-011-0179-9.



- MacKinnon J, Phillipps K, Balen B. 2010. Birds in Sumatra, Java, Bali, and Kalimantan (Including Sabah, Sarawak and Brunei Darussalam). Buring Indonesia, Bogor, Indonesia. [Indonesian].
- Heidelberg Cement Group. 2018. Indonesia. <https://www.heidelbergcement.com/en/indonesia>.
- Iskandar J, Iskandar BS, Partasasmita R. 2016. The local knowledge of the rural people on species, role and hunting of birds: Case study in Karangwangi Village, West Java, Indonesia. *Biodiversitas* 17(2): 435-446. DOI 10.13057/biodiv/d170206.
- ITP. 2015. Sustainability Report 2014. Improving Efficiency, Achieving Sustainability. <http://www.indocement.co.id/v5/IndocementContent/Attachment/Report/Sustainability%20Report/03%20SR%20INDOCEMENT%202014.pdf>.
- ITP. 2016. Sustainability Report 2015. Ensuring Sustainability for the Next Decades. <http://www.indocement.co.id/v5/IndocementContent/Presentation/2016/160418%20-%20AR%20SR/Indocement%20Sustainability%20Report%202015.pdf>.
- ITP. 2017. Sustainability Report 2016. Keeping a Blue Sky: Being Different to Shape the Future. [http://www.indocement.co.id/v5/IndocementContent/Presentation/2017/ARSR2016/SR%20Indocement%202016\\_lowres.pdf](http://www.indocement.co.id/v5/IndocementContent/Presentation/2017/ARSR2016/SR%20Indocement%202016_lowres.pdf).
- ITP. 2018a. Indocement in Brief. <http://www.indocement.co.id/v5/en/company/indocement-in-brief/>
- ITP. 2018b. Indocement Opens New Cement Terminal & Aims for Low-End Market. Indonesia-Investments. <https://www.indonesia-investments.com/news/todays-headlines/indocement-opens-new-cement-terminal-aims-for-low-end-market/item8676?>
- ITP. 2018c. PT Indocement Tunggal Prakarsa Tbk. Q1 2018 Results. [http://www.indocement.co.id/v5/IndocementContent/Press%20Release/2018/Indocement%20Q1%202018%20Results\\_20180518.pdf](http://www.indocement.co.id/v5/IndocementContent/Press%20Release/2018/Indocement%20Q1%202018%20Results_20180518.pdf).
- ITP. 2018d. Sustainability Report 2017. Agility for Sustainability. <http://www.indocement.co.id/v5/IndocementContent/GCG/2018/SR%20Indocement%202017.pdf>.
- IUCN. 2018. The IUCN Red List of Threatened Species. Version 2018-1. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 14 November 2018.
- Kaban A, Mardiasuti A, Mulyani YA. 2017. Response of bird community to various plantation forests in Gunung Walat, West Java, Indonesia. *Hayati J Biosci* 24: 72-78.
- Kosickia JZ, Chylarecki P. 2012. Effect of climate, topography and habitat on species-richness of breeding birds in Poland. *Basic Appl Ecol* 13: 475-483. DOI 10.1016/j.baae.2012.07.007.
- Smythies BE. 1981. The Birds of Borneo. The Sabah Society & The Malayan Nature Society, Kuala Lumpur.
- Soendjoto MA, Riefani MK, Mahrudin, Zen M. 2014. Dynamics of avifauna species in the area of PT Arutmin Indonesia - North Pulau Laut Coal Terminal, Kotabaru, South Kalimantan. In: Karyanto P et al. (Eds). Proceedings of National Conference XI on Biology Education. Sebelas Maret University, Surakarta, 7 June 2014. Pp. 512-520. [Indonesian].
- Soendjoto MA, Riefani MK, Triwibowo D, Anshari MN, Metasari D. 2015a. Wild animals in the reclamation area of PT Adaro Indonesia, South Kalimantan revegetated less than two years. In: Sajidan et al. (Eds). Proceeding of National Conference on Conservation and Utilization of Nature Resources, Sebelas Maret University, Surakarta, Indonesia, 13 Januari 2015. pp. 192-198. [Indonesian].
- Soendjoto MA, Riefani MK, Triwibowo D, Metasari D. 2018. Birds observed during the monitoring period of 2013-2017 in the revegetation area of ex-coal mining sites in South Kalimantan, Indonesia. *Biodiversitas* 19(1): 323-329. DOI 10.13057/biodiv/d190144.
- Soendjoto MA, Riefani MK, Triwibowo D, Wahyudi F. 2015b. Avifauna in the Reclamation Area of PT Adaro Indonesia. Banjarbaru: Universitas Lambung Mangkurat Press. [Indonesian].
- Sokos CK, Birtsas PK, Connelly JW, Papaspyropoulos KG. 2013. Hunting of migratory birds: disturbance intolerant or harvest tolerant? *Wildl Biol* 19: 113-125. DOI 10.2981/12-032.
- Strange M. 2002. A Photographic Guide to the Birds of Southeast Asia, Including the Philippines & Borneo. Periplus Editions (HK) Ltd., Hong Kong.
- Sulaiman IM, Apeverga PT, Ringim AS, Ibrahim KM. 2014. Effects of anthropogenic activities on bird diversity and abundance at the Hadejia - Nguru Wetlands, Nigeria. *Stand Sci Res Essays* 2(10): 546-551.
- Symes WS, Edwards DP, Miettinen J, Rheindt FE, Carrasco LR. 2018. Combined impacts of deforestation and wildlife trade on tropical biodiversity are severely underestimated. *Nat Communications* 9(4052).
- Tanalgo KC, Pineda JAF, Agravante ME, Amerol ZM. 2015. Bird diversity and structure in different land-use types in lowland South-Central Mindanao, Philippines. *Trop Life Sci Res*. 26(2): 85-103.
- Teuscher M, Vorlaufer M, Wollni M, Brose U, Mulyani Y, Clough Y. 2015. Trade-offs between bird diversity and abundance, yields and revenue in smallholder oil palm plantations in Sumatra, Indonesia. *Biol Conserv* 186: 306-318.
- Walter M, Mardiasuti A, Mühlenberg M. 2004. The effect of land use on bird species richness in Sulawesi, Indonesia. *Conserv Biol* 15:1339-1346. DOI 10.1111/j.1523-1739.2004.00127.x.
- Wang X, Kuang F, Tan K, Ma Z. 2018. Population trends, threats, and conservation recommendations for waterbirds in China. *Avian Res* 9:14. DOI 10.1186/s40657-018-0106-9.
- Wetlands International. 2012. Waterbird Population Estimates, Fifth Edition. Summary Report. Wetlands International, Wageningen, The Netherlands.

A close-up photograph of a monkey's face, showing its large, reddish-brown nose and brown eyes. The monkey's fur is a mix of light brown and tan. The background is blurred, suggesting a natural outdoor setting.

ISSN: 1412-033X  
E-ISSN: 2085-4722

# BIODIVERSITAS

Journal of Biological Diversity

Volume 20 - Number 1 - January 2019

# BIODIVERSITAS

Journal of Biological Diversity  
Volume 20 - Number 1 - January 2019

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

## EDITORIAL BOARD (COMMUNICATING EDITORS):

**Abdel Fattah N.A. Rabou** (Palestine), **Agnieszka B. Najda** (Poland), **Alan J. Lymbery** (Australia), **Alireza Ghanadi** (Iran), **Ankur Patwardhan** (India), **Bambang H. Saharjo** (Indonesia), **Daiane H. Nunes** (Brazil), **Darlina Md. Naim** (Malaysia), **Ghulam Hassan Dar** (India), **Faiza Abbasi** (India), **Hassan Pourbabaie** (Iran), **I Made Sudiana** (Indonesia), **Ivan Zambrana-Flores** (United Kingdom), **Joko R. Witono** (Indonesia), **Katsuhiko Kondo** (Japan), **Krishna Raj** (India), **Livia Wanntorp** (Sweden), **M. Jayakara Bhandary** (India), **Mahdi Reyahi-Khoram** (Iran), **Mahendra K. Rai** (India), **Mahesh K. Adhikari** (Nepal), **Maria Panitsa** (Greece), **Muhammad Akram** (Pakistan), **Mochamad A. Soendjoto** (Indonesia), **Mohib Shah** (Pakistan), **Mohamed M.M. Najim** (Srilanka), **Morteza Eighani** (Iran), **Pawan K. Bharti** (India), **Paul K. Mbugua** (Kenya), **Rasool B. Tareen** (Pakistan), **Seweta Srivastava** (India), **Seyed Aliakbar Hedayati** (Iran), **Shahabuddin** (Indonesia), **Shahir Shamsir** (Malaysia), **Shri Kant Tripathi** (India), **Stavros Lalas** (Greece), **Subhash Santra** (India), **Sugiyarto** (Indonesia), **T.N. Prakash Kammardi** (India)

## EDITOR-IN-CHIEF:

Sutarno

## EDITORIAL MEMBERS:

English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun\_s@yahoo.com), **Artini Pangastuti** (pangastuti\_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

## MANAGING EDITORS:

**Ahmad Dwi Setyawan** (unsjournals@gmail.com)

## PUBLISHER:

The Society for Indonesian Biodiversity

## CO-PUBLISHER:

Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

## ADDRESS:

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, Email: unsjournals@yahoo.com

## ONLINE:

biodiversitas.mipa.uns.ac.id

## EXPERTISE AND CORRESPONDING EMAIL OF THE COMMUNICATING EDITORS:

**GENETIC DIVERSITY:** **Agnieszka B. Najda** (agnieszka.najda@up.lublin.pl), **Alan J. Lymbery** (a.lymbery@murdoch.edu.au), **Darlina Md. Naim** (darlinamdnd@usm.my), **Mahendra K. Rai** (pmkrai@hotmail.com).

**SPECIES DIVERSITY:** **Joko R. Witono** (jrwitono@yahoo.com), **Katsuhiko Kondo** (k3kondo@nodai.ac.jp), **Livia Wanntorp** (livia.wanntorp@nrm.se), **Mahesh K. Adhikari** (mkg\_adh@wlink.com.np), **Maria Panitsa** (mpanitsa@upatras.gr), **Mohib Shah** (mohibshah@awkum.edu.pk), **Paul K. Mbugua** (paulkmbugua@gmail.com), **Rasool B. Tareen** (rbtareen@yahoo.com).

**ECOSYSTEM DIVERSITY:** **Abdel Fattah N.A. Rabou** (arabou@iugaza.edu), **Alireza Ghanadi** (aghannadi@yahoo.com), **Ankur Patwardhan** (ankurpatwardhan@gmail.com), **Bambang H. Saharjo** (bhsaharjo@gmail.com), **Daiane H. Nunes** (nunesdaiane@gmail.com), **Faiza Abbasi** (faeza.abbasi@gmail.com), **Ghulam Hassan Dar** (profdar99@gmail.com), **Hassan Pourbabaie** (hassan\_pourbabaie@yahoo.com), **I Made Sudiana** (sudianai@yahoo.com), **Ivan Zambrana-Flores** (izambrana@gmail.com), **Krishna Raj** (krishnarajsec@yahoo.co.uk), **Mahdi Reyahi-Khoram** (phdmrk@gmail.com), **Mochamad A. Soendjoto** (masoendjoto@gmail.com), **Mohamed M.M. Najim** (mnajim@kln.ac.lk), **Morteza Eighani** (morteza\_eighani@yahoo.com), **Pawan K. Bharti** (gurupawanbharti@rediffmail.com), **Seweta Srivastava** (seweta.21896@lpu.co.in), **Seyed Aliakbar Hedayati** (Hedayati@gau.ac.ir), **Shahabuddin** (shahabsaleh@gmail.com), **Shahir Shamsir** (shahirshamsir@gmail.com), **Shri Kant Tripathi** (sk\_tripathi@rediffmail.com), **Stavros Lalas** (slalas@teilar.gr), **Subhash Santra** (scsantra@yahoo.com), **Sugiyarto** (sugiyarto\_ys@yahoo.com), **T.N. Prakash Kammardi** (prakashtnk@yahoo.com).

**ETHNOBIOLOGY:** **M. Jayakara Bhandary** (mbjaikar@gmail.com), **Muhammad Akram** (makram\_0451@hotmail.com).



Society for Indonesia  
Biodiversity



Sebelas Maret University  
Surakarta

Published by Smujo International for Society For Indonesia Biodiversity and Sebelas Maret University Surakarta



# BIODIVERSITAS

Journal of Biological Diversity  
Volume 20 - Number 1 - January 2018

---

<b>The sufficiency of existed protected areas in conserving the habitat of proboscis monkey (<i>Nasalis larvatus</i>)</b> WARDATUTTHOYYIBAH, SATYAWAN PUDYATMOKO, SENA ADI SUBRATA, MUHAMMAD ALI IMRON	1-10
<b>The prevalence and species of root-knot nematode which infect on potato seed in Central Java, Indonesia</b> MUTALA'LIAH, SIWI INDARTI, ARIF WIBOWO	11-16
<b>Genetic diversity and phylogenetic analysis of two Indonesian local cattle breeds based on cytochrome b gene sequences</b> TETY HARTATIK, DWI NUR HAPPY HARIYONO, YUDI ADINATA	17-22
<b>Impact of the green revolution on the gender's role in wet rice farming: A case study in Karangwangi Village, Cianjur District, West Java, Indonesia</b> RUHYAT PARTASMITA, BUDIAWATI S. ISKANDAR, SITI NURAENI, JOHAN ISKANDAR	23-36
<b>The study of suweg (<i>Amorphophallus paeoniifolius</i>) and other undergrowth species in teak plantation forest of Temengeng, Blora, Indonesia</b> SYAMSUL HIDAYAT	37-42
<b>Typology of native species as the shade tree for merbau (<i>Intsia bijuga</i>) plantations in Papua, Indonesia based on ecological species group</b> ELIESER VIKTOR SIRAMI, DJOKO MARSONO, RONGGO SADONO, MUHAMMAD ALI IMRON	43-53
<b>Taxonomy confirmation of <i>Puntius cf. binotatus</i> from Gunung Tujuh Lake, Jambi, Indonesia based on Cytochrome Oxidase-I (COI) gene</b> DEWI IMELDA ROESMA, DJONG HON TJONG, WILA KARLINA, DYTA RABBANI AIDIL	54-60
<b>Spatio-temporal distribution of microalgae producing chlorophyll and carotenoid pigments in Bali Strait, Indonesia</b> ANNA FAUZIAH, DIETRICH G BENGEN, MUJIZAT KAWAROE, HEFNI EFFENDI, MAJARIANA KRISANTI	61-67
<b>Expression of oil palm (<i>Elaeis guineensis</i>) polyisoprenoids in response to <i>Ganoderma boninense</i> infection</b> DADANG AFANDI, MOHAMMAD BASYUNI, LOLLIE AGUSTINA P. PUTRI, DIANA CHALIL, INDRA SYAHPUTRA	68-76
<b>Conflict between human and wild Sumatran Elephant (<i>Elephas maximus sumatranus</i> Temminck, 1847) in Aceh Province, Indonesia</b> INDIRA NURUL QOMARIAH, TUTIA RAHMI, ZURAIDAH SAID, ARIEF WIJAYA	77-84
<b>Short Communication: Age and growth of elongated mudskipper, <i>Pseudapocryptes elongatus</i> (Cuvier, 1816) from Sundarbans, India</b> GOPALAN MAHADEVAN, PALANIVEL BHARATHIRAJAN, VELAYUTHAM RAVI, MOJTABA POULADI, MARYAM MIRZAEI VAND KHANGHAH	85-90
<b>Bamboo diversity of Sulawesi, Indonesia</b> DITA ERVIANTI, ELIZABETH A. WIDJAJA, AGUNG SEDAYU	91-109
<b>Characteristics and causes of forest and land fires in Kapuas District, Central Kalimantan Province, Indonesia</b> ACHMAD SIDDIK THOHA, BAMBANG HERO SAHARJO, RIZALDI BOER, MUHAMMAD ARDIANSYAH	110-117

<b>Local monitoring of flowering and fruiting of Jernang, <i>Daemonorops</i> species in Sumatra, Indonesia</b> LARS SCHMIDT, HIMMAH RUSTIAMI, IDA THEILADE	118-125
<b>Qualitative morphological diversity of female Pelung Chickens in West Java, Indonesia</b> INDRAWATI Y. ASMARA, DANI GARNIDA, WIWIN TANWIRIAH, RUHYAT PARTASASMITA	126-133
<b>Growth evaluation of native tree species planted on post coal mining reclamation site in East Kalimantan, Indonesia</b> DEWI AYU LESTARI, ABBAN PUTRI FIQA, FAUZIAH, SUGENG BUDIHARTA	134-143
<b>Folic acid content and fruit characteristics of five Indonesian dessert banana cultivars</b> RITA NINGSIH, RITA MEGIA	144-151
<b>Pollen morphology of eight tribes of Asteraceae from Universitas Indonesia Campus, Depok, Indonesia</b> ANDI SALAMAH, RACHMI LUTHFIKASARI, ASTARI DWIRANTI	152-159
<b>The practice of hydrological restoration to rehabilitate abandoned shrimp ponds in Bunaken National Park, North Sulawesi, Indonesia</b> RIGNOLDA DJAMALUDDIN, BENJAMIN BROWN, ROY R. LEWIS III	160-170
<b>Relationship between environmental parameters and the plankton community of the Batuhideung Fishing Grounds, Pandeglang, Banten, Indonesia</b> NOVERITA D. TAKARINA, WAWAN NURLIANSYAH, WISNU WARDHANA	171-180
<b>The natural forest gaps maintenance diversity of understory birds in Mae Sa-Kog Ma Biosphere Reserve, northern Thailand</b> SUPALAK SIRI, YUWADEE PONPITUK, MONGKOL SAFOOWONG, DOKRAK MAROD, PRATEEP DUENGKAE	181-189
<b>Local wisdom of Talang Mamak Tribe, Riau, Indonesia in supporting sustainable bioresource utilization</b> PRIMA WAHYU TITISARI, ELFIS, IRINA SAFITRI ZEN, KHAIRANI, NADIATUL JANNA, NUNUT SUHARNI, TIKA PERMATA SARI	190-197
<b>The importance of in-situ conservation area in mining concession in preserving diversity, threatened and potential floras in East Kalimantan, Indonesia</b> ABBAN PUTRI FIQA, FAUZIAH, DEWI AYU LESTARI, SUGENG BUDIHARTA	198-210
<b>Short Communication: Further morphological evidence for separating <i>Mukia</i> Arn. from <i>Cucumis</i> L.</b> MENTARI PUTRI PRATAMI, TATIK CHIKMAWATI, RUGAYAH	211-217
<b>Short Communication: Bird species in the cement factory complex of Tarjun, South Kalimantan, Indonesia</b> MAULANA KHALID RIEFANI, MOCHAMAD ARIEF SOENDJOTO, ANDY M. MUNIR	218-225
<b>Distribution and habitat preferences of Sumatran elephant (<i>Elephas maximus sumatranus</i>) in Riau, Indonesia</b> YOHANES WISHNU SUKMANTORO, HADI S. ALIKODRA, AGUS P. KARTONO, EFRANSJAH	226-235
<b>Body size characteristics and polymorphism in GH and GHRH genes of Simeulue Buffalo of Aceh, Indonesia</b> KARTINI ERIANI, RAHMI RAHMI, IKHWAN JAMIL, ROSNIZAR ROSNIZAR, AL AZHAR	236-242
<b>Terrestrial gastropod diversity, distribution and abundance in areas with and without anthropogenic disturbances, Northeast Algeria</b> HADJER BELHIOUANI, MOHAMED EL-HADEF EL-OKKI, FATIMA-ZOHRA AFRI-MEHENNAOUI, LEILA SAHLI	243-249
<b>Short Communication: Dye-yielding plant resources of Maharashtra, India: A checklist</b> PATIL SONALI HINDURAO, KURLAPKAR DILIP DAMODAR, GAIKWAD DATTATRAYA KRISHNA	250-266

<b>Initiating the establishment of commercial stand at Tasik Besar Serkap, Riau, Indonesia: An early growth assessment of two peat swamp tree species</b> HENTI HENDALASTUTI RACHMAT, ATOK SUBIAKTO	267-272
<b>The response of East Kalimantan, Indonesia local rice cultivars against iron stress</b> NURHASANAH, HELMA SUCI LESTARI, WIDI SUNARYO	273-282
<b>Local biodiversity conservation in Sigi, Central Sulawesi, Indonesia: Analysis of the effect of elevation, land accessibility, and farmers' income and perception on vegetation diversity in agroforestry systems</b> NINA DWI LESTARI, DIDIK SUPRAYOGO, ARIEF RACHMANSYAH	283-291
<b>Short Communication: Genetic diversity among <i>Fusarium</i> isolates from cereals in Iran assessed using RAPD marker</b> SARA SIAHPOUSH, MOSTAFA DARVISHNIA	292-296
<b>Using micro-morphological fruit characters in resolving some of ambiguities in Iranian <i>Acer</i> L. (Sapindaceae) species</b> FATEMEH ASADI, FARIBA SHARIFNIA, FAHIMEH SALIMPOUR, AHMAD MAJD	297-304
<b>Comparative study of floristic diversity along altitude in the northern slope of the central Alborz Mountains, Iran</b> HALIME MORADI, FARIDEH ATTAR	305-312
<b>Presence of <i>Anisakis</i> nematode larvae in Indian mackerel (<i>Rastrelliger</i> spp.) along the Indian Ocean southern coast of East Java, Indonesia</b> EKO SETYOBUDI, IZANATUR ROHMAH, RIZKA FAUZIANA SYARIFAH, LISA RAMATIA, MURWANTOKO, DINI WAHYU KARTIKA SARI	313-319
<b>Response of polyisoprenoid concentration and profile in three groups of mangrove seedlings of coping with long-term salinity</b> MOHAMMAD BASYUNI, HIROSHI SAGAMI, SHIGEYUKI BABA, HIROSUKE OKU	320-326
<b>Morphology, palynology, and stipe anatomy of four common ferns from Pekanbaru, Riau Province, Indonesia</b> NERY SOFIYANTI, DYAH IRIANI, FITMAWATI, AFNI ATIKA MARPAUNG	327-336
<b>Short Communication: Early stability test of mutant candidates of Bangka local cassava, Indonesia</b> TRI LESTARI, ERIES DYAH MUSTIKARINI, RION APRIYADI, SAIPUL ANWAR	337-342
<b>Species diversity of Orchids in Bukit Barisan Selatan National Park, Lampung, Indonesia</b> Esti Munawaroh, Yuzammi	343-349
<b>Analysis of land use change and its relation to land potential and elephant habitat at Besitang Watershed, North Sumatra, Indonesia</b> WANDA KUSWANDA, AHMAD DANY SUNANDAR	350-358
<b>Comparison of true mangrove stands in Dudepo and Poneo Islands, North Gorontalo District, Indonesia</b> FAIZAL KASIM, MIFTAHUL KHAIR KADIM, SITTI NURSINAR, ZULKIFLI KARIM, ALDIN LAMALANGO	259-266
<b>Physiological status of high and low metabolism <i>Hevea</i> clones in the difference stage of tapping panel dryness</b> RADITE TRISTAMA, PUTRI ARISMA SIREGAR MAWADDAH, LUBIS ADE-FIPRIANI, JUNAIDI	267-273
<b>Short Communication: Exploration of bidara laut (<i>Strychnos lucida</i>) mother trees in Gunung Tunak Ecotourism Park, West Nusa Tenggara, Indonesia</b> ALI SETYAYUDI, KRISNAWATI, RYKE NANDINI	274-278
<b>The diversity of <i>Smilax</i> (Smilacaceae) in Besiq-Bermai and Bontang Forests, East Kalimantan, Indonesia</b> SITI SOFIAH, LULUT DWI SULISTYANINGSIH	279-287



**Salinity and subsequent freshwater influences on the growth, biomass, and polyisoprenoids distribution of *Rhizophora apiculata* seedlings** 288-295

MOHAMMAD BASYUNI, MASRIDA WASILAH, POPPY ANJELISA ZAITUN HASIBUAN, NURDIN SULISTİYONO, SUMARDI, YUNTHA BIMANTARA, RAHMAH HAYATI, HIROSHI SAGAMI, HIROSUKE OKU

**Population status of endemic tree kokoleceran (*Vatica bantamensis*) in Ujung Kulon National Park, Indonesia** 296-302

IYAN ROBIANSYAH, DODO, ARIEF HAMIDI

*Front cover: Nasalis larvatus male*  
(PHOTO: CHARLES J SHARP)

---

Published bimonthly

PRINTED IN INDONESIA

ISSN: 1412-033X



9 771412 033603

E-ISSN: 2085-4722



9 772085 472607

# Short Communication: Bird species in the cement factory complex of Tarjun, South Kalimantan, Indonesia

---

ORIGINALITY REPORT

---

**32%**  
SIMILARITY INDEX

**28%**  
INTERNET SOURCES

**17%**  
PUBLICATIONS

**11%**  
STUDENT PAPERS

---

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

4%  
★ [www.scribd.com](http://www.scribd.com)  
Internet Source

---

Exclude quotes    On  
Exclude bibliography    On

Exclude matches    Off