

Gastropod Density and Diversity in the Mangrove Forest of Pagatan Besar Village, Tanah Laut Regency, Indonesia

by Mochamad Arief Soendjoto

Submission date: 25-May-2023 11:08AM (UTC+0700)

Submission ID: 2101350111

File name: 019_Jurnal_Ilmu_Kelautan_24_4_179-185_Nugroho_et_al._Lengkap.pdf (930.19K)

Word count: 3983

Character count: 21272

Gastropod Density and Diversity in the Mangrove Forest of Pagatan Besar Village, Tanah Laut Regency, Indonesia

Bimo Aji Nugroho^{1*}, Mochamad Arief Soendjoto² and Muhammad Zaini¹

¹Biology Education Study Program, Lambung Mangkurat University
Jl. H. Hasan Basry, Banjarmasin, 70123, Indonesia

²Forestry Faculty, Lambung Mangkurat University
Jl. A. Yani Km. 36 Banjarbaru 70714, Indonesia

Email: bimoaji772@gmail.com

Abstract

Public awareness and knowledge on mangrove forest biodiversity, especially gastropods, is very limited. This study aims to compare the density and diversity index of gastropod species according to their distance from the coastline and to analyze the relationship of the two with the physical environment around them. Three lanes were placed in the southern part of the mangrove forest. The first lane (LT-1) was placed about 5 m from the coastline towards the sea. The second lane (LT-2) was placed 10 m towards the sea from LT-1 and the third lane (LT-3) was 10 m from LT-2 towards the sea. In each lane there were 10 (1 m x 1 m) plots and the distance between adjacent plots was 10 m. A total of 1.432 Gastropod individuals consisting of 16 species and 7 families were collected from 30 plots. In detail 16 species (927 individuals) were collected from LT-1, 16 species (500 indiv) from LT-2, and only 3 species or 5 individuals from LT-3. The density of LT-1, LT-2, and LT-3 species respectively was 92.7; 50.0; and 0.5 indiv.m⁻², while the diversity index successively was 2.56 (moderate); 2.53 (moderate); and 0.95 (low). Both species density and diversity in lanes that are close to the coastline are higher than in lanes located in the direction of the sea. The base substrate on the path that was close to the coastline was clay, while on the lane that laid in the direction of the sea was sandy loam.

Keywords: density, biodiversity, gastropod, lane, mangrove

Introduction

Pagatan Besar mangrove forest in the South Kalimantan Province has been managed sustainably. This forest is not destroyed, mangrove trees are not cut down and not converted into activities or for various other purposes. This forest has been developed by the community as an ecotourism area since 2016. Mangrove planting programs have been carried out to support the development of ecotourism areas. For example, muddy beaches in the southern part of the forest are planted with *Avicennia marina* up to 95%. Other species that are also planted are *Rhizophora* and *Bruguiera*.

In regard to sustainable management, the development of the Pagatan Besar Mangrove Forest needs to be socialized to the community (Soendjoto, 2003). For example, regarding the size of the forest area (Riefani et al., 2019), as well as biodiversity within the forest, i.e. plants (Soendjoto and Arifin, 1999), and animals (Riefani and Arsyad, 2019) including marine organisms, such as gastropods.

Several reports showed that gastropods can be found in mangrove forests (Pribadi et al. 2009; Ernanto et al., 2010; Darmi et al., 2017). This class of animals generally functions ecologically (Setyono, 2006). In aquatic and terrestrial ecosystems, Gastropods are known as food for echinoderms, fish, birds and mammals (Pawar and Al-Tawaha, 2017). On the other hand, they play a role of mechanically breaking down (decomposition) organic matter, such as leaves through their feeding activities (Kabir et al., 2014; Tavares et al., 2015). Some gastropod are even of economic value to the community. Meat of the genus *Strombus*, for example, can be consumed (Suwignyo, 2005; Rosady et al., 2016) and the shell can be used as a raw material for decoration (Supusepa, 2018). The genus *Cymatium*, *Haliotis*, *Monodonta*, *Tectus*, *Trochus*, *Turbo*, *Vasum* and *Vexillum* can also be used as food (Supusepa, 2018).

Knowledge about gastropoda population is needed in maintaining sustainable management of mangrove forest in the Pagatan Besar Mangrove Forest. This study aims to compare the density index and diversity of Gastropod species according to their

*) Corresponding author
© Ilmu Kelautan, UNDIP

distance from the coastline and analyze the relationship between the two and their physical environment. This finding can also be used to increase public awareness about the sustainability of mangrove forest ecotourism areas.

Materials and Methods

Gastropod samples were taken in March-April 2018 from three lanes placed in the southern part of the Pagatan Besar Mangrove Forest, Takisung District, Tanah Laut Regency. Riefani et al. (2019) mentions two parts of the Pagatan Besar Mangrove Forest, namely the northern and southern parts. The northern part which is the estimated area of 5.97 ha stretches along 825 m starting from the mouth of the Pagatan Besar River (Tabanio) to the south to the first ulin walkpath. According to Soendjoto (2003), the northern part of the forest naturally formed after a tidal wave in 1985. The southern part, which covers about 4.72 ha, stretches 587 m from the first ulin walkpath to south to the fourth walkpath or muddy area that has no mangrove plants. The southern forest is a concentration of the development of ecotourism area. The muddy area is an area for planting mangrove trees by the communities.

The three lanes were extended from north to south. The first lane (LT-1) was placed about 5 m from the coastline towards the sea. The coastline is a virtual line that is indicated by land covered with grass or sandy land which in daily tides is considered the highest tidal boundary; in certain cases, for example *rob*, this land may be inundated with sea water. The second lane (LT-2) was placed 10 m towards the sea from LT-1 and the third lane (LT-3)

was 10 m towards the sea from LT-2 (Figure 1.). In each lane was placed 10 (1m x 1m) plots and the closest inter-plot distance was 10 m.

From each plot Gastropod individuals were collected and grouped according to species. Species were identified according to Nybakken and Bertness (2005), and species diversity index was done with Shannon-Wiener index. Abiotic factors consisting of water temperature, salinity, pH of the substrate, and type of substrate were measured or sampled from 1 plot per lane.

Results and Discussion

Gastropods (n=1.432) found at Pagatan Besar mangrove forest consist of 16 species or 7 families (Figure 2.) with density and diversity as presented in Table 1. The number of these species is less than those found by Siwi (2017) on the Beach Si Runtoh, Baluran National Park, East Java (19 species); Supusepa (2018) in Negeri Tial and Negeri Suli, Maluku (30 and 39 species, respectively); or even Rumahlatu and Leiwakabessy (2017) in the coastal waters of Ambon Island, Maluku (65 species). However, the number of species in the forest is still greater than that reported by Ernanto et al. (2010) in Muara Sungai Batang, South Sumatra (10 species), Isnaningsih and Patria (2018) in Tanjung Lesung, Banten (7 species), Maturbongs and Elviana (2016) in Kambapi Coast, Papua (10 species), Nurrudin et al. (2015) in Tanjung Jabung Barat, Jambi (15 species), Rangan (2010) in Minahasa, North Sulawesi (9 species), and Puryono and Suryanti (2019) in Pemalang, Central Java (8 species).



Figure 1. The placement of line transects on the Pagatan Besar mangrove forest

Tabel 1. Diversity of Gastropod in the Pagatan Besar mangrove forest

Family and spesies name	LT-1		LT-2		LT-3	
	n	K	n	K	n	K
A. Potamididae						
1. <i>Pirenella retifera</i>	148	14.8	69	6.9	0	0
2. <i>Tectonatica pusilla</i>	92	9.2	21	2.1	0	0
3. <i>Nassarius stolatus</i>	71	7.1	15	1.5	0	0
4. <i>Cerithidea cingulata</i>	134	13.4	78	7.8	0	0
5. <i>Littoraria melanostoma</i>	64	6.4	52	5.2	3	0.3
6. <i>Vexillum rugosum</i>	28	2.8	17	1.7	0	0
7. <i>Cerithideopsilla alata</i>	38	3.8	21	2.1	1	0.1
B. Naticidae						
8. <i>Cerithidea obtusa</i>	39	3.9	8	0.8	0	0
C. Nassariidae						
9. <i>Cassidula aurisfelis</i>	30	3.0	34	3.4	0	0
D. Littorinidae						
10. <i>Cassidula aurisfelis</i>	21	2.1	17	1.7	0	0
11. <i>Cassidula nucleus</i>	10	1.0	1	0.1	0	0
E. Costellariidae						
12. <i>Littoraria cingulate</i>	12	1.2	9	0.9	0	0
F. Ellobiidae						
13. <i>Neritina cornucopia</i>	42	4.2	27	2.7	0	0
14. <i>Neritina violacea</i>	67	6.7	49	4.9	1	0.1
G. Neritidae						
15. <i>Cerithidea decollata</i>	78	7.8	41	4.1	0	0
16. <i>Cerithidea alata</i>	53	5.3	41	4.1	0	0
The number of individuals (N)	927	-	500	-	5	-
The number of species (S)	16	-	16	-	3	-
Density of all species (K)	-	92.7	-	50.0	-	0.5
Diversity Index (H')	2.56	-	2.53	-	0.95	-

Remark: LT = line transect; K = density (individu.m⁻²)

Species density at LT-1 is higher than LT-2 density and so subsequently density at LT-2 compared to LT-3. From Figure 3, the differences in LT-1 and LT-2 and LT-3 species densities are even striking. All species were found in LT-1 and LT-2, but only three species were found in LT-3, namely *Littoraria melanostoma*, *Cerithideopsilla alata*, and *Neritina violacea*. The same thing happens with the species diversity index. The diversity index on LT-1 is higher than that of LT-2 and the diversity index on LT-2 is higher than that of LT-3. The qualitative criteria for LT-1 diversity index are the same as that of LT-2 (moderate). There is only LT-3 having different criteria, low diversity. From these two conditions, the results showed that both Gastropod species density and diversity index were higher in the lane that was closer to the land than the lane that was further from the land or closer to the sea. In other words, the more toward the sea, the lower the density. The same thing happened to the diversity index Gastropoda. These results were obtained under conditions of water temperature of 27–31°C, salinity of 28–31 ppm, pH of substrate of 5.8, 6.2, as well as bottom substrate types of clay and sandy loam (Table 2.).

The difference in density and diversity index of Gastropods from each lane is thought to be caused

by differences in organic materials—in terms of the amount and diversity—contained in the substrate or scattered on the floor of mangrove forests. Organic material which is basically a source of food for Gastropods comes from parts of plants that fall in the forest or are carried by ocean / river currents and then accumulate in the mangrove forest. Leaves are an organic material favored by gastropods *Melampus coffeus* because they contain little lignocellulose (Tavares et al., 2015). Gastropod abundance is positively affected by the content of organic matter in sediments. but negatively by the content of organic matter in seawater in the mangrove ecosystems of Payum. Merauke waters (Merly and Elviana. 2017).

There are no publications regarding the number and species of plants in the southern part of the Pagatan Besar Mangrove Forest. but a preliminary survey shows that this forest is dominated by api-api (*Avicennia marina*) which are visually 95% still at a sapling level (diameter at breast height less than 10 cm). In the northern forest. the dominance of api-api was reported by Soendjoto and Arifin (1999). In addition to api-api. plant species in the northern forest consisted of bakau (*Rhizophora mucronata*). buta-buta (*Excoecaria agallocha*). nyiri batu (*Xylocarpus*




Figure 2. Sixteen Gastropod species found in the Pagatan Besar mangrove forest, Tanah Laut Regency, Indonesia

granatum). rambai padi (*Sonneratia caseolaris*), rambai bogam (*S. alba*), and 4 species of non-woody plants. As a comparison, the amount of organic matter in litter in the Sepi Bay mangrove forest, Lombok reaches 9.9 tons.ha⁻¹.y⁻¹ (Zamroni and Rohyani, 2008).

Organic material is not only from plants, but can also come from parts of animals died directly in the forest or are carried by ocean / river currents. There is no data on the number and species of animals in this condition. However, 60 species of birds (Riefani and Arsyad, 2019) and 4 mudskipper species (Soendjoto, 2019) use the southern forest as their habitat.

Factors affecting Gastropods are not only organic material or biotic components, but also abiotic components or physical environment conditions. Gastropod diversity not only involves food sources, competition and competition, but also the environment (Dahuri *et al.*, 2008). There is a positive correlation between Gastropod and physical-chemical factors (temperature, salinity, pH, DO) in

the coastal waters of Ambon Island (Rumahlatu and Leiwakabessy, 2017). The number of Gastropod species and individuals in sandy clay substrate is greater than those in sand substrate (Riniatsih and Kushartono, 2009). Gastropods are more commonly found in clay substrate compared to sandy (Rahmasari, 2015).

Sludge substrate is easier to trap or bind organic material as a food source for Gastropods than sand substrate. Sand has greater porosity so that the organic material in it is more easily moved or transported by natural facilities which in the case of mangrove forests are in the form of tides or river water flow. The condition of the substrate is also influenced by the movement of sea or river water that carries certain temperatures, salinity, and pH.

In addition, the condition of the substrate is influenced by sunlight penetration in mangrove forests. Substrates in thick or lush forests are less exposed to light penetration than that in thin or non-forested forests. Thin or non-forested forest refers to lanes far from land or close to the sea.

Table 2. Environment condition of Pagatan Besar mangrove forest during the collecting of Gastropoda samples

Parameter	LT-1	LT-2	LT-3	Comparison
Water temperature (°C)	27	28	31	23.7–30 ¹ ; 28–29 ² ; 26–27 ³ ; 29.9–31.4 ⁴
Salinity (ppm)	28	31	30	26–31 ¹ ; 33 ² ; 29–32 ³ ; 30.31–32.02 ⁴
pH substrate	5.8	6.0	6.2	3–7 ¹ ; 5.8–6 ²
Bottom substrate type	Clay	Sandy loam	Sandy loam	Sandy loam ³

Remark: 1. Merly and Elviana (2017); 2. Rahmasari *et al.* (2015); 3. Riniatsih and Kushartono (2009); 4. Rumahlatu dan Leiwakabessy (2017)




Figure 3. Density and diversity index of gastropods on three line transects in the research area

Conclusion

Gastropods obtained from the Pagatan Besar Mangrove Forest (southern part) consist of 16 species and 7 families. The density and diversity of species in lanes that are close to the coastline (whose substrate is clay) is higher than those close to the sea (whose substrate is sandy loam).

References

- Dahuri, R., Rais, J., Ginting, S.P. & Sitepu, M.J., 2008. Pengelolaan Sumberdaya Pesisir dan Lautan Secara Terpadu. PT. Pradnya Paramita. Jakarta. 326 pages.
- Darmi, Setyawati, T.R. & Yanti, A.H., 2017. Jenis-jenis Gastropoda di Kawasan Hutan Mangrove Muara Sungai Kuala Baru Kecamatan Jawai Kabupaten Sambas. *Protobiont.*, 6(1):29-34.
- Ernanto, R., Agustriani, F. & Aryawati, R., 2010. Struktur komunitas Gastropoda pada ekosistem mangrove di Muara Sungai Batang Ogan Komering Ilir Sumatera Selatan. *Maspari J.*, 01:73-78.
- Isnainingsih, N.R. & Patria, M.P., 2018. Peran komunitas Moluska dalam mendukung fungsi kawasan mangrove di Tanjung Lesung, Pandeglang, Banten. *J. Biotropika*, 6(2): 35-44.
- Kabir, M., Abolfathi, M., Hajimoradloo, A., Zahedi, S., Kathiresan, K., & Goli, S., 2014. Effect of mangroves on distribution, diversity and abundance of molluscs in mangrove ecosystem: a review. *AACL Bioflux*, 7(4): 286-300.
- Maturbongs, M.R. & Elviana, S., 2016. Komposisi, kepadatan dan keanekaragaman jenis Gastropoda di Kawasan Mangrove Pesisir Pantai Kambapi pada musim peralihan I. *J. Ilmiah Agribisnis Perikan.*, 9(2): 19-23. doi : 10.29239/j.agrikan.9.2.19-23
- Merly, S.L. & Elviana, S., 2017. Korelasi sebaran Gastropoda dan bahan organik dasar pada ekosistem mangrove di Perairan Pantai Payum, Merauke. *Dinamika Maritim*, 6(1): 18-22.
- Nurrudin, Hamidah, A. & Kartika, W.D., 2015. Keanekaragaman jenis Gastropoda di sekitar Tempat Pelelangan Ikan (TPI) Parit 7 Desa Tungkal I Tanjung Jabung Barat. *Biospecies*, 8(2): 51-60.
- Nybakken, J.W. & Bertness, M.D., 2005. *Marine Biology an Ecological Approach*, 6th edition. San Francisco: Pearson Education, Inc.
- Pawar, P.R. & Al-Tawaha, A.R.M.S., 2017. Biodiversity of marine gastropods along the Uran coast, Navi Mumbai, west coast of India. *American-Eurasian J. Sustain. Agricul.*, 11(2): 19-30.
- Pribadi, R., Hartati, R. & Suryono, C.A., 2009. Komposisi jenis dan distribusi Gastropoda di Kawasan Hutan Mangrove Segara Anakan Cilacap. *Ilmu Kelautan*, 14(2):102-111. doi : 10.14710/ik.jkms.14.2.102-111
- Puryono, S. & Suryanti, S., 2019. Gastropod diversity in Mangrove Forests of Mojo Village, Ulujami District, Pemalang Regency, Indonesia. *J. Ecol. Eng.*, 20(1): 165-173. doi : 10.12911/22998993/93940
- Rahmasari, T., Purnomo, T. & Ambarwati, R. 2015. Keanekaragaman dan kelimpahan Gastropoda di Pantai Selatan Kabupaten Pamekasan. *Biosaintifika*, 7(1):48-54. doi: 10.15294/biosaintifika.v7i1.3535
- Rangan, J.K., 2010. Inventarisasi Gastropoda di lantai hutan mangrove Desa Rap-Rap Kabupaten Minahasa Selatan Sulawesi Utara. *J. Perikan. Kel.*, 6(1): 63-66. doi: 10.35800/jpkt.6.1.2010.163
- Riefani, M.K. & Arsyad, M. 2019. Spesies burung di Kawasan Ekowisata Mangrof Pagatan Besar, Kabupaten Tanah Laut, Indonesia. *Pros. Sem. Nas. Lingkungan Lahan Basah*, 4(1): 192-196.
- Riefani, Mahrudin, M.K. & Soendjoto, M.A., 2019. Pemberdayaan masyarakat untuk melestarikan Kawasan Ekowisata Mangrove Desa Pagatan Besar, Kabupaten Tanah Laut. *Pro Sejahtera*, 1: 121-126.
- Riniatsih, I. & Kushartono, E.W., 2009. Substrat dasar dan parameter oseanografi sebagai penentu keberadaan Gastropoda dan Bivalvia di Pantai Sluke Kabupaten Rembang. *Ilmu Kelautan*, 14(1): 50-59.
- Rosady, V.P., Astuty, S. & Prihadi, D.J., 2016. Kelimpahan dan kondisi habitat siput gonggong (*Strombus turturella*) di Pesisir Kabupaten Bintan, Kepulauan Riau. *J. Perikan. Kel.*, 7(2): 35-44.
- Rumahlatu, D. & Leiwakabessy, F., 2017. Biodiversity of gastropoda in the coastal waters of Ambon Island, Indonesia. *AACL Bioflux*, 10(2): 285-296
- Setyono, D.E.D., 2006. Karakteristik biologi dan produk kekerangan laut. *Oseana*, 31(1): 1- 7.

- Siwi, F.R, Sudarmadji & Suratno, 2017. Keanekaragaman dan kepadatan Gastropoda di Hutan Mangrove Pantai Si Runtoh Taman Nasional Baluran. *J. Ilmu Dasar*, 18(2):119-124.
- Soendjoto, M.A., 2003. Hutan mangrove Pegatan Besar: Hikmah dari sebuah musibah. *Warta Konservasi Lahan Basah*, 11(2): 26-27.
- Soendjoto, M.A., 2019. Potential and problems in development of the ecotourism area (Case in the Pagatan Besar Mangrove Forest, Tanah Laut Regency, Indonesia). *Pros. Sem. Nas. Lingkungan Lahan Basah*, 4(3): 634-641.
- Soendjoto, M.A. dan P. Arifin., 1999. Hutan mangrove Desa Pegatan Besar, Kalimantan Selatan: Vegetasi dan manfaatnya bagi masyarakat. *Manusia dan Lingkungan*, 6(17): 42-51.
- Supusepa, J., 2018. Inventaris jenis dan potensi Gastropoda di Negeri Suli dan Negeri Tial. *J. Triton*, 14(1): 28-34.
- Suwignyo, S., Widigdo, B., Wardiatno, Y. & Krisanti, M., 2005. *Avertebrata Air. Jilid 1*. Penebar Swadaya, Jakarta.
- Tavares, D.S., Maia, R.C., Rocha-Barreira, C. & Matthews-Cascon, H., 2015. Ecological relations between mangrove leaf litter and the spatial distribution of the gastropod *Melampus coffeus* in a fringe mangrove forest. *Iheringia, Sér. Zool.*, 105(1): 35-40. doi: 10.1590/1678-4766 201510513540.
- Zamroni, Y., I.S. Rohyani., 2008. Produksi serasah hutan mangrove di Perairan Pantai Teluk Sepi, Lombok Barat. *Biodiversitas*, 9(4): 284-287. doi: 10.13057/biodiv/d090409.

Index Edition

ISSN 0853 - 7291

Volume 24 No. 4 December 2019

ILMU KELAUTAN

Indonesian Journal of Marine Sciences



IK Vol. 24 No. 4 Page 153 - 194 Semarang, December 2019 ISSN

IK	Vol. 24	No. 4	Page 153 - 194	Semarang, December 2019	ISSN 0853 - 7291
----	---------	-------	----------------	-------------------------	------------------



THE JOURNAL IS JOINTLY PUBLISHED BY:
MARINE SCIENCE DEPARTMENT, FPIK, DIPONEGORO UNIVERSITY, &
ASSOCIATION OF INDOONESIAN COASTAL MANAGEMENT EXPERTS
Accredited DIKTI : 12/M/KP/II/2015




Editorial Team

Editor In Chief














• **Prof. Dr. Ambariyanto Ambariyanto** (ScopusID: [7409724866](#))
Faculty of Fisheries and Marine Science, Diponegoro University, Indonesia

Associate (Handling) Editor

- **Dr. Ir. Retno Hartati, M.Sc** (ScopusID: [57193736954](#))
 Marine Science Study Program, Faculty of Fisheries and Marine Sciences, Diponegoro University, Indonesia
- **Dr. Ir. Widianingsih M.Sc** (ScopusID: [57193740093](#))
 Marine Science Study Program, Faculty of Fisheries and Marine Sciences, Diponegoro University, Indonesia
- **Prof. Dr. Ir. Delianis Pringgenis, M.Sc** (ScopusID: [24290899400](#))
 Marine Science Study Program, Faculty of Fisheries and Marine Sciences, Diponegoro University, Indonesia
- **Dr. Ir. Bambang Yulianto, DEA** (ScopusID: [57191832540](#))
Marine Science Study Program, Faculty of Fisheries and Marine Sciences, Diponegoro University, Semarang, Indonesia
- **Dr. Ir. Chrisna Adhi Suryono, M.Phill** (ScopusID: [55939217500](#))
Marine Science Study Program, Faculty of Fisheries and Marine Sciences, Diponegoro University, Indonesia
- **Ir. Ali Djunaedi, M.Phill** (ScopusID: [7801315575](#))
Marine Science Study Program, Faculty of Fisheries and Marine Sciences, Diponegoro University, Semarang, Indonesia
- **Robertus Triaji Mahendrajaya, S.Kel. M.Si** (ScopusID: [57201186750](#))
 Marine Science Study Program, Faculty of Fisheries and Marine Sciences, Diponegoro University, Indonesia

International Editorial Board

- **Prof. Ian M Dutton** (ScopusID: [6602374559](#))
 School of Fisheries and Ocean Sciences, University of Alaska Fairbanks, United States
- **Prof. Ove Hoegh-Guldberg** (ScopusID: [26643606100](#))
 Global Change Institute, Queensland University, Brisbane, Australia., Australia
- **Prof. Yasuhiro Igarashi** (ScopusID: [7401635211](#))
 Biotechnology Research Center and Department of Biotechnology, Toyama Prefectural University, Japan
- **Prof. Hamid Talib Al-Saad** (ScopusID: [6603939687](#))
 Department of Marine Chemistry and Pollution, University of Basrah, Iraq

- **Craig J. Starger, PhD** (ScopusID: [6506597568](#))
 American Association for the Advancement of Science, US AID, Washington DC 20523, United States
- **Prof. Agus Sabdono, PhD.** (ScopusID: [16234515500](#))
 Marine Science Study Program, Faculty of Fisheries and Marine Sciences, Diponegoro University, Indonesia
- **Prof. Agoes Soegiarto PhD** (ScopusID: [6506295541](#))
 Biology Department, Faculty of Science and Technology, Universitas Airlangga. Surabaya Indonesia., Indonesia
- **Prof. Ocky Karna Radjasa, PhD** (ScopusID: [15824260100](#))
 Research and Community Services Institutes Diponegoro University, Indonesia
- **Prof. Dr. Ir. Indra Jaya, MSc** (ScopusID: [26039180400](#))
 Faculty of Fisheries and Marine Science, Bogor Agriculture Institute, Indonesia
- **Prof. Dr. Diah Permata Wijayanti, MSc.** (ScopusID: [56315616200](#))
 Coral Reef Research Center, Diponegoro University, Indonesia
- **Prof. Dr. Feliatra DEA** (ScopusID: [6506798004](#))
 Faculty of Fisheries and Marine Science, University of Riau, Indonesia
- **Agus Trianto, ST, MSc, PhD.** (ScopusID: [9432351700](#))
 Marine Science Study Program, Faculty of Fisheries and Marine Sciences, Diponegoro University, Indonesia

Editorial Office

- **Ir. Ita Riniatsih, M.Sc** (ScopusID: [57201188055](#))
Marine Science Study Program, Faculty of Fisheries and Marine Sciences, Diponegoro University, Semarang, Indonesi
-

Vol 24, No 4 (2019): Ilmu Kelautan

Table of Contents

Research Articles

- [Microplastic in The Bali Strait: Comparison of Two Sampling Methods](#) 153-158
Defri Yona, Zefanya Nandaningtyas, Bernads Daniel Marolop Siagian, Syarifah Hikmah
Julinda Sari, Agung Yunanto, Feni Iranawati, Mochamad Arif Zainul Fuad, Junika
Chintia Ayu Putri, Mela Dita Maharani
| Language: [EN](#) | DOI: [10.14710/ik.ijms.24.4.153-158](https://doi.org/10.14710/ik.ijms.24.4.153-158)
-
- [Oxygen Consumption Rate of Polychaeta Nereis sp. Different Sizes and Type of Feed](#) 159-163
Eko Setio Wibowo, Endah Sri Palupi, I G A Ayu Ratna Puspitasari, Atang Atang
| Language: [EN](#) | DOI: [10.14710/ik.ijms.24.4.159-163](https://doi.org/10.14710/ik.ijms.24.4.159-163)
- [Statistical multi-metric based eutrophication index: Case of study at Batam Marine Reserve Park](#) 164-170
Hedi Indra Januar, Asri Pratitis, Rini Susilowati
| Language: [EN](#) | DOI: [10.14710/ik.ijms.24.4.164-170](https://doi.org/10.14710/ik.ijms.24.4.164-170)
- [Parasites Identification and Histopathology Changes on Blood Cooke \(Anadara granosa Linnaeus, 1758\)](#) 171-178
Yuni Karnisa, Desrina Desrina, Ita Widowati
| Language: [EN](#) | DOI: [10.14710/ik.ijms.24.4.171-178](https://doi.org/10.14710/ik.ijms.24.4.171-178)
- [Gastropod Density and Diversity in the Mangrove Forest of Pagatan Besar Village, Tanah Laut Regency, Indonesia](#) 179-185
Bimo Aji Nugroho, Mochamad Arief Soendjoto, Muhammad Zaini
| Language: [EN](#) | DOI: [10.14710/ik.ijms.24.4.179-185](https://doi.org/10.14710/ik.ijms.24.4.179-185)
- [Biological Parameters of Blacksaddled Coralgrouper \(Plectropomus laevis Lacepède, 1801\) Eggs and Larvae](#) 186-194
Regina Melianawati, Bejo Slamet
| Language: [EN](#) | DOI: [10.14710/ik.ijms.24.4.186-194](https://doi.org/10.14710/ik.ijms.24.4.186-194)
-

Gastropod Density and Diversity in the Mangrove Forest of Pagatan Besar Village, Tanah Laut Regency, Indonesia

ORIGINALITY REPORT

13%
SIMILARITY INDEX

12%
INTERNET SOURCES

8%
PUBLICATIONS

4%
STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

5%
★ **smujo.id**
Internet Source

Exclude quotes On
Exclude bibliography On

Exclude matches Off