

Grain diversity and cultivation of Indonesian swamp rice germplasm: building the foundation for an ex-situ conservation programme

by Dindin Hidayatul Mursyidin

Submission date: 31-May-2023 03:17AM (UTC-0500)

Submission ID: 2105864842

File name: 10-Grain_diversity_and_cultivation.pdf (2.7M)

Word count: 8581

Character count: 43836

Grain diversity and cultivation of Indonesian swamp rice germplasm: building the foundation for an ex-situ conservation programme

Dindin H. Mursyidin¹, Izhar Khairullah², Muhammad Saleh²

11

¹Laboratory of Genetics and Molecular Biology, University of Lambung Mangkurat, South Kalimantan, Indonesia

²National Research and Innovation Agency, Republic of Indonesia

SUMMARY

Tropical swamps are ecosystems with high genetic resources amongst plants, animals and microbes that are also in some demand as agricultural land. Traditional rice cultivars, known as swamp rice, are very interesting in this context, mainly because of their high adaptability to extreme local conditions such as waterlogging, low pH, heavy metal poisoning and salinity. We have collected 107 accessions of swamp rice from seven Indonesian provinces across the two large islands of Sumatra and Kalimantan. In this study we aimed to determine the range of grain trait diversity amongst this swamp rice germplasm, and to cultivate the different forms under managed conditions at the Research Station of the Indonesian Swamp Agricultural Research Institute (ISARI) South Kalimantan. The germplasm was classified into six categories on the basis of grain shape, i.e., very long-slender, very long-intermediate, long-slender, long-intermediate, medium-slender and medium-intermediate. The very long-slender class, identified as the *indica* subspecies, was dominant (71 %). The germplasm was planted successfully and grew well, with most plants >100 cm tall but showing height differences between different accessions. The accession *Ketan* from Lampung was tallest and *Betek* from South Sumatra was shortest. Our study provides an initial foundation for a future ex-situ conservation effort for swamp rice in Indonesia.

KEY WORDS: genetic diversity, landrace, *Oryza sativa*, peat swamp

INTRODUCTION

The principal distinguishing characteristic of tropical swamp ecosystems is the seasonally recurrent shallow waterlogging that leads to swampy areas being known as saturated land (Kennish 2016, Trettin *et al.* 2019). These swamps are also notable for hosting high numbers of endemic species of flora, fauna and microbes (Mitsch 2013, Giesen *et al.* 2018, Harrison *et al.* 2020) and performing numerous beneficial ecosystem functions. In general, they act as giant reservoirs that store floodwater and maintain surface water flow during dry periods, whilst filtering or purifying the water and thus improving water quality (Harrison *et al.* 2020). Furthermore, swamps provide opportunities for economic benefits to, for example, the industrial, agricultural and tourism sectors (Page *et al.* 2012, Harrison *et al.* 2020).

Although swampland is regarded as marginal land for agriculture, it is currently used extensively for this purpose in efforts to feed the world's rapidly growing human population (Sulaiman *et al.* 2019, Qurani & Lakitan 2021). For example, in Asia, over 200 million ha of swampland are utilised for agricultural purposes (Wang 2020), including 20.7 million ha of

tidal and 13.3 million ha of inland swamps in Indonesia (Gunawan *et al.* 2012, Mursyidin *et al.* 2019, Qurani & Lakitan 2021). These areas are mostly located on the large islands of Sumatra and Kalimantan (Gunawan *et al.* 2012, Sulakhudin & Hatta 2018).

The traditional rice cultivar known as swamp rice is very interesting in the context of the inherent value of natural swamps combined with the need to utilise them for agriculture. Swamp rice germplasm has developed over a long period under the intensively interacting influences of local environmental adaptation and the selection of desirable traits by farmers (Sanghera *et al.* 2013, Mursyidin *et al.* 2017). Consequently, swamp rice exhibits some agronomic traits that might be beneficially utilised in conservation and breeding programs aiming to improve crop yield stability, stress tolerance, adaptability to local conditions, etc. (Aziz *et al.* 2018, Mursyidin *et al.* 2019); for example, tolerance to acidity, salinity and metal contamination (Rao *et al.* 2018, Mursyidin *et al.* 2021).

Subspecies of Indonesian swamp rice are phenotypically distinguished on the basis of grain shape. Historically, the *indica* subspecies moved



from Sumatra to Kalimantan (Borneo) about 300 years ago (Kiple & Ornelas 2000). On the other hand, *javanica* rice comes from Java Island, and continues to change genetically through domestication. Hence, it seems that the islands of Sumatra and Kalimantan provide an important route for the domestication and distribution of traditional rice accessions worldwide (Mursyidin *et al.* 2017). According to Calingacion *et al.* (2014), grain shape is also fundamental to developing new rice cultivars for swamp areas within Indonesia because local farmers and consumers have strong preferences with regard to this trait. For example, most farmers in South Kalimantan Province cultivate and consume medium-sized rice grain, e.g. the so-called *Siam* group.

With the increasing adoption of high-yielding ‘green revolution’ cultivars, the traditional swamp rice germplasm has begun to disappear (Nourollah 2016). Consequently, urgent action to preserve and conserve this germplasm is now needed. In general, the main goal of conservation is to ensure the sustainability of species, habitats and biological communities, including interactions amongst species or between species and their environments (Offord 2016, Malhotra *et al.* 2018). According to Offord (2016), there are three possible conservation strategies, namely ex-situ, in-situ, and on-farm. Ex-situ conservation involves collecting samples from the place of origin and storing them somewhere. For

example, the International Rice Research Institute (IRRI) has established a gene bank for rice, by collecting and storing dried rice seeds in preservation tanks at a temperature of -20 °C (Jackson 2016). While this kind of conservation has certain limitations, it is relatively safe and cost efficient when implemented for some plant species (Galetti 2018, Malhotra *et al.* 2018).

For Indonesia, we have collected more than one hundred accessions of swamp rice (*Oryza sativa* L.) from different parts of the country (Mursyidin *et al.* 2017). The objective of this study was to characterise this germplasm morphologically on the basis of grain shape (diversity) and cultivate it under managed conditions at a location distant from its origin. Thus, we aimed to establish an initial foundation for a future ex-situ conservation effort for swamp rice in Indonesia.

METHODS

This study encompassed 107 accessions of swamp rice collected from tidal and inland swamps across seven regions (provinces) of Indonesia (Figure 1). Table A1 in the Appendix catalogues the complete collection of swamp rice accessions, including their origins and growing habitats. This seed was planted, cultivated and harvested in May–October 2020 at the



11 Figure 1. Map of Indonesia, showing the seven regions (provinces) where the 107 swamp rice accessions employed in this study were collected. Detailed information about the accessions (names, habitats) is listed in Table A1 in the Appendix.

6 research station of the Indonesian Swampland Agriculture Research Institute (ISARI) in Tanjung Harapan Village (Alalak District, Barito Kuala Regency, South Kalimantan Province; Figure 2). The land to be planted was prepared using a hoe and hand tractor to loosen the soil, then flooded with water twice a month to prevent weeds from growing. Seeds were sown in a moist (but not saturated) seedbed using the 'tugal' method, which involves making holes ± 5 cm deep using a galam (*Melaleuca leucadendron*, a typical plant of swamps) stick with a pointed tip. The rice grain (20–25 seeds per accession) was placed into the seedbed holes, then the seedbed was covered with soil plus a layer of reed leaves to deter predators, especially birds, and kept moist by regular watering. After sprouting, the seedlings were transplanted to the edge of the planting site using a hoe and cleaver; then, when they were about four weeks old, they were planted out in a pattern of two rows, 2.5 m long, for each cultivar (see Figure 5 later). The plants were maintained by fertilising with urea (90 kg ha⁻¹) and the NPK fertiliser 'PHONSKA' (150 kg ha⁻¹) one week after planting out, and weeding (removing weeds growing around each plant) four weeks after fertiliser application. The rice field was protected from rats and birds by plastic fences and para-nets. After harvesting, the grain of each rice accession was measured and determined using the Khan *et al.* (2009) criteria (Table 2), then classified as subspecies *indica*, *javanica* or *japonica* (Zhang *et al.* 2011). Plant height was also measured, as an additional rice descriptor (Tran *et al.* 2012).

RESULTS

Grain diversity

Based on initial observation, our accessions of Indonesian swamp rice were classified into the *indica* and *javanica* subspecies plus an intermediate form. We further classified the latter into three categories, i.e., close to one of the subspecies *indica*, *javanica* or *japonica* (Figure 3). Interestingly, our morphological observations identified five accessions of *indica* with an awn on the grain tip (E in Figure 3). Further analysis revealed that the germplasm encompassed six shape classes, i.e., very long-slender, very long-intermediate, long-slender, long-intermediate, medium-slender, and medium-intermediate (Table A2). Figure 4A shows this grouping more clearly, and that the very long-slender *indica* subspecies was dominant (71 %). Within the intermediate form, the category 'close to *japonica*' was more frequent (68 %) than the two others (Figure 4B).



Figure 2. Map showing the location of the ex-situ conservation site for Indonesian swamp rice germplasm at the ISARI Research Station (red circle; 3° 10' 15.77" S, 114° 36' 12.34" E).

Table 2. Morphological classification of rice grain into four categories based on its relative length (L) and width (W). Source: Khan *et al.* (2009).

Grain size	Length (mm)	Grain shape	Grain ratio (L/W)
very long	>7.50	slender	>3.00
long	6.61–7.50	intermediate	2.10–3.00
medium	5.50–6.60	bold	1.10–2.00
short	≤ 5.50	round	<1.00

Rice cultivation

Generally, the Indonesian swamp rice germplasm was planted successfully and grew well, although some plants did not grow optimally (Figure 5). Figure 6 shows the variation in plant height. The plants grown from most of the swamp rice accessions reached >100 cm in height, the tallest being *Ketan* (from Lampung) and the shortest *Betek* (from South Sumatra).






Figure 3. Grain diversity of Indonesian swamp rice germplasm, showing differences among accessions. The labels A–F indicates the subspecies: A = *indica*; B = *javanica*; C = close to *indica*; D = close to *indica*; E = *indica* with awn on grain tip; F = close to *japonica*. The name of each accession is provided in Table A1.




Figure 4. Percentage of Indonesian swamp rice germplasm accessions based on grain shape (A) and sub-species (B).

DISCUSSION

Grain diversity

On the basis of grain shape, Indonesian swamp rice shows distinctive characteristics (Figure 3, Table A2) and is dominated by the *indica* subspecies (Figure 4). Genetically, the shape of the rice grain is determined by several traits, e.g., weight, length, width and thickness (Zheng *et al.* 2015). At molecular level, several genes (QTL), such as *Dwarf1 (D1)*, *GS3* and *GW2*, control this trait (Fu *et al.* 2015, Zhou *et al.* 2015). Zhou *et al.* (2015) reported that *GS3* affects the grain length and weight, whereas *GW2* is related to grain width and weight.

Regardless of the characteristics of the grain studied, information on genetic diversity is essential for plant genetic conservation, preventing the genetic erosion of breeding populations, and selecting superior parents for plant breeding (Wu *et al.* 2021). In other words, studies of genetic diversity facilitate an understanding of the relationships between accessions and help towards identifying

redundancies or admixtures in the germplasm (Delfini *et al.* 2021). Generally, genetic diversity is indispensable to the formation of an initial population for natural selection, the evolutionary direction and future adaptive changes (Govindaraj *et al.* 2015, Lloyd *et al.* 2016). In conservation efforts, an understanding of genetic diversity is essential for maximising the effectiveness and efficiency of the programme.

Rice cultivation

Although all of the swamp rice accessions that we planted can grow, their growth is not optimal. The main obstacle is soil agrophysical differences between their original habitats and the ISARI research station, which has a potentially sour sulfate soil with Type B water overflow (by large tides only; Vepraskas & Craft 2016). According to Vepraskas & Craft (2016), this soil type is characterised by a pyrite content of >2 %, unoxidised, and located at a depth of <50 cm from the soil surface.




Figure 5. Cultivation of Indonesian swamp rice germplasm at the ISARI Research Station in South Kalimantan. A: seedlings grown from seeds planted in slightly dry soil on the edge of a swamp rice field; B: seedlings after translocation to the planting site; C: four-week old rice seedlings after planting out; D: flowering plants protected from birds by para-net installation.

Table A1 shows that the swamp rice was collected two different types of land, namely tidal swamps and inland swamps. According to Craft (2016) and Kennish (2016), there are significant differences between these two habitats. The tidal ecosystem is naturally characterised by soils that are water-saturated or shallowly stagnant year-round or for several months per year. The dominant soil formation process is gleization which is characterised by a grey-bluish reduced soil horizon and the formation of a peat layer on the surface. In contrast, the inland swamp is located near the middle of the catchment, upstream of any tidal influence but subject to periodic flooding of the river during the rainy season. This land is gradually inundated by stagnant water which subsequently recedes with the change from rainy

season to the dry season of the following year (Craft 2016, Kennish 2016).

According to Michael (2020), problems in acid sulfate soils are due to the presence of a pyrite (FeS_2) layer that undergoes oxidation and causes soil acidification. In very acidic conditions ($\text{pH} < 4$), the solubility of aluminum increases drastically and can poison the plants (Khairullah 2020). Poisoning usually occurs in dry soil or during a prolonged drought (Shamshuddin *et al.* 2013). Conversely, when the soil is flooded, the increase in pH can lead to the reduction of ferric iron to ferrous. This phenomenon occurs mainly in actual acid sulfate soils (when pyrite has oxidised) that are re-inundated by rain or tidal water (Khairullah *et al.* 2021, Shamshuddin *et al.* 2013). Ferrous iron




Figure 6. Plant height variation (cm) of Indonesian swamp rice germplasm growing at the ISARI Research Station, South Kalimantan, Indonesia.

concentrations of 300–400 ppm are highly poisonous to rice plants and cause low nutrient availability (Craft 2016, Panhwar *et al.* 2016).

In the inland swamps, the main problem is the fluctuations of inundation between rainy and dry seasons (Mehner & Tockner 2022). Consequently, most farmers have difficulty predicting the degree of waterlogging, especially when planting and when the plants are entering the vegetative growth phase. So far, farmers have adapted their agronomic methods by delaying planting and by translocating seedlings more than once (Paiman *et al.* 2020). In addition to the problem of waterlogging, the land conditions of inland swamps are very acidic, causing high availability of Al and Fe (Maruapey *et al.* 2020). On the other hand, the availability of plant macronutrients such as nitrogen, phosphorus and potassium is reduced (Maruapey *et al.* 2020). In the context of agronomy, efforts to increase crop productivity include improving fertilisation methods in terms of dosage, timing and method of administration (Mehner & Tockner 2022).

So how does the rice plant grow in swamp locations? Physiologically, its success is probably due to the mechanism of lengthening the stems and storing energy during submergence, which is an adaptation of rice plants to swampland conditions (Setter *et al.* 1997). Local adaptations of rice to heavy metal poisoning, especially by Fe, include avoidance mechanisms and tissue tolerance. Asch *et al.* (2005) explain that the avoidance mechanism may be related to the ability of rice to oxidise Fe^{2+} to Fe^{3+} on the root surface, resulting in the formation of harmless iron plaques. As a result of this mechanism, there are also differences in the distribution of Fe in roots, stems and leaves that enable the rice plant to grow in swamplands (Sahrawat 2000, Audebert 2006).

The measurements of plant height show that most of our rice accessions grow to heights of more than 100 cm (Figure 6), which makes the germplasm suitable for use as a parent or for further development

in waterlogged swamp habitats. According to Wu *et al.* (2011), plant height must be correlated with the size (diameter) of the stem to minimise lodging. In breeding programmes, lodging is a limiting factor on potential yields because it can reduce the canopy for photosynthesis, increase respiration and susceptibility to disease, and limit the translocation of nutrients and carbon to the grain (Wu *et al.* 2011). Ünán *et al.* (2013) report that lodging is correlated with plant height. At molecular level, the plant height variations are controlled by a gene regulator, i.e., *rice plasticity 1 (RPL1)*, located on chromosome 6 (Nayar 2014).

Regardless of the land conditions, several pests and diseases also present obstacles to the growth of swamp rice at the ISARI site. The most troublesome are white rice stem borers, brown and green leafhoppers, and ear bugs (*Leptocoris oratorius*, Hemiptera) including blast and tungro. According to Susanti *et al.* (2016), these pests and diseases are common in most Indonesian swamp rice farms. To eliminate or reduce such problems they suggest the use of natural reagents and predators, as well as some safe chemicals; for example, spiders (Arachnida), extracts of ‘purun’ or water chestnut (*Eleocharis dulcis*), silicates from rice husk ash, and potassium, as well as sex pheromones (Susanti *et al.* 2016).

Future programmes

Since this is the first step towards ex-situ conservation, some follow-up activities are urgent. For example, the conservation effort should be supported by creating a database or core collections (Kumar *et al.* 2020). This would reduce conservation costs, increase management efficiency, and continue to support the utilisation of genetic diversity through the precise and rapid identification of sources of genetic diversity for specific trait improvement programmes (Kumar *et al.* 2020, Tanaka *et al.* 2021). According to Lioy *et al.* (2015), the development of core collections is an efficient approach to exploring,



characterising and utilising the genetic diversity of large populations. This activity also has strategic value because it can facilitate exploration of the germplasm resource for new genes (Tanaka *et al.* 2021).

Conceptually, the core collection is a set of accessions with maximum genetic diversity and minimum repetition, including existing ecologically and genetically different ones (Kumar *et al.* 2020, Vilayheuang *et al.* 2020). In other words, a core collection is composed of a small percentage of representative genotypes drawn from most of the genetic diversity of the entire population (Tanaka *et al.* 2021). The size of core collections varies between 5 % and 30 % of the overall population (Liu *et al.* 2015). However, the core collection should be reviewed regularly (Kumar *et al.* 2020).

Characterisation of the germplasm collection must also be done systematically and periodically (Kumar *et al.* 2020), using both morphological (phenotypic) and molecular markers; e.g., SNPs that are faster, more accurate, and stable against environmental influences (Mursyidin *et al.* 2018). According to Liu *et al.* (2015), analysing differences in genetic diversity between core collections and the entire population is essential to testing the effectiveness of such activities. In other words, characterisation is carried out to avoid duplication, contamination, mixing or mislabelling of germplasm collections and to monitor the viability of these genetic resources for the future (Ahmed & Iftekharuddaula 2017, Vilayheuang *et al.* 2020).

Finally, to maintain the viability of germplasm and to determine when germination tests should be made, cold storage of the seed and plant rejuvenation are also needed (Maxted *et al.* 2020).

ACKNOWLEDGEMENTS

The authors are grateful to the Director of ISARI, South Kalimantan, Indonesia, for facilitating this study. We also thank Zainuddin, Bejo Utomo and Riduansyah Syamsudin for their help with cultivation on the rice field and for characterising the grain traits.

AUTHOR CONTRIBUTIONS

DHM initiated the study, carried out the data analyses, created most of the Figures and Tables, and wrote the first draft of the manuscript. IK designed the field experiment and documentation. MS did the fieldwork and morphological observations. All authors contributed to interpretation of the results and to writing the final manuscript.

REFERENCES

- Ahmed, M.S., Iftekharuddaula, K.Md. (2017) Selection of core collection from Jesso-Balam rice (*Oryza sativa* L.) accessions using quantitative, qualitative and molecular characters - a review. *The Agriculturists*, 15(1), 170–181.
- Asch, F., Becker, M., Kpongor, D.S. (2005) A quick and efficient screen for tolerance to iron toxicity in lowland rice. *Journal of Plant Nutrition and Soil Science*, 168, 764–773.
- Audebert, A. (2006) Iron partitioning as a mechanism for iron toxicity tolerance in lowland rice. In: Audebert, A., Narteh, L.T., Millar, D., Beks, B. (eds.) *Iron Toxicity in Rice-Based Systems in West Africa*, WARDA (Africa Rice Center), Cotonou, 34–46.
- Azeez, M.A., Adubi, A.O., Durodola, F.A. (2018) Landraces and crop genetic improvement. In: Grillo, O. (ed.) *Rediscovery of Landraces as a Resource for the Future*, IntechOpen, London, UK, 1–20.
- Calingacion, M., Laborte, A., Nelson, A., Resurreccion, A. and 44 others (2014) Diversity of global rice markets and the science required for consumer-targeted rice breeding. *PLoS ONE*, 9(1), e85106, 12 pp.
- Craft, C.B. (2016) *Creating and Restoring Wetlands: From Theory to Practice*. Elsevier, New York, USA, 336 pp.
- Delfini, J., Moda-Cirino, V., dos Santos Neto, J., Ruas, P.M., Sant'Ana, G.C., Gepts, P., Gonçalves, L.S.A. (2021) Population structure, genetic diversity and genomic selection signatures among a Brazilian common bean germplasm. *Scientific Reports*, 11(1), 2964, 12 pp.
- Fu, J.R., Zhu, L.X., Sun, X.T., Zhou, D.H., Ouyang, L.J., Bian, J.M., He, H.H., Xu, J. (2015) Genetic analysis of grain shape and weight after cutting rice husk. *Genetics and Molecular Research*, 14(4), 17739–17748.
- Galetti, P.M.Jr. (2018) Conservation genetics. In: Caplan, M.J. (ed.) *Reference Module in Biomedical Sciences*, Elsevier Inc., New York, USA, 1–4.
- Giesen, W., Wijedasa, L.S., Page, S.E. (2018) Unique Southeast Asian peat swamp forest habitats have relatively few distinctive plant species. *Mires and Peat*, 22, 01, 13 pp.
- Govindaraj, M., Vetriventhan, M., Srinivasan, M. (2015) Importance of genetic diversity assessment in crop plants and its recent advances: An overview of its analytical perspectives. *Genetics Research International*, 2015, 431487, 14 pp.
- Gunawan, H., Kobayashi, S., Mizuno, K., Kono, Y.



- (2012) Peat swamp forest types and their regeneration in Giam Siak Kecil-Bukit Batu Biosphere Reserve, Riau, East Sumatra, Indonesia. *Mires and Peat*, 10, 05, 17 pp.
- Harrison, M.E., Ottay, J.B., D'Arcy, L.J., Cheyne, S.M. and 27 others (2020) Tropical forest and peatland conservation in Indonesia: Challenges and directions. *People and Nature*, 2(1), 4–28.
- Jackson, S.A. (2016) Rice: The first crop genome. *Rice*, 9, 14, 3 pp.
- Kennish, M.J. (2016) *Encyclopedia of Estuaries*. Springer, New York, USA, 778 pp.
- Khairullah, I. (2020) Indigenous knowledge cultivation of local rice varieties “Siam Mutiara” and “Siam Saba” at tidal swampland. *BIO Web of Conferences*, 20, 01007, 7 pp.
- Khairullah, I., Alwi, M., Masganti (2021) Increasing productivity of rice through iron toxicity control in acid sulfate soils of tidal swampland. *IOP Conference Series: Earth and Environmental Science*, 648, 012151, 13 pp.
- Khan, A.S., Imran, M., Ashfaq, M. (2009) Estimation of genetic variability and correlation for grain yield components in rice (*Oryza sativa* L.). *American-Eurasian Journal of Agricultural & Environmental Sciences*. 6(5), 585–590.
- Kiple, K.F., Ornelas, K.C. (eds.) (2000) *The Cambridge World History of Food*. Cambridge University Press, Cambridge, UK, 2210 pp.
- Kumar, A., Kumar, S., Singh, K.B.M., Prasad, M., Thakur, J.K. (2020) Designing a mini-core collection effectively representing 3004 diverse rice accessions. *Plant Communications*, 1(5), 100049, 14 pp.
- Liu, W., Shahid, M.Q., Bai, L., Lu, Z., Chen, Y., Jiang, L., Diao, M., Liu, X., Lu, Y. (2015) Evaluation of genetic diversity and development of a core collection of wild rice (*Oryza rufipogon* Griff.) populations in China. *PLoS ONE*, 10(12), e0145990, 23 pp.
- Lloyd, M.M., Makukhov, A.D., Pespeni, M.H. (2016) Loss of genetic diversity as a consequence of selection in response to high pCO₂. *Evolutionary Applications*, 9(9), 1124–1132.
- Malhotra, N., Panatu, S., Singh, B., Negi, N., Singh, D., Singh, M., Chandora, R. (2018) Genetic resources: Collection, conservation, characterization and maintenance. In: Singh, M. (ed.) *Lentils: Potential Resources for Enhancing Genetic Gains*, Elsevier, New York, USA, 21–41.
- Maruapey, A., Wicaksana, N., Karuniawan, A., Windarsih, G., Utami, D.W. (2020) Swampy rice lines for iron toxicity tolerance and yield components performance under inland swamp at Sorong, West Papua, Indonesia. *Biodiversitas*, 21(11), 5394–5402.
- Maxted, N., Hunter, D., Rios, R.O. (eds.) (2020) *Plant Genetic Conservation*. Cambridge University Press, Cambridge, UK, 249–277.
- Mehner, T., Tockner, K. (2022) *Encyclopedia of Inland Waters*. Elsevier Inc., New York, USA, 2838 pp.
- Michael, P.S. (2020) Simple carbon and organic matter addition in acid sulfate soils and time-dependent changes in pH and redox under varying moisture regimes. *Asian Journal of Agriculture*, 4(1), 23–29.
- Mitsch, W.J. (2013) Wetland creation and restoration. In: Levin, S.A. (ed.) *Encyclopedia of Biodiversity*, Elsevier Inc., New York, USA, 367–383.
- Mursyidin, D.H., Nazari, Y.A., Daryono, B.S. (2017) Tidal swamp rice cultivars of South Kalimantan Province, Indonesia: A case study of diversity and local culture. *Biodiversitas*, 18(1), 427–432.
- Mursyidin, D.H., Purnomo, P., Sumardi, I., Daryono, B.S. (2018) Molecular diversity of tidal swamp rice (*Oryza sativa* L.) in South Kalimantan, Indonesia. *Diversity*, 10, 22, 10 pp.
- Mursyidin, D.H., Purnomo, Sumardi, I., Daryono, B.S. (2019) Phenotypic diversity of the tidal swamp rice (*Oryza sativa* L.) germplasm from South Kalimantan, Indonesia. *Australian Journal of Crop Science*, 13(3), 386–394.
- Mursyidin, D.H., Purnomo, Daryono, B.S. (2021) The ligule ultrastructure of the tidal swamp rice (*Oryza sativa*) landraces of South Kalimantan, Indonesia, and their genetic diversity and relationship. *Biodiversitas*, 22(12), 5280–5285.
- Nayar, N.M. (2014) *Origins and Phylogeny of Rices*. Academic Press, Oxford, UK, 322 pp.
- Nourollah, A. (2016) Genetic diversity, genetic erosion, and conservation of the two cultivated rice species (*Oryza sativa* and *Oryza glaberrima*) and their close wild relatives. In: Ahuja, M.R., Jain, S.M. (eds.) *Genetic Diversity and Erosion in Plants*, Springer International Publishing, Cham, Switzerland, 35–73.
- Offord, C.A. (2016) Germplasm conservation. In: Thomas, B., Murray, B.G., Murphy, D.J. (eds.) *Encyclopedia of Applied Plant Sciences*, Elsevier Inc., New York, USA, 281–288.
- Page, S., Hooijer, A., Rieley, J., Banks, C., Hoscilo, A. (2012) The tropical peat swamps of Southeast Asia: human impacts on biodiversity, hydrology, and carbon dynamics. In: Gower, D.J., Johnson, K.G., Richardson, J.E., Rosen, B.R., Rüber, L., Williams, S.T. (eds.) *Biotic Evolution and Environmental Change in Southeast Asia*, Cambridge University Press, Cambridge, UK, 406–433.



- Paiman, Ardiyanta, Ansar, M., Effendy, I., Sumbodo, B.T. (2020) Rice cultivation of superior variety in swamps to increase food security in Indonesia. *Reviews in Agricultural Science*, 8, 300–309.
- Panhwar, Q.A., Naher, U.A., Shamsuddin, J., Radziah, O., Hakeem, K.R. (2016) Management of acid sulfate soils for sustainable rice cultivation in Malaysia. In: Hakeem, K.R., Akhtar, J., Sabir, M. (eds.) *Soil Science: Agricultural and Environmental Perspectives*. Springer International Publishing, Cham, Switzerland, 91–104.
- Qurani, I.Z., Lakitan, B. (2021) *Inland Swamp Agriculture: Opportunities and Challenges*. TJF Brief April 2021, Tay Juhana Foundation, Jakarta, Indonesia, 5 pp.
- Rao, I.S., Neeraja, C.N., Srikanth, B., Subrahmanyam, D., Swamy, K.N., Rajesh, K., Vijayalakshmi, P., Kiran, T.V., Sailaja, N., Revathi, P., Rao, P.R., Rao, L.V.S., Surekha, K., Babu, V.R., Voleti, S.R. (2018) Identification of rice landraces with promising yield and the associated genomic regions under low nitrogen. *Scientific Reports*, 8, 9200, 13 pp.
- Sahrawat, K.L. (2000) Elemental composition of the rice plant as affected by iron toxicity under field conditions. *Communications in Soil Science and Plant Analysis*, 31, 2819–2827.
- Sanghera, G.S., Kashyap, S.C., Parray, G.A. (2013) Genetic variation for grain yield and related traits in temperate red rice (*Oryza sativa* L.) ecotype. *Notulae Scientia Biologica*, 5(3), 1–7.
- Shamsuddin, J., Elisa, A.A., Shazana, M.A.R.S., Fauziah, I.C. (2013) Rice defense mechanisms against the presence of excess amount of Al³⁺ and Fe²⁺ in the water. *Australian Journal of Crop Science*, 7(3), 314–320.
- Setter, T.L., Laureles, E.V., Mazaredo, A.M. (1997) Lodging reduces yield of rice by self-shading and reduction in canopy photosynthesis. *Field Crops Research*, 49, 95–106.
- Sulaiman, A.A., Sulaeman, Y., Minasny, B. (2019) A framework for the development of wetland for agricultural use in Indonesia. *Resources*, 8, 34, 16 pp.
- Sulakhudin, Hatta, M. (2018) Increasing productivity of newly opened paddy field in tidal swampy areas using a local specific technology. *Indonesian Journal of Agricultural Science*, 19(1), 9–16.
- Susanti, M.A., Trahmrin, M., Asikin, S. (2016) Management of insect major pest of rice in tidal swampland. *Prosiding Seminar Nasional Lahan Basah (Proceedings of the National Seminar on Wetlands)*, 1, 1–10.
- Tanaka, N., Shenton, M., Kawahara, Y., Kumagai, M., Sakai, H., Kanamori, H., Yonemaru, J.I., Fukuoka, S., Sugimoto, K., Ishimoto, M., Wu, J., Ebana, K. (2021) Investigation of the genetic diversity of a rice core collection of Japanese landraces using whole-genome sequencing. *Plant & Cell Physiology*, 61(12), 2087–2096.
- Tran, D.S., Tran, T.T.H., Nguyen, T.L.H., Ha, M.L., Dinh, B.Y., Kumamaru, T., Satoh, H. (2012) Variation on grain quality in Vietnamese rice cultivars collected from central Vietnam. *Journal of the Faculty of Agriculture, Kyushu University*, 57(2), 365–371.
- Trettin, C.C., Jurgensen, M.F., Dai, Z. (2019) Effects of climate change on forested wetland soils. In: Busse, M., Giardina, C., Morris, D., Page-Dumroese, D. (eds.) *Global Change and Forest Soils*, Elsevier B.V., Amsterdam, Netherlands, 171–188.
- Ünan, R., Sezer, İ., Şahin, M., Mur, L.A.J. (2013) Control of lodging and reduction in plant length in rice (*Oryza sativa* L.) with the treatment of trinexapac-ethyl and sowing density. *Turkish Journal of Agriculture and Forestry*, 37, 257–264.
- Vepraskas, M.J., Craft, C.B. (2016) *Wetland Soils: Genesis, Hydrology, Landscape, and Classification*. CRC Press, Boca Raton, USA, 523 pp.
- Vilayheuang, K., Borrayo, E., Kawase, M., Watanabe, K.N. (2020) Development of Laos Khao Kai Noi rice landrace (*Oryza sativa* L.) core collection as a model for rice genetic resources management in the Laos National Genebank. *IOP Conference Series: Earth and Environmental Science*, 482(1), 012039, 6 pp.
- Wang, Y. (2020) *Wetlands and Habitats*. CRC Press, Boca Raton, USA, 311 pp.
- Wu, L.-L., Liu, Z.-L., Wang, J.-M., Zhou, C.-Y., Chen, K.-M. (2011) Morphological, anatomical, and physiological characteristics involved in development of the large culm trait in rice. *Australian Journal of Crop Science*, 5(11), 1356–1363.
- Wu, F., Ma, S., Zhou, J., Han, C., Hu, R., Yang, X., Nie, G., Zhang, X. (2021) Genetic diversity and population structure analysis in a large collection of white clover (*Trifolium repens* L.) germplasm worldwide. *PeerJ*, 9, e11325, 17 pp.
- Zhang, P., Li, J., Li, X., Liu, X., Zhao, X., Lu, Y. (2011) Population structure and genetic diversity in a rice core collection (*Oryza sativa* L.) investigated with SSR markers. *PLoS ONE*, 6(12), e27565, 13 pp.
- Zheng, J., Zhang, Y., Wang, C. (2015) Molecular functions of genes related to grain shape in rice. *Breeding Science*, 65(2), 120–126.



Zhou, Y., Miao, J., Gu, H., Peng, X., Leburu, M., Yuan, F., Gu, H., Gao, Y., Tao, Y., Zhu, J., Gong, Z., Yi, C., Gu, M., Yang, Z., Liang, G. (2015) Natural variations in SLG7 regulate grain shape in rice. *Genetics*, 201(4), 1591–1599.

Submitted 28 Aug 2022, revision 03 Nov 2022
Editor: Olivia Bragg

Author for correspondence:

Dr Dindin Hidayatul Mursyidin MSc, University of Lambung Mangkurat, Faculty of Mathematics and Natural Sciences, Laboratory of Genetics and Molecular Biology, South Kalimantan, 70714, Indonesia.
Tel: +62 (0)511-4773112; E-mail: dindinhm@gmail.com



Appendix

Table A1. ⁷ List of Indonesian swamp rice accessions used in the study, including their origins and natural habitats.

Province of origin	Name of accession	Code	Habitat
Jambi	<i>Mentol</i>	72	Tidal swamp
	<i>Mentong</i>	36	Tidal swamp
	<i>Menuh</i>	103	Tidal swamp
	<i>Sawah Kanyut</i>	73	Tidal swamp
South Sumatra	<i>Belut</i>	78	Tidal swamp
	<i>Betek</i>	33	Tidal swamp
	<i>Bonai Tinggi</i>	74	Tidal swamp
	<i>Cempak Merah</i>	60	Tidal swamp
	<i>Cemurai</i>	75	Tidal swamp
	<i>Mayes</i>	59	Inland swamp
	<i>Padang</i>	55	Inland swamp
	<i>Sawo Rampak</i>	77	Inland swamp
	<i>Pelita Rampak</i>	16	Inland swamp
	<i>Sardani</i>	56	Tidal swamp
Lampung	<i>Serai Rampak</i>	105	Inland swamp
	<i>Awan Kuning</i>	76	Tidal swamp
	<i>Kamajaya</i>	80	Tidal swamp
	<i>Karat Kaleng</i>	5	Tidal swamp
	<i>Katimuri</i>	58	Tidal swamp
	<i>Kebau</i>	70	Tidal swamp
	<i>Kencana Baliman</i>	61	Tidal swamp
	<i>Keromojoyo</i>	71	Tidal swamp
	<i>Ketan Bujule</i>	79	Tidal swamp
	<i>Ketek Muri</i>	107	Tidal swamp
	<i>Ketek Semut</i>	99	Tidal swamp
	<i>Kuning Sore</i>	38	Tidal swamp
	<i>Kutut</i>	25	Tidal swamp
	<i>Lembo Sawo</i>	2	Tidal swamp
	<i>Lembu Sawah</i>	54	Tidal swamp
	<i>Lembu Sawah Tinggi</i>	34	Tidal swamp
	<i>Petek Kuning</i>	3	Tidal swamp
	<i>Punai Baru</i>	66	Tidal swamp
	<i>Senapi</i>	52	Tidal swamp
	<i>Senapi Super</i>	98	Tidal swamp
<i>Tampu</i>	53	Tidal swamp	
<i>Tumbara</i>	24	Tidal swamp	
West Kalimantan	<i>Randah Pala</i>	37	Tidal swamp
	<i>Selumbang</i>	65	Tidal swamp
	<i>Umbung Buluh</i>	19	Tidal swamp
Central Kalimantan	<i>Padi Merah</i>	18	Tidal swamp
	<i>Pahakang</i>	17	Tidal swamp
	<i>Pulut Air</i>	13	Tidal swamp
	<i>Pulut Kemenyan</i>	31	Tidal swamp
	<i>Raanti</i>	62	Tidal swamp
	<i>Raden Pulatar</i>	97	Tidal swamp
	<i>Kapuas</i>	104	Tidal swamp

³ *Mires and Peat*, Volume 28 (2022), Article 35, 17 pp., <http://www.mires-and-peat.net/>, ISSN 1819-754X
International Mire Conservation Group and International Peatland Society, DOI: 10.19189/Map.2022.OMB.566



Province of origin	Name of accession	Code	Habitat
South Kalimantan	<i>Adil Kuning</i>	100	Tidal swamp
	<i>Badagai</i>	101	Tidal swamp
	<i>Baliman Putih</i>	21	Tidal swamp
	<i>Banih Kuning</i>	39	Tidal swamp
	<i>Bayar Pahit</i>	87	Tidal swamp
	<i>Biduin</i>	69	Tidal swamp
	<i>Ketan</i>	83	Tidal swamp
	<i>Ketan BundeI</i>	92	Tidal swamp
	<i>Ketan Hitam</i>	11	Tidal swamp
	<i>Ketan Kuning</i>	14	Tidal swamp
	<i>Ketan Merah</i>	29	Tidal swamp
	<i>Ketan Merah Bule</i>	50	Tidal swamp
	<i>Ketan Selung</i>	28	Tidal swamp
	<i>Ketan Serang</i>	44	Tidal swamp
	<i>Kuning</i>	81	Tidal swamp
	<i>Lakatan Gadur</i>	12	Tidal swamp
	<i>Lakatan Hirang</i>	15	Tidal swamp
	<i>Lalantik</i>	32	Tidal swamp
	<i>Lemo</i>	67	Tidal swamp
	<i>Pandak Baru</i>	48	Tidal swamp
	<i>Pandak Kembang</i>	85	Tidal swamp
	<i>Pandak Tinggi</i>	7	Tidal swamp
	<i>Puput</i>	106	Tidal swamp
	<i>Putih</i>	96	Tidal swamp
	<i>Randah Pandang</i>	41	Tidal swamp
	<i>Raden Rata</i>	40	Tidal swamp
	<i>Siam Adus/Abu</i>	94	Tidal swamp
	<i>Siam Berandal</i>	63	Tidal swamp
	<i>Siam Ganai</i>	1	Tidal swamp
	<i>Siam Gumpal</i>	93	Tidal swamp
	<i>Siam Halus</i>	9	Tidal swamp
	<i>Siam Karang Dukuh Kuning</i>	64	Tidal swamp
	<i>Siam Karang Dukuh Putih</i>	8	Tidal swamp
	<i>Siam Kretik/Keriting</i>	46	Tidal swamp
	<i>Siam Mayang</i>	47	Tidal swamp
	<i>Siam Mutiara</i>	26	Tidal swamp
	<i>Siam Pangling</i>	95	Tidal swamp
	<i>Siam Panting</i>	23	Tidal swamp
	<i>Siam Parupuk</i>	35	Tidal swamp
	<i>Siam Perak</i>	102	Tidal swamp
	<i>Siam Perak Halus</i>	4	Tidal swamp
	<i>Siam Pontianak Halus</i>	22	Tidal swamp
	<i>Siam Pontianak Tinggi</i>	42	Tidal swamp
	<i>Siam Puntal</i>	88	Tidal swamp
	<i>Siam Rukut</i>	10	Tidal swamp
	<i>Siam Sabar</i>	27	Tidal swamp
	<i>Siam Salawi</i>	90	Tidal swamp
	<i>Siam Super</i>	91	Tidal swamp
	<i>Siam Tanggung</i>	86	Tidal swamp
<i>Siam Teladan</i>	89	Tidal swamp	
<i>Siam Unus Gampa</i>	43	Tidal swamp	
<i>Super</i>	49	Tidal swamp	



Province of origin	Name of accession	Code	Habitat
South Kalimantan	<i>Unus Organik</i>	57	Tidal swamp
	<i>Unyil</i>	6	Tidal swamp
East Kalimantan	<i>Kamajaya Hitam</i>	45	Tidal swamp
	<i>Kamajaya Putih</i>	51	Tidal swamp
	<i>Semeru</i>	84	Tidal swamp
	<i>Serai</i>	82	Tidal swamp
	<i>Siam Bohai Pendek</i>	68	Tidal swamp
	<i>Siyam</i>	30	Tidal swamp
	<i>Tiga Dara Sawah Beling</i>	20	Tidal swamp



Table A2. Characteristics of Indonesian swamp rice germplasm based on the grain length, width, and ratio, including grain type and subspecies.

Name of accession	Code	Province of origin	Grain length (mm)	Grain width (mm)	Grain ratio (length/width)	Grain type	Subspecies
Petek Kuning	3	Lampung	9.43 ± 0.31	1.80 ± 0.10	5.24	very long-slender	indica
Siam Halus	9	South Kalimantan	9.37 ± 0.23	1.83 ± 0.06	5.11	very long-slender	indica
Lakatan Hirang	15	South Kalimantan	9.30 ± 0.17	2.53 ± 0.06	3.67	very long-slender	indica
Siam Berandal	63	South Kalimantan	9.20 ± 0.61	1.93 ± 0.23	4.76	very long-slender	indica
Siam Perak	102	South Kalimantan	9.20 ± 0.30	2.43 ± 0.15	3.78	very long-slender	indica
Banli Kuning	39	South Kalimantan	9.13 ± 0.47	2.50 ± 0.26	3.65	very long-slender	indica
Pulut Kemenyan	31	Central Kalimantan	9.07 ± 0.55	2.83 ± 0.06	3.20	very long-slender	indica
Sawah Kanyut	73	Jambi	9.07 ± 0.40	2.23 ± 0.25	4.06	very long-slender	indica
Pahakang	17	Central Kalimantan	9.00 ± 0.53	2.10 ± 0.20	4.29	very long-slender	indica
Super	49	South Kalimantan	9.00 ± 0.20	2.67 ± 0.15	3.38	very long-slender	indica
Siam Gatal	1	South Kalimantan	8.97 ± 0.75	2.20 ± 0.20	4.08	very long-slender	indica
Mentong	36	Jambi	8.97 ± 0.38	2.50 ± 0.26	3.59	very long-slender	indica
Siam Salawi	90	South Kalimantan	8.93 ± 0.31	2.47 ± 0.12	3.62	very long-slender	indica
Pulut Air	13	Central Kalimantan	8.93 ± 0.15	1.70 ± 0.10	5.25	very long-slender	indica
Ketan	83	Lampung	8.90 ± 0.61	2.33 ± 0.23	3.81	very long-slender	indica
Siam Sabar	27	South Kalimantan	8.90 ± 0.26	1.83 ± 0.06	4.85	very long-slender	indica
Ketan Merah Bule	50	South Kalimantan	8.90 ± 0.17	2.23 ± 0.15	3.99	very long-slender	indica
Senapi Super	98	Lampung	8.87 ± 0.60	2.43 ± 0.06	3.65	very long-slender	indica
Siam Puntal	88	South Kalimantan	8.87 ± 0.49	1.87 ± 0.06	4.75	very long-slender	indica
Ketan Bijule	79	South Kalimantan	8.87 ± 0.29	2.47 ± 0.12	3.60	very long-slender	indica
Lambu Sawah	54	Lampung	8.83 ± 0.76	2.37 ± 0.12	3.73	very long-slender	indica
Kanjajaya	80	Lampung	8.83 ± 0.12	2.50 ± 0.26	3.53	very long-slender	indica
Kapuas	104	Central Kalimantan	8.80 ± 0.92	2.37 ± 0.15	3.72	very long-slender	indica
Pandak Tinggi	7	South Kalimantan	8.80 ± 0.26	1.83 ± 0.15	4.80	very long-slender	indica
Pandak Baru	48	South Kalimantan	8.77 ± 0.29	2.33 ± 0.12	3.76	very long-slender	indica
Siam Perak Halus	4	South Kalimantan	8.73 ± 0.06	1.97 ± 0.12	4.44	very long-slender	indica
Semeru	84	East Kalimantan	8.70 ± 0.85	2.30 ± 0.26	3.78	very long-slender	indica
Kuning	81	South Kalimantan	8.67 ± 0.60	2.47 ± 0.31	3.51	very long-slender	indica
Raanti	62	Central Kalimantan	8.67 ± 0.51	2.37 ± 0.12	3.66	very long-slender	indica
Kuttut	25	Lampung	8.67 ± 0.31	1.93 ± 0.06	4.48	very long-slender	indica
Randah Pala	37	West Kalimantan	8.63 ± 0.91	2.43 ± 0.40	3.55	very long-slender	indica
Siam Pangling	95	South Kalimantan	8.60 ± 0.46	2.03 ± 0.12	4.23	very long-slender	indica
Ketan Hitam	11	South Kalimantan	8.60 ± 0.40	2.20 ± 0.20	3.91	very long-slender	indica
Ketan Kuning	14	South Kalimantan	8.57 ± 0.64	2.47 ± 0.21	3.47	very long-slender	indica
Unus Organik	57	South Kalimantan	8.50 ± 0.50	1.90 ± 0.00	4.47	very long-slender	indica



Name of accession	Code	Province of origin	Grain length (mm)	Grain width (mm)	Grain ratio (length/width)	Grain type	Subspecies
<i>Pelita Rampak</i>	16	South Sumatera	8.47 ± 0.38	2.37 ± 0.06	3.58	very long-slender	<i>indica</i>
<i>Siam Paripuk</i>	35	South Kalimantan	8.43 ± 0.31	2.33 ± 0.06	3.61	very long-slender	<i>indica</i>
<i>Lelanik</i>	32	South Kalimantan	8.40 ± 1.08	2.23 ± 0.12	3.76	very long-slender	<i>indica</i>
<i>Umbung Buluh</i>	19	West Kalimantan	8.40 ± 0.17	2.20 ± 0.17	3.82	very long-slender	<i>indica</i>
<i>Siam Mayang</i>	47	South Kalimantan	8.37 ± 0.25	1.93 ± 0.06	4.33	very long-slender	<i>indica</i>
<i>Badagai</i>	101	South Kalimantan	8.33 ± 0.64	1.97 ± 0.12	4.24	very long-slender	<i>indica</i>
<i>Adil Kuning</i>	100	South Kalimantan	8.33 ± 0.21	1.90 ± 0.00	4.38	very long-slender	<i>indica</i>
<i>Siam Unus Gampa</i>	43	South Kalimantan	8.27 ± 0.93	2.27 ± 0.12	3.65	very long-slender	<i>indica</i>
<i>Siam Panting</i>	23	South Kalimantan	8.27 ± 0.40	1.80 ± 0.00	4.59	very long-slender	<i>indica</i>
<i>Raden Rata</i>	40	South Kalimantan	8.27 ± 0.15	2.23 ± 0.06	3.70	very long-slender	<i>indica</i>
<i>Randah Pandang</i>	41	South Kalimantan	8.23 ± 0.86	2.60 ± 0.26	3.17	very long-slender	<i>indica</i>
<i>Kuning Sore</i>	38	Lampung	8.20 ± 0.30	2.20 ± 0.17	3.73	very long-slender	<i>indica</i>
<i>Kencana Baliman</i>	61	Lampung	8.20 ± 0.10	2.53 ± 0.06	3.24	very long-slender	<i>indica</i>
<i>Siyam</i>	36	East Kalimantan	8.20 ± 0.10	2.70 ± 0.10	3.04	very long-slender	<i>indica</i>
<i>Siam Kretik/Keriting</i>	46	South Kalimantan	8.17 ± 0.78	2.37 ± 0.12	3.45	very long-slender	<i>indica</i>
<i>Siam Mutara</i>	26	South Kalimantan	8.17 ± 0.64	2.33 ± 0.21	3.50	very long-slender	<i>indica</i>
<i>Siam Pontanak Tinggi</i>	42	South Kalimantan	8.17 ± 0.45	2.00 ± 0.10	4.09	very long-slender	<i>indica</i>
<i>Siam Tanggung</i>	86	South Kalimantan	8.17 ± 0.35	1.97 ± 0.12	4.15	very long-slender	<i>indica</i>
<i>Pandak Kembang</i>	85	South Kalimantan	8.17 ± 0.06	2.00 ± 0.10	4.09	very long-slender	<i>indica</i>
<i>Katimuri</i>	58	Lampung	8.13 ± 0.64	2.47 ± 0.15	3.30	very long-slender	<i>indica</i>
<i>Siam Teladan</i>	89	South Kalimantan	8.13 ± 0.21	1.97 ± 0.15	4.13	very long-slender	<i>indica</i>
<i>Ketan Bundel</i>	92	South Kalimantan	8.10 ± 0.30	2.27 ± 0.06	3.57	very long-slender	<i>indica</i>
<i>Lakaian Gaddur</i>	12	South Kalimantan	8.07 ± 0.35	2.00 ± 0.10	4.04	very long-slender	<i>indica</i>
<i>Bidain</i>	69	South Kalimantan	8.07 ± 0.32	2.37 ± 0.06	3.41	very long-slender	<i>indica</i>
<i>Tiga Dara Sawah Beling</i>	20	East Kalimantan	8.07 ± 0.27	2.23 ± 0.12	3.61	very long-slender	<i>indica</i>
<i>Bayar Pahit</i>	87	South Kalimantan	8.03 ± 0.45	2.33 ± 0.15	3.44	very long-slender	<i>indica</i>
<i>Tampu</i>	53	Lampung	8.03 ± 0.15	2.60 ± 0.10	3.09	very long-slender	<i>indica</i>
<i>Siam Rukul</i>	10	South Kalimantan	8.00 ± 0.79	2.43 ± 0.12	3.29	very long-slender	<i>indica</i>
<i>Padi Merah</i>	18	Central Kalimantan	8.00 ± 0.10	2.20 ± 0.10	3.64	very long-slender	<i>indica</i>
<i>Belut</i>	78	South Sumatera	7.97 ± 0.32	2.37 ± 0.21	3.37	very long-slender	<i>indica</i>
<i>Siam Karang Dukuh Kuning</i>	64	South Kalimantan	7.97 ± 0.32	2.30 ± 0.17	3.47	very long-slender	<i>indica</i>
<i>Cempak Merah</i>	60	South Sumatera	7.93 ± 0.55	2.47 ± 0.06	3.21	very long-slender	<i>indica</i>
<i>Punai Baru</i>	66	Lampung	7.90 ± 0.20	2.37 ± 0.06	3.34	very long-slender	<i>indica</i>
<i>Siam Bohai Pendek</i>	68	East Kalimantan	7.90 ± 0.17	2.77 ± 0.23	2.86	very long-intermediate	close to <i>indica</i>
<i>Ketan Selang</i>	28	South Kalimantan	7.87 ± 0.49	2.43 ± 0.12	3.23	very long-slender	<i>indica</i>
<i>Ketek Muri</i>	107	Lampung	7.87 ± 0.40	2.40 ± 0.20	3.28	very long-slender	<i>indica</i>



Name of accession	Code	Province of origin	Grain length (mm)	Grain width (mm)	Grain ratio (length/width)	Grain type	Subspecies
<i>Siam Super</i>	91	South Kalimantan	7.83 ± 0.42	2.20 ± 0.36	3.56	very long-slender	<i>indica</i>
<i>Kamajaya Hitam</i>	45	East Kalimantan	7.80 ± 0.30	2.53 ± 0.12	3.08	very long-slender	<i>indica</i>
<i>Lembu Sawah Tinggi</i>	34	Lampung	7.73 ± 0.25	2.37 ± 0.12	3.27	very long-slender	<i>indica</i>
<i>Awan Kuang</i>	76	Lampung	7.73 ± 0.21	2.43 ± 0.06	3.18	very long-slender	<i>indica</i>
<i>Baliman Putih</i>	21	South Kalimantan	7.70 ± 0.20	2.20 ± 0.17	3.50	very long-slender	<i>indica</i>
<i>Siam Karang Dukuh Putih</i>	8	South Kalimantan	7.67 ± 0.06	2.10 ± 0.10	3.65	very long-slender	<i>indica</i>
<i>Mentol</i>	72	Jambi	7.63 ± 0.38	2.60 ± 0.36	2.93	very long-slender	<i>indica</i>
<i>Sawo Rampak</i>	77	South Sumatera	7.53 ± 0.38	2.33 ± 0.06	3.23	very long-slender	<i>indica</i>
<i>Betek</i>	33	South Sumatera	7.50 ± 0.20	2.37 ± 0.12	3.17	long-slender	8 <i>indica</i>
<i>Serai</i>	82	East Kalimantan	7.50 ± 0.10	2.13 ± 0.06	3.52	long-slender	close to <i>indica</i>
<i>Lemo</i>	67	South Kalimantan	7.30 ± 0.30	2.13 ± 0.06	3.42	long-slender	close to <i>indica</i>
<i>Siam Pontianak Halus</i>	22	South Kalimantan	7.10 ± 0.36	1.90 ± 0.26	3.74	long-slender	close to <i>indica</i>
<i>Tumbara</i>	24	Lampung	6.97 ± 0.61	2.33 ± 0.06	2.99	long-intermediate	<i>javanica</i>
<i>Lembo Sawo</i>	2	Lampung	6.80 ± 0.17	2.40 ± 0.20	2.83	long-intermediate	<i>javanica</i>
<i>Siam Adus/Abu</i>	94	South Kalimantan	6.77 ± 0.25	2.33 ± 0.12	2.90	long-intermediate	<i>javanica</i>
<i>Ketan Merah</i>	29	South Kalimantan	6.73 ± 0.35	2.37 ± 0.06	2.84	long-intermediate	<i>javanica</i>
<i>Mayes</i>	59	South Sumatera	6.70 ± 0.36	2.50 ± 0.10	2.68	long-intermediate	<i>javanica</i>
<i>Kebau</i>	70	Lampung	6.67 ± 0.23	2.47 ± 0.12	2.70	long-intermediate	<i>javanica</i>
<i>Menuh</i>	103	Jambi	6.60 ± 0.20	3.00 ± 0.10	2.20	medium-intermediate	8 <i>javanica</i>
<i>Kamajaya Putih</i>	51	East Kalimantan	6.57 ± 0.21	2.60 ± 0.17	2.53	medium-intermediate	close to <i>Japonica</i>
<i>Senapi</i>	52	Lampung	6.53 ± 0.31	2.57 ± 0.12	2.54	medium-intermediate	close to <i>Japonica</i>
<i>Serai Rampak</i>	105	South Sumatera	6.50 ± 0.30	2.63 ± 0.21	2.47	medium-intermediate	8 <i>close to Japonica</i>
<i>Padang</i>	55	South Sumatera	6.50 ± 0.17	2.37 ± 0.15	2.75	medium-intermediate	close to <i>Japonica</i>
<i>Siam Gumpal</i>	93	South Kalimantan	6.47 ± 0.25	2.53 ± 0.06	2.55	medium-intermediate	close to <i>Japonica</i>
<i>Sardani</i>	56	South Sumatera	6.45 ± 0.35	2.57 ± 0.15	2.51	medium-intermediate	close to <i>Japonica</i>
<i>Raden Pulatar</i>	97	Central Kalimantan	6.43 ± 0.15	2.23 ± 0.21	2.88	medium-intermediate	close to <i>Japonica</i>
<i>Unyil</i>	6	South Kalimantan	6.40 ± 0.30	2.23 ± 0.42	2.87	medium-intermediate	close to <i>Japonica</i>
<i>Papat</i>	106	South Kalimantan	6.37 ± 0.32	2.23 ± 0.06	2.85	medium-intermediate	close to <i>Japonica</i>
<i>Selumbang</i>	65	West Kalimantan	6.37 ± 0.31	2.00 ± 0.10	3.19	medium-slender	8 <i>close to Indica</i>
<i>Karat Kaleng</i>	5	Lampung	6.37 ± 0.29	2.20 ± 0.17	2.90	medium-intermediate	close to <i>Japonica</i>
<i>Cemurai</i>	75	South Sumatera	6.33 ± 0.49	2.33 ± 0.21	2.71	medium-intermediate	close to <i>Japonica</i>
<i>Bonai Tinggi</i>	74	South Sumatera	6.30 ± 0.17	2.33 ± 0.23	2.70	medium-intermediate	close to <i>Japonica</i>
<i>Ketek Semut</i>	99	Lampung	6.00 ± 0.30	2.27 ± 0.06	2.65	medium-intermediate	close to <i>Japonica</i>
<i>Putih</i>	96	South Kalimantan	6.00 ± 0.26	2.13 ± 0.06	2.81	medium-intermediate	close to <i>Japonica</i>
<i>Keronjojoyo</i>	71	Lampung	5.90 ± 0.26	2.27 ± 0.15	2.60	medium-intermediate	close to <i>Japonica</i>
<i>Ketan Serang</i>	44	South Kalimantan	5.77 ± 0.31	2.43 ± 0.06	2.37	medium-slender	close to <i>Javanica</i>



Grain diversity and cultivation of Indonesian swamp rice germplasm: building the foundation for an ex-situ conservation programme

ORIGINALITY REPORT

14%

SIMILARITY INDEX

10%

INTERNET SOURCES

6%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1	www.theseus.fi Internet Source	3%
2	Submitted to Institut Pertanian Bogor Student Paper	2%
3	helda.helsinki.fi Internet Source	1%
4	123docz.net Internet Source	1%
5	Dindin Hidayatul Mursyidin. "Phylogenetic relationship of superior durian (<i>Durio zibethinus</i>) cultivars native to South Kalimantan, Indonesia", <i>Pesquisa Agropecuária Tropical</i> , 2022 Publication	1%
6	kasetartjournal.ku.ac.th Internet Source	1%
7	smujo.id Internet Source	1%

8

Submitted to Indira Gandhi Agricultural
University

Student Paper

1 %

9

www.cropj.com

Internet Source

1 %

10

journals.plos.org

Internet Source

1 %

11

www.smujo.id

Internet Source

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On