

Growth Patterns and Relative Condition Factor of Macrobrachium Species from Sungai Batang River, Indonesia

by Ahmadi .

Submission date: 07-Jun-2023 09:39PM (UTC+0700)

Submission ID: 2111079787

File name: terns_and_relative_condition_factor_of_Macrobrachium_species.pdf (819.79K)

Word count: 7981

Character count: 37683

Growth Patterns and Relative Condition Factor of *Macrobrachium* Species from Sungai Batang River, Indonesia

Ahmadi¹, Stephen Rossiter², Pahmi Ansyari³

^{1,3} Faculty of Marine and Fisheries, Lambung Mangkurat University, Indonesia

² School of Biological and Chemical Sciences, Queen Mary University, United Kingdom

Corresponding Author, E-mail address: ahmadizarigani@gmail.com

Abstract: The present study provides the first reference for the growth patterns and relative condition factor of *Macrobrachium* species from Sungai Batang River, Indonesia. A total of 328 specimens, caught by using wire-stage trap, consisted of 83 males (25%) and 245 females (75%) with the sex ratio of 1:3. *Macrobrachium* species showed a negative allometric growth pattern ($b = 1.94-2.14$). The highest percentage of catch falls between 90-94 mm TL (38.55%) and weighted between 9-10 g (46.99%). The average total length and body weight of male were 1.1 and 1.4 times greater than those of female computed by comparing the said body sizes of shrimp. Male had total length, carapace length, abdominal length and chelae length longer than female. The mean relative condition factor values ranged of 0.96 ± 0.17 and 1.01 ± 0.16 , reflecting well-being of the species. Outcome of research could be useful for sustainable fisheries management and conservation measures for *Macrobrachium* species in this river. In addition, a new fishing method was also introduced to improve fishing efficiency.

Key words: *Macrobrachium* species, growth pattern, relative condition factor, wire-stage trap, Sungai Batang River

1. Introduction

In all parts of the world, the length-weight relationships and condition factors of shrimps have been well-documented not only for marine species such as *Acetes japonicus* (Wong, Ong, & Khoo, 2015), *Penaeus monodon* (Uddin, Ghosh, & Maity, 2016) and *Plesionika ensis* (Sousaa et al., 2019), but also for freshwater species like

Macrobrachium rosenbergii (Lalrinsanga *et al.*, 2015), *M. macrobrachion* (Kingdom, Hart, Erondy, & Kwen, 2014), *M. lamarrei* (Ara, Noby, Ahmed, & Fatema, 2014), *M. hainanense* (Mantel & Dudgeon, 2004), *M. malcolmsonii* (Hossain *et al.*, 2012) and *M. assamense peninsulare* (Kumar, Kotnala, & Rana, 2014). Penaeid shrimp and Palaemonid prawns support beneficially aquaculture, capture fishery and shrimp processing industries in countries all over the world, including Indonesia. The shrimp species are being caught by various fishing gears such as trawl nets (Wong, Ong, & Khoo, 2015), cast nets (Sethi, Ram, & Venkatesan, 2012), gillnet (Pravin & Ravindran, 2011), trammel nets (Ahmadi & Kristina, 2017), fyke-nets (Farias, Pereira-Júnior, Domingos, & Dantas, 2019), electrofishing (Ng, 2017), basket traps (Kingdom, Hart, Erondy, & Kwen, 2014), bottom and floating traps (Sousa *et al.*, 2019), and light traps (Ahmadi, 2018a).

The length-weight relationship has important implication in fisheries management and aquaculture practice (Klassen, Gawlik, & Botson, 2014). It is applicable for ¹⁴ estimating growth rates, age structure and other aspect of shrimp population dynamics (Tsoumani *et al.*, 2006), calculating the standing stock biomass (Enin, 1994) and determining condition factor ⁶³ (Li *et al.*, 2016). The length-weight regressions often ⁶ used to estimate weight from length due to the technical constraint in the field. Morphometric traits have been reported to give an accurate estimate of body weight and it can serve as a useful tool for selecting broodstock on weight basis (Daud & Ang, 1995). Variability in the length-weight relationship, as an indicator of condition in marine fishes and crustaceans, is a feature that can reflect fluctuations in the uptake and allocation of energy. These morphometric variations ⁴⁵ can be affected by many factors such as reproductive cycle (Chu, Chen, Huang, & Wong, 1995), genetic differences (Cadrin,

2000) and environment factors such as water temperature, nutrient supply (Castilho et al., 2007) and also interspecific characteristic between species (Bissaro, Gomes-Jr, & Beneditto, 2013). ³ Fulton's condition factor (K) is often used to quantify an animal's physical wellbeing and thought to be a useful complement to growth estimates of crustaceans. ¹⁹ Information on the length-weight relationship and condition factor are useful for fish farmers in selecting suitable species or varieties of faster growth rate for culture and estimating its duration for producing marketable size of the shrimps (Piratheepa, Edrisinghe, & Chitradivelu, 2013).

Like other freshwater bodies, Sungai Batang River in South Kalimantan Province also plays important role particularly for fisheries and aquaculture. A great attention has been devoted to describe the morphometric characteristic of some endemic fish species such as snakehead (Ahmadi, 2018b), *mystus nigriceps* (Ahmadi, 2018c), and three spot gourami (Aminah & Ahmadi, 2018). Whereas *Macrobrachium* species, locally called "udang sapit" was still poorly studied (Ahmadi & Rizani, 2012). It can be distinguished clearly from other shrimp species in term of cheliped traits, which are longer than its body length (Figure 1). To more understand the biological aspect and its implication for fisheries management, we investigated ⁴ the length-weight relationship, growth patterns and condition factor of this species, as well as provided some recommendation for future research in this area of study.

⁴⁷ 2. Materials and Methods

Study Site

Field survey was carried out in Sungai Batang River, South Kalimantan (Figure 2), located on 03°22'36 S and 114°49'29 E ⁵ determined by GPS 60 (Garmin Co. Ltd.,

Taiwan). *Macrobrachium* species were caught by fishermen using the wire stage-trap (heart-shaped, 65 cm high and 45 cm wide with 4 cm wide opening of entrance slit, and a piece of coconut used as bait). The traps were deployed at shallow water along the riverbank before sunset and retrieve the next morning. According to fishermen, the number of daily catch was low (1-3 shrimp per trap). Sampling activity was conducted on February to March 2019.

Data collection and Statistical Analysis

A total of 328 individuals of *Macrobrachium* species consisted of 83 males (25%) and 245 females (75%) with the sex ratio of 1:3 were investigated. Male has a pair of long second pereopods (chelipeds) that were rather dark in color with a pair of the small movable fingers of the leg. Male had chelae about 1.5 times longer than female in the same length size. Rostrum was over half of total carapace length and bent in the middle with 11 dorsal teeth and 8 ventral teeth. The upper eyes of male were parallel with the maximum height of rostrum, while those of female were slightly beneath the rostrum lines. Male and female had a pair of second antenna about 1.2 and 1.7 times longer than its chelae measured at 90 mm TL. The genital pores of the male were situated at the base of the fifth walking legs, while those females were at the base of the third walking legs. Distinct black bands located on the dorsal side at the junctions of the fifth abdominal segments. The tip of its telson reaches distinctly beyond the posterior spines of the telson. The catches were counted, identified and measured for total length (TL), carapace length (CL), abdominal length (AL), telson length (TsL), chelae length (ChL), and weight (W). Total length was measured from the tip of the rostrum to the tip of the telson to the nearest mm, keeping the abdomen fully stretched. Carapace length

was measured from the anterior tip of the rostrum to the posterior edge of the carapace; Abdominal length (AL) was measured from the anterior margin of the first abdominals somite to tip of the telson; Telson length (TsL) was measured from the posterior end of the carapace to the base of the telson; Chelae length was from the base of the first pair of walking legs to the anterior tip the pincers. The body weight was measured by Digital balance with precision of 0.01 g (Dretec KS-233, Japan). The sexes were determined by visual examination of chelae second pereopods or chelipeds. The size distribution of catch was set at class interval of 5 mm length and of 2 g weight, and the number of catch was stated in percentage. The length-weight relationship of shrimp was expressed in the allometric form (Li *et al.*, 2016):

$$W = aL^b \quad (1)$$

The length-weight relationship of shrimp can also be represented in logarithmic equation:

$$\text{Log } W = \text{Log } a + b \text{ Log } L \quad (2)$$

Where: W is the weight (g), L is the total length (mm), a is the constant showing the initial growth index and b is the slope showing growth coefficient. If shrimp retains the same shape, it grows isometrically ($b = 3$). When weight increases more than length ($b > 3$), it shows positively allometric. When the length increases more than weight ($b < 3$), it indicates negatively allometric (Li *et al.*, 2016). The proper fit of the growth model was given by determination coefficient (R^2). The regression coefficient (r) of morphological variables between male and female was figured by the square regression. The class interval for each 5 mm total length and 2 g weight was also computed regardless the sex. The previous length-weight relationship data were used for

calculating the Fulton's condition factor of shrimp by using the following formula (Gayaniilo & Pauly, 1997):

$$K = 100(W/L^3) \quad (3)$$

Where K is the Fulton's condition factor, L is total length (cm) and W is weight (g). The factor of 100 is used to bring K close to a value of one. Relative condition factor (Kn) was further estimated by following Le Cren (1951) formula:

$$Kn = W/\wedge W \quad (4)$$

Where Kn is relative condition factor reflecting 'Condition', 'fatness' or well-being of shrimp, W is the observed weight and $\wedge W$ is the calculated weight derived from length-weight relationship. The metric indicates that the higher the Kn value the better the condition of shrimp. The t-test was applied for comparing the body sizes and condition factor between male and female. All tests were analyzed at the 0.05 level of significance using SPSS-16 software.

3. Results

The body sizes, growth patterns, and condition factors of *Macrobrachium* species from Sungai Batang River were presented in Table 1. There was significantly difference between male and female ($P < 0.05$). The size of male ranged from 71 to 100 mm (88.28 ± 5.87) TL and from 4 to 12 g (8.40 ± 1.56) weight; while the size of female ranged from 60 to 100 mm (82.96 ± 6.41) TL and from 2 to 10 g (6.17 ± 1.46) weight. The pooled mean and standard deviation were 84.30 ± 6.68 mm TL and 6.73 ± 1.78 g weight. Total length, carapace length, abdominal length, and chelae length of male were considerably longer than those of female ($P < 0.001$) as described in Table 2. The mean total length of male (88.28 ± 5.87 mm) was significantly higher than that of female

(82.96 ± 6.41 mm). The carapace length of male varied from 30 to 50 mm (38.48 ± 3.12); while the carapace length of female ranged from 25 to 45 mm (35.59 ± 3.26). Male had abdomen slightly longer than female. The mean abdominal length of male and female were 36.19 ± 7.11 mm (20-53 mm) and 33.74 ± 4.47 mm (16-52 mm) respectively. There was no significant difference in the telson length between male and female ($P > 0.05$). Male had chelae about 1.5 times longer than female of the same size. The chelae length of male ranged from 75 to 148 mm (98.60 ± 15.41); while the chelae length of female ranged from 35 to 95 mm (63.26 ± 13.19). During sampling period, we also collected 26 females carrying the eggs (77.27 ± 6.37 mm TL and 5.11 ± 1.66 g weights) and they are also incorporated in statistical analysis as the whole.

The values of observed length and corresponding weight of *Macrobrachium* species are plotted in Figure 3A. The slopes (b) of the regressions were less than 3 for both the sexes, hence growth of the individual sex in river was found to be negative allometric, even for the pooled data, which means that the length increases more than weight. The estimated b values given in the allometric equations were 1.94 for male ($W = 0.0014TL^{1.9379}$) and 2.14 for female ($W = 0.0005TL^{2.1408}$); with the R^2 value falls between 0.435 and 0.449 respectively. The correlation coefficient (r) of male and female was 0.659 and 0.670, found to be higher than 0.5, showing the length-weight relationship was positively correlated. The value of $b = 2.44$ for the pooled also departs significantly from isometry ($W = 0.0001TL^{2.4433}$, $R^2 = 0.509$). Data on the length-weight relationship were presented in Table 1. Further analysis showed that male had the W/TL ratio significantly higher than female ($P < 0.001$). The relationship between total length and W/TL ratio was plotted in Fig. 3B. The W/TL ratio of male ranged from 0.054 to 0.133 (0.09 ± 0.02) and that of female ranged from 0.023 to 0.113 (0.07 ± 0.01). The

CL/TL ratio of male ranged from 0.378 to 0.549 (0.44 ± 0.03), while the CL/TL ratio of female varied from 0.295 to 0.533 (0.43 ± 0.02). Morphologically carapace length of male was longer than that of female, however, the CL/TL ratios between the sexes were not statistically significant difference ($P > 0.05$). Both male and female being equal in the ratio of abdominal length to the total length was observed ($P > 0.05$). The AL/TL ratio of male ranged from 0.288 to 0.494 (0.41 ± 0.04), while the AL/TL ratio of female ranged from 0.246 to 0.591 (0.41 ± 0.04). Apparently, male's telson length was equal to female has; however, the ratio of telson length to total length (TsL/TL) between the sexes was dimensionally differ. The mean TsL/TL ratios of male and female were 0.15 ± 0.02 ($0.108 - 0.200$) and 0.16 ± 0.02 ($0.114 - 0.231$) respectively. Male had the ChL/TL ratio greater than female ($P < 0.001$). The ChL/TL ratio of male ranged from 0.825 to 1.648 (1.12 ± 0.17), while the ChL/TL ratio of female ranged from 0.452 to 1.188 (0.76 ± 0.13). Those comparative values of body size ratios can be seen in Table 3.

The regression plots of the transformed data on the length of body parts (CL, AL, TsL, ChL) and total length relationship for the sexes indicated a linear relationship between the two variables (Figures. 4A-D). There was a variation in the intercept (a) and slope (b) of model-fitting with linear regression. The a values varied between -3.328 and 1.056, while the b values fallen between 0.479 and 2.141. The R^2 values obtained from these relationships ranged from 0.051 to 0.497, and the respective logarithmic equations and its attributes given for each relationship were presented in Table 3.

The mean Fulton's condition factor (K) of male was significantly higher than that of female ($P < 0.001$). The K values of male ranged from 1.175 to 1.276 (1.22 ± 0.23), while the K values of female ranged from 1.052 to 1.107 (1.08 ± 0.21). Male also had

the relative condition factor (Kn) greater than female ($P < 0.01$). The Kn values of male ranged from 0.981 to 1.049 (1.01 ± 0.16), while the K values of female ranged from 0.934 to 0.978 (0.96 ± 0.17). In other words, the condition factor of *Macrobrachium* species was statistically significant increase corresponding to length classes. The detail estimated parameters (the main body sizes and its aspect ratios, Fulton's condition factor and relative condition factor), as well as the significance test for the said parameters between male and female were described in Table 4.

From fishing point of view, the highest number of catches was found between 90-94 mm TL (38.55%) for male and between 80-84 mm TL (32.24%) for female (Figure 5A). To facilitate the largest amount of shrimp caught, weight was recorded at the interval class of 9-10 g (46.99%) for male and of 5-6 g (45.71%) for female (Figure 5B). None of shrimp catch over 104 mm TL and 12 g weight was observed. The other way, we found no male was captured at the size of less than 70 mm TL or between 1-2 g weights. On the other hand, *Macrobrachium* species in the present study was mostly caught by the wire stage-trap in the middle size classes compared to lower or higher classes. The t-test analysis showed that there was no significant difference in individual class interval of length-weight size range between male and female ($P > 0.05$), except for class interval of 11-12 g weight represented by male ($P < 0.0001$).

4. Discussion

There was sexual dimorphism in the sizes of male and female. In present study, male *Macrobrachium* species was significantly larger and heavier than female. Male with larger chelae win competitive interactions for shelter and food (Capelli & Munjal, 1982). Meanwhile the size of individual species had a significant effect on energy

utilization efficiency on a linear basis (Glencross, 2008). On the other side, the female ratio was ¹² 3 times higher than male caught in the investigated area, indicating that female was more active than male in term of spawning migration during wet season. Although the traps are female-biased gear, male ratio in catch would increase when population density decreases and competitive interaction are often. The similar results were also documented for *M. malcolmsonii* in ⁶¹ Indus River, Pakistan (Soomro *et al.*, 2012) or *M. felicinium* ⁸ in the lower Taylor Creek, Nigeria (Kingdom, Hart, Erundu, & Kwen, 2014). In contrast, no significant difference between the sexes observed for *M. lar* in the stream ² of Andaman and Nicobar Islands (Sethi, Ram, & Venkatesan, 2012) and *Metapenaeus dobsoni* in the Vishakhapatnam coast (Murthy & Ramaseshaiah, 1996).

The most important result of this study was *Macrobrachium* species from Sungai Batang River grew ¹ negatively allometric, where the *b* values (1.94–2.14) were significantly lower than the critical isometric value ($b < 3$). ⁶ Females have a greater regression slope than males. ¹⁸ A negative allometric growth condition in the present study was also reported for female *M. lamarrei* (Ara, Nobil, Ahmed, & Fatema, 2014), *M. dux* (Arimoro & Meye, 2007), female *M. malcolmsonii* (Hossain *et al.*, 2012), *M. lar* (Sethi, Ram, & Venkatesan, 2012) or *P. monodon* (Komi, Wilson, & Francis, 2017). It was contrary to *M. vollenhovenii* (Nwosu & Wolfi, 2006), *M. rosenbergii* (Lalrinsanga *et al.*, 2012) or *Litopenaeus vannamei* (Gautam *et al.*, 2014) that grew positively allometric, as well as *M. felicinium* (Okayi & Iorkyaa, 2004) or male *Acetes japonicas* (Wong, Ong, & Khoo, 2015) that displayed isometric growth pattern (see Table 5). Although male had the mean carapace size substantially greater than female, however, it was found not to be correlated with the growth pattern since both male and female grew

negatively isometric. According to Li *et al.* (2016), the big carapace was not conducive to effective growth.

The sizes of *Macrobrachium* species collected from the Sungai Batang River was also similar to *M. hainanense* in Chinese streams (Mantel & Dudgeon, 2004) and *M. Lamarrei* in Nepal rivers (Sharma & Subba, 2005), but it was smaller than *Macrobrachium* species in Barito River (Ahmadi & Rizani, 2012). The W/TL ratio of *Macrobrachium* species in the present study was higher than *M. vollehovenii* (Kingdom, Hart, Erondy, & Kwen, 2014), *M. macrobrachion* (Enin, 1994), *Pandalus borealis* (Wieland, 2002), or *Litopenaeus vannamei* (Khademzadeh & Haghi, 2017), but it was lower than *M. lar* (Sethi, Ram, & Venkatesan, 2012), *P. monodon* (Komi, Wilson, & Francis, 2017) or *Acetes japonicas* (Wong, Ong, & Khoo, 2015). The weight-length relationships are not constant over the entire year and vary according to factors such as food availability, feeding rate, gonad development and spawning period, fecundity, temperature, salinity, and inherited body shape (Li *et al.*, 2016).

The values of Fulton's condition factor ($K = 1.08-1.22$) of *Macrobrachium* species in the present study agreed favorably with the values recorded for other shrimp species from different geographical areas (Table 4). The K values were greater than *M. malcolmsonii* from Pakistan (Soomro *et al.*, 2012), *M. macrobrachion* from Nigeria (Kingdom, Hart, Erondy, & Kwen, 2014) or *Litopenaeus vannamei* from Iran (Khademzadeh & Haghi, 2017), but those were lower than *M. felicinum* from Nigeria (Okayi & Iorkyaa, 2004) or *P. monodon* from South China Sea (Li *et al.*, 2016). The K value greater than 1 indicating *Macrobrachium* species in Sungai Batang River were in good condition (Le-Cren, 1951), suggested that result of this study was valid. Our sampling period coincided with the reproductive period of this species that was between

February and March indicated by female carrying the eggs caught by the traps. It was also consistent with our previous work in Barito River (Ahmadi, 2018a).¹ Variation in the value of the mean K may be attributed to the biological interaction involving intraspecific competition for food and space (Arimoro & Meye, 2007) between species including sex, stages of maturity, condition of the stomach contents and availability of food (Uddin, Ghosh, & Maity, 2016; Komi, Wilson, & Francis, 2017).

It is interesting to note that *Macrobrachium* species caught by the wire stage-trap from Sungai Batang River were identical to those captured by the lighted traps from Barito River (Ahmadi, 2018a). It is quite reasonable because both rivers were geographically connected one to another. For both species, the W/TL, CL/TL and ChL/TL ratios⁶ of males were considerably higher than those of females. The principal differences were found in the total length, chelae length and body color. The mean body sizes of male and female *Macrobrachium* species from Sungai Batang River were 88.28 ± 5.87 mm TL and 82.96 ± 6.41 g weight; while those of Barito River were 66.2 ± 14.17 mm TL and 49.7 ± 11.68 g weight. Males *Macrobrachium* species from Sungai Batang River had chelae about 1.5 times longer than females; while those of Barito River had chelae about⁵ 1.3 times longer than females of the same size. Moreover, the body color of *Macrobrachium* species from Sungai Batang River was much brighter than *Macrobrachium* species from Barito River with high turbid water due to sedimentation. Recording of individual length and weight data of *Macrobrachium* species from Sungai Batang River should be continued to provide more detailed statistical analysis on possible differences in the⁴⁸ length-weight and the length-length relationships, and also condition factor³ at different developmental stages. This study presents the first reference for growth patterns and relative condition factor of *Macrobrachium* species sampled

from Sungai Batang River. From fishing practices point of view, the local fishermen were still unsatisfied with the current number of shrimp catch collected by the baited wire stage-trap, therefore it is recommended to use the lighted traps for catching *Macrobrachium* species as they were successfully applied in Barito River, and the results are still open for discussion.

5. Conclusion

Macrobrachium species in Sungai Barito River grew negatively allometric and were in good condition; males were significantly larger and heavier than females. The b values were generally in good agreement with the results obtained from other geographical areas. This study provides a useful method in assessing the well-being and growth performance, as well as conservation and management measures for this species.

Acknowledgments

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. Many thank Mrs. Siti Ubaidah for her assistance during sampling period and measurement. The authors also thank reviewers for significantly improving the contents of manuscript at publishable level.

References

- Ahmadi & Rizani, A. (2012). Catch efficiency of Low-powered incandescent light and LED light traps fishing in Barito River of Indonesia. *Kasertart University Fisheries Research Bulletin*, 36(3), 1-15.
- Ahmadi & Kristina, E. (2017). Catching efficiency of multifilament trammel nets with different twisted twine for shrimp fishing in Kotabaru, Indonesia. *Ege Journal of Fisheries and Aquatic Science*, 34(4), 363-368.

- Ahmadi. (2018a). Morphometric characteristic of *Macrobrachium* sp. from Barito River, Indonesia. *International Journal of Innovative Studies in Aquatic Biology and Fisheries*, 4(2), 19-31.
- Ahmadi. (2018b). The length-weight relationship and condition factor of the threatened Snakehead (*Channa striata*) from Sungai Batang River, Indonesia. *Polish Journal of Natural Sciences*, 33(4), 607-623.
- Ahmadi. (2018c). Length-weight relationship and condition factor of the *Mystus nigriceps* from Sungai Batang River, Indonesia. *International Journal of Innovative Studies in Aquatic Biology and Fisheries*, 4(3), 26-35.
- Aminah, S., & Ahmadi. (2018). Experimental fishing with LED light traps for Three-spot gourami (*Trichogaster trichopterus*) in Martapura, Indonesia. *International Journal of Fisheries and Aquatic Studies*, 6(1), 37-42.
- Ara, M. G., Nobil, M. N., Ahmed, Z. F., & Fatema, M. K. (2014). Length-weight relationship and growth pattern inference of a small indigenous freshwater prawn, *Macrobrachium lamarrei* (H. Milne-Edwards, 1837) in Bangladesh. *Research in Agriculture, Livestock, and Fisheries* 1(1), 137-145.
- Arimoro, F. O., & Meye, J. A. (2007). Some aspects of the biology of *Macrobrachium dux* (Lenz, 1910) (Crustacea: Decapoda: Natantia) in river Orogodo, Niger delta, Nigeria. *Acta Biologica Colombiana*, 12(1), 111-122.
- Bissaro, F. G., Gomes-Jr, J. L., & Beneditto, A. P. M. D. (2013). Morphometric variation in the shape of the cephalothorax of shrimp *Xiphopenaeus kroyeri* on the east coast of Brazil. *Journal of the Marine Biological Association of the United Kingdom*, 93(3), 683-691.
- Cadrin, S. X. (2000). Advances in morphometric identification of fishery stock. *Review of Fish Biology and Fisheries*, 10, 91-112.
- Capelli, G. M., & Munjal, B. L. (1982). Aggressive interactions and resource competition in relation to species displacement among crayfish of the genus *Orconectes*. *Journal of Crustacean Biology*, 2, 486-492
- Castilho, A. L., Gavio, M. A., Costa, R. C., Boschi, E. E., Bauer, R. T., & Fransozo, A. (2007). Latitudinal variation in population structure and reproductive pattern of the endemic South American shrimp *Artemesia longinaris* (Decapoda: Penaeoidea). *Journal of Crustacean Biology*, 27, 548-552.
- Chu, K. H., Chen, Q. C., Huang, L. M., & Wong, C. K. (1995). Morphometric analysis of commercially important penaeid shrimps from the Zhejiang estuary, China. *Fisheries Research*, 23, 83-93.

- Daud, S. K., & Ang, K. J. (1995). Selection of broodstock of tiger prawn, *Penaeus monodon* Fabricius, on the basis of morphometric traits. *Pertanika Journal of Tropical Agricultural Science*, 18(1), 15-20.
- Dineshbabu, A.P. (2006). Length-weight relationship and growth of the speckled shrimp *Metapenaeus monoceros* (Fabricius) off Saurashtra. *Journal of Marine Biological Association of India*, 48(2), 180-184.
- Enin, U. I. (1994). Length-weight parameters and condition factor of two West African prawns. *Review of Hydrobiology Tropical*, 27(2), 121-127.
- Farias, E. G. G., Pereira-Júnior, A. C., Domingos, M. M., & Dantas, D. V. (2019). Proposed bycatch-reduction modifications of shrimp fyke nets used in South American lagoons. *Acta Ichthyologica et Piscatoria*, 49(1), 1-7.
- Gayanilo, F. C., & Pauly, D. (1997). FAO-ICLARM Stock Assessment Tools (FiSAT). FAO Computerised Information Series (fisheries) No. 8, Rome, p. 262.
- Gautam, K, Nazar, A. R., Anand, E., Ganesh, Mahendran, S., & Mahadevan, G. (2014). Study of length and weight relationship of *Litopenaeus vannamei* (Boone, 1931) from East Coast of India. *International Journal of Science Inventions Today*, 3(4), 365-376.
- Glencross, B. (2008). A factorial growth and feed utilization model for Barramundi, *Lates calcarifer* based on Australian production conditions. *Aquaculture Nutrition* 14(4), 360-373.
- Hossain, M. Y., Ohtomi, J., Jaman, A., Jasmine, S. & Vadas Jr, R. L. (2012). Life history traits of the monsoon river prawn *Macrobrachium malcolmsonii* (Milne-Edwards, 1844) (Palaemonidae) in the Ganges (Padma) river, northwestern Bangladesh. *Journal of Freshwater Ecology*, 27(1): 131-142.
- Khademzadeh, O., & Haghi, M. (2017). Length-weight relationship and condition factor of white leg shrimp *Litopenaeus vannamei* (Boone, 1931) in culture systems of Choebdeh, West-South of Iran. *International Journal of Fisheries and Aquatic Studies*, 5(1), 298-301.
- Kingdom, T., Hart, A. I., Erundu, E. S., & Kwen, K. (2014). Morphology and condition indices of *Macrobrachium* species in the in the lower Taylor Creek, Niger Delta, Nigeria. *International Journal of Fisheries and Aquatic Studies*, 1(6), 95-103.
- Klassen, J. A., Gawlik, D. E., & Botson, B. A. (2014). Length-weight and length-length relationships for common fish and crayfish species in the Everglades, Florida, USA. *Journal of Applied Ichthyology*, 30, 564-566.
- Komi, Wilson, G., & Francis, A. (2017). Length-weight Relationship, condition factor and aspects of growth parameters of the Black Tiger Shrimp (*Penaeus monodon*)

- in the Andoni River System, Niger Delta, Nigeria. *Global Journal of Science Frontier Research: D Agriculture and Veterinary*, 17(2), 9-18.
- Kumar, K., Kotnala, C. B., & Rana, A. R. (2014). Morphometric study on *Macrobrachium assamense peninsulare* (Tiwari, 1958) from Rawasan stream of Garhwal Himalaya, Uttarakhand, India. *International Journal of Advanced Research*, 2(8), 317-322.
- Lalrinsanga, P. L., Pillai, B. R., Patra, G., Mohanty, S., Naik, N. K., & Sahu, S. (2012). Length weight relationship and condition factor of giant freshwater prawn *Macrobrachium rosenbergii* (De Man, 1879) based on developmental stages, culture stages and sex. *Turkish Journal of Fisheries and Aquatic Sciences*, 12, 917-924.
- Lawal-Are, A. O., & Akinjogunla, V. F. (2012). *Penaeus notialis* (Pink Shrimps): Length-weight relationships and condition factor in Lagos Lagoon, South West, Nigeria. *Science and Technology*, 2(3), 32-40.
- Le-Cren, E. D. (1951). The Length-weight relationship and seasonal cycle in gonad-weight and condition in the Perch (*Perca fluviatilis*). *Journal of Animal Ecology*, 20, 201-219.
- Li, Y., Zhou, F., Ma, Z., Huang, J., Jiang, S., Yang, Q., Li, T., & Qin, J. G. (2016). Length-weight relationship and condition factor of giant tiger shrimp, *Penaeus monodon* (Fabricius, 1798) from four breeding families. *SpringerPlus*, 5(1279), 2-5. DOI 10.1186/s40064-016-2979-6.
- Mantel, S.K., & Dudgeon, D. (2004). Growth and production of a tropical predatory shrimp, *Macrobrachium hainanense* (Palaemonidae) in two Hong Kong streams. *Freshwater Biology*, 49, 1320-1336.
- Murthy, V. S. R., & Ramaseshaiah, M. (1996). Length-weight relationship and other conversion factors in *Metapenaeus dobsoni* (Miers) from the Vishakhapatnam Coast. *Indian Journal of Fisheries*, 43(2), 179-185.
- Ng, P. K. L. (2017). Collecting and processing freshwater shrimps and crabs. *Journal of Crustacean Biology*, 37(1), 115-122, <https://doi.org/10.1093/jcbiol/ruw004>.
- Nwosu, F. M., & Wolfi, M. (2006). Population dynamics of the Giant African River Prawn *Macrobrachium vollenhovenii* Herklots 1857 (Crustacea, Palaemonidae) in the Cross River Estuary, Nigeria. *West Africa Journal of Applied Ecology*, 9, 1-14.
- Okayi, G., & Iorkyaa, A. (2004). Length-weight relationship and condition of freshwater shrimps *Atya gabonensis* and *Macrobrachium felicinium* from the Mu river, Makurdi, Nigeria. *Animal Research International*, 1(3), 153-154.

- Olawusi-Peters Olamide, O., Ajibare Adefemi, O., & Bello-Olusoji Oluayo, A. (2014). Length-weight relationship and condition factor of shrimps in coastal waters of Ondo state, South West, Nigeria. *International Journal of Fisheries and Aquatic Studies*, 1(3), 137-142.
- Piratheepa, S., Edrisinghe, U., & Chitravadivelu, K. (2013). Investigation on Length-Weight Relationship of *Penaeus monodon* (Fabricius, 1798) in Kakkaithevu coastal waters in the Northern part of Sri Lanka. *Tropical Agricultural Research*, 25(1), 133-140.
- Pravin & Ravindran. (2011). Catch efficiency of gillnets in shrimp filtration farms at Vypeen Island, Kerala, South India. *Indian Journal of Fisheries*, 58(2), 153-157.
- Sethi, S. N., Ram, N., & Venkatesan, V. (2012). Length-weight relationship of *Macrobrachium lar* (Fabricius, 1798), an endemic freshwater prawn in streams and ponds of Andaman and Nicobar Islands. *Indian Journal of Fisheries*, 59(4), 157-161.
- Sharma, A., & Subba, B. R. (2005). General biology of freshwater prawn *Macrobrachium lamarrei* (H. Milne-Edwards) of Biratnagar, Nepal. *Our Nature*, 3, 31-41.
- Soomro, A. N., Baloch, W. A., Chandio, T. J., Achakzai, W. M., & Saddozai, S. (2012). Condition factor and length-weight relationship of Monsoon River Prawn *Macrobrachium malcolmsonii malcolmsonii* (H. Milne-Edwards, 1844) (Palaemonidae) in Lower Indus River. *Pakistan Journal of Zoology*, 44(5), 1279-1283.
- Sousaa, R., Gouveia, L., Pinto, A. R., Timóteoa, V., Delgado J., & Henriques, P. (2019). Weight-length relationships of six shrimp species caught off the Madeira Archipelago, Northeastern Atlantic. *Brazilian Journal of Biology*, 79(1), 133-138.
- Tsoumani, M., Liasko, R., Moutsaki, P., Kagalou, I., & Leonardos, I. (2006). Length-weight relationship of an invasive cyprinid fish (*Carassius gibelio*) from 12 Greek lakes in relation to their trophic states. *Journal of Applied Ichthyology*, 22, 281-284.
- Uddin, S. K. N., Ghosh, S., & Maity, J. (2016). Length weight relationship and condition factor of *Penaeus monodon* (Fabricius, 1798) from Digha coast, West Bengal, India. *International Journal of Fisheries and Aquatic Studies*, 4(3), 168-172.
- Wieland, K. (2002). A new length-weight Relationship for northern shrimp (*Pandalus borealis*) off West Greenland (NAFO Subareas 0+1). Scientific Council Meeting, November 2002. NAFO SCR Doc.02/144 Serial No. N4773. pp. 1-4.

Wong, B. Y., Ong, H. K. A., & Khoo, G. (2015). Length-weight relationships of *Acetes* spp. sampled along the West Coast of Peninsular Malaysia. *Sains Malaysiana*, 44(3), 379-386.

Table 1. Body sizes, growth patterns, and condition factors of *Macrobrachium* species sampled from Sungai Batang River.

| Sex | n | Total length (mm) | | Weight (g) | | Allometric equations | a | b | R ² | r | Growth pattern | K | | Kn | |
|--------|-----|-------------------|-----|--------------|-----|----------------------|-------------|--------|----------------|-------|----------------|-----|-------------|-------------|-----------|
| | | Min | Max | Mean ± SD | Max | | | | | | | Min | Mean ± SD | Mean ± SD | Mean ± SD |
| Male | 83 | 71 | 100 | 88.28 ± 5.87 | 4 | 12 | 8.40 ± 1.56 | 0.0014 | 1.938 | 0.435 | 0.659 | A- | 1.22 ± 0.24 | 1.01 ± 0.16 | |
| Female | 245 | 60 | 100 | 82.96 ± 6.41 | 2 | 10 | 6.17 ± 1.46 | 0.0005 | 2.141 | 0.449 | 0.670 | A- | 1.08 ± 0.21 | 0.96 ± 0.17 | |
| Pooled | 328 | 60 | 100 | 84.30 ± 6.68 | 2 | 12 | 6.73 ± 1.78 | 0.0001 | 2.443 | 0.509 | 0.714 | A- | 1.12 ± 0.23 | 0.97 ± 0.16 | |

a = constant, b = exponent, R² = coefficient of determination, r = coefficient of correlation, A = allometric, K = Fulton's condition factor, Kn = relative condition factor

Table 2. Statistic parameters of main body parts observed for *Macrobrachium* species from Sungai Batang River.

| Sex | Statistic Parameters | TL | W | CL | AL | TsL | ChL | K | Kn |
|--------|----------------------|--------|--------|--------|--------|--------|--------|--------|--------|
| Male | N | 83 | 83 | 83 | 83 | 83 | 83 | 83 | 83 |
| | Minimum | 71 | 4 | 30 | 20 | 10 | 75 | 0.88 | 0.68 |
| | Maximum | 100 | 12 | 50 | 53 | 18 | 148 | 2.51 | 1.66 |
| | Mean | 88.28 | 8.40 | 38.48 | 36.19 | 13.65 | 98.60 | 1.22 | 1.01 |
| | Std. Deviation | 5.87 | 1.56 | 3.12 | 7.11 | 1.88 | 15.41 | 0.24 | 0.16 |
| | Std. Error Mean | 0.64 | 0.17 | 0.34 | 0.78 | 0.21 | 1.69 | 0.03 | 0.02 |
| Female | N | 245 | 245 | 245 | 245 | 245 | 245 | 245 | 245 |
| | Minimum | 60 | 2 | 25 | 16 | 10 | 35 | 0.29 | 0.27 |
| | Maximum | 100 | 10 | 45 | 52 | 20 | 95 | 2.31 | 1.58 |
| | Mean | 82.96 | 6.17 | 35.60 | 33.74 | 13.62 | 63.26 | 1.08 | 0.96 |
| | Std. Deviation | 6.41 | 1.46 | 3.26 | 4.47 | 1.83 | 13.19 | 0.21 | 0.17 |
| | Std. Error Mean | 0.41 | 0.09 | 0.21 | 0.28 | 0.12 | 0.84 | 0.01 | 0.01 |
| Pooled | Sig. (2-tailed) | 0.0000 | 0.0000 | 0.0000 | 0.0003 | 0.9114 | 0.0000 | 0.0000 | 0.0058 |

TL = total length, W = weight, CL = carapace length, AL = abdominal length, TsL = telson length, ChL = chelae length,

K = Fulton's condition factor, Kn = relative condition factor

Table 3. Descriptive statistic and estimated parameters of length-weight and length-length relationships of *Macrobrachium* species from Sungai Batang River.

| Sex | n | Total length (mm) | | Weight (g) | | Log W | Logarithmic equations | a | b | R ² | r | Range of W/TL | Mean±SD W/TL |
|--------|-----|-------------------|--------|-----------------------|--------|---------|-----------------------|---------|-------|----------------|-------|-----------------|----------------|
| | | Min | Max | log TL | log TL | | | | | | | | |
| Male | 83 | 71 | 100 | 1.94 ± 0.03 | 4 | 12 | 0.93 ± 0.09 | -2.853 | 1.938 | 0.435 | 0.659 | 0.054 - 0.133 | 0.09 ± 0.02 |
| Female | 245 | 60 | 100 | 1.92 ± 0.03 | 2 | 10 | 0.78 ± 0.11 | -3.3278 | 2.141 | 0.449 | 0.670 | 0.023 - 0.113 | 0.07 ± 0.01 |
| Sex | n | Total length (mm) | | Carapace length (mm) | | Log CL | Logarithmic equations | a | b | R ² | r | Range of CL/TL | Mean±SD CL/TL |
| Min | Max | log TL | log TL | Min | Max | | | | | | | | |
| Male | 83 | 71 | 100 | 1.94 ± 0.03 | 30 | 50 | 1.58 ± 0.04 | 0.134 | 0.745 | 0.392 | 0.626 | 0.378 - 0.549 | 0.44 ± 0.03 |
| Female | 245 | 60 | 100 | 1.92 ± 0.03 | 25 | 45 | 1.55 ± 0.04 | -0.061 | 0.840 | 0.496 | 0.704 | 0.295 - 0.533 | 0.43 ± 0.03 |
| Sex | n | Total length (mm) | | Abdominal length (mm) | | Log AL | Logarithmic equations | a | b | R ² | r | Range of AL/TL | Mean±SD AL/TL |
| Min | Max | log TL | log TL | Min | Max | | | | | | | | |
| Male | 83 | 71 | 100 | 1.94 ± 0.03 | 20 | 53 | 1.55 ± 0.06 | -1.209 | 1.421 | 0.467 | 0.684 | 0.282 - 0.494 | 0.41 ± 0.04 |
| Female | 245 | 60 | 100 | 1.92 ± 0.03 | 16 | 52 | 1.52 ± 0.06 | -0.904 | 1.266 | 0.497 | 0.705 | 0.246 - 0.591 | 0.41 ± 0.04 |
| Sex | n | Total length (mm) | | Telson length (mm) | | Log TsL | Logarithmic equations | a | b | R ² | r | Range of TsL/TL | Mean±SD TsL/TL |
| Min | Max | log TL | log TL | Min | Max | | | | | | | | |
| Male | 83 | 71 | 100 | 1.94 ± 0.03 | 10 | 18 | 1.13 ± 0.06 | -0.362 | 0.767 | 0.138 | 0.371 | 0.108 - 0.200 | 0.15 ± 0.02 |
| Female | 245 | 60 | 100 | 1.92 ± 0.03 | 10 | 20 | 1.13 ± 0.06 | -0.594 | 0.899 | 0.260 | 0.510 | 0.114 - 0.231 | 0.16 ± 0.02 |
| Sex | n | Total length (mm) | | Chelae length (mm) | | Log ChL | Logarithmic equations | a | b | R ² | r | Range of ChL/TL | Mean±SD ChL/TL |
| Min | Max | log TL | log TL | Min | Max | | | | | | | | |
| Male | 83 | 71 | 100 | 1.94 ± 0.03 | 75 | 148 | 1.99 ± 0.06 | 1.056 | 0.479 | 0.051 | 0.225 | 0.825 - 1.648 | 1.12 ± 0.17 |
| Female | 245 | 60 | 100 | 1.92 ± 0.03 | 35 | 95 | 1.79 ± 0.09 | -1.614 | 1.776 | 0.411 | 0.641 | 0.452 - 1.188 | 0.76 ± 0.13 |

a = constant, *b* = exponent, *R*² = coefficient of determination, *r* = coefficient of correlation

Table 4. Estimated parameters and significance test for LWR and its ratios of *Macrobrachium* species from Sungai Batang River

| Estimated Parameters | Levene's Test for Equality of Variances | | | t-test for Equality of Means | | | | | | |
|---------------------------|---|-------|-----|------------------------------|-----------------|-----------------------|---------|---------|---|--|
| | F | Sig. | df | Sig. (2-tailed) | Mean Difference | Std. Error Difference | Lower | Upper | 95% Confidence Interval of the Difference | |
| Total Length (TL) | 1.510 | 0.220 | 326 | 0.000 | 5.3179 | 0.7972 | 3.7496 | 6.8862 | | |
| Weight (W) | 0.428 | 0.513 | 326 | 0.000 | 2.2302 | 0.1890 | 1.8584 | 2.6021 | | |
| Carapace length (CL) | 0.018 | 0.893 | 326 | 0.000 | 2.8860 | 0.4100 | 2.0794 | 3.6926 | | |
| Abdominal length (AL) | 34.015 | 0.000 | 326 | 0.000 | 2.4540 | 0.6678 | 1.1403 | 3.7677 | | |
| Telson length (TsL) | 0.007 | 0.934 | 326 | 0.911 | 0.0261 | 0.2344 | -0.4350 | 0.4872 | | |
| Chelae length (ChL) | 0.024 | 0.876 | 326 | 0.000 | 35.3412 | 1.7502 | 31.8980 | 38.7844 | | |
| Fulton's condition factor | 0.011 | 0.918 | 326 | 0.000 | 0.1462 | 0.0278 | 0.0914 | 0.2009 | | |
| Relative condition factor | 0.271 | 0.603 | 326 | 0.006 | 0.0584 | 0.0211 | 0.0170 | 0.0999 | | |
| W/TL | 0.287 | 0.593 | 326 | 0.000 | 0.0209 | 0.0018 | 0.0173 | 0.0246 | | |
| CL/TL | 0.426 | 0.514 | 326 | 0.060 | 0.0069 | 0.0037 | -0.0003 | 0.0141 | | |
| AL/TL | 0.771 | 0.381 | 326 | 0.611 | 0.0026 | 0.0051 | -0.0075 | 0.0127 | | |
| TsL/TL | 1.068 | 0.302 | 326 | 0.000 | -0.0096 | 0.0025 | -0.0144 | -0.0047 | | |
| ChL/TL | 5.626 | 0.018 | 326 | 0.000 | 0.3603 | 0.0183 | 0.3243 | 0.3963 | | |

Table 5. Comparative parameters of length-weight relationship and condition factors of *Macrobrachium* species from Sungai Batang River and other shrimp species from different geographical areas.

| Species | Country | n | Sex | W/TL | a | b | R ² | K | Growth Type | References |
|--------------------------|------------|------|-----|--------|--------|-------|----------------|------|-------------|---|
| <i>Macrobrachium</i> sp. | Indonesia | 83 | M | 0.952 | 0.001 | 1.938 | 0.435 | 1.22 | A- | Present study |
| <i>Macrobrachium</i> sp. | Indonesia | 245 | F | 0.744 | 0.001 | 2.141 | 0.449 | 1.08 | A- | Present study |
| <i>Macrobrachium</i> sp. | Indonesia | 161 | M | 0.710 | 0.000 | 3.210 | 0.975 | 1.39 | A+ | Ahmadi, 2018 |
| <i>Macrobrachium</i> sp. | Indonesia | 175 | F | 0.322 | 0.000 | 2.734 | 0.937 | 1.14 | A- | Ahmadi, 2018 |
| <i>M. lamarrei</i> | Bangladesh | 1018 | P | ... | 0.000 | 2.924 | 0.970 | - | A- | Ara, Nofri, Ahmed & Fatema, 2014 |
| <i>M. malcolmsonii</i> | Pakistan | 62 | M | 5.464 | 0.003 | 2.97 | 0.97 | 0.61 | A- | Soomro <i>et al.</i> , 2012 |
| <i>M. malcolmsonii</i> | Pakistan | 47 | F | 2.779 | 0.009 | 3.04 | 0.94 | 0.86 | I | Soomro <i>et al.</i> , 2012 |
| <i>M. felicinum</i> | Nigeria | 55 | P | 1.043 | 0.002 | 3.003 | 0.998 | 2.03 | I | Okayi & Iorkyaa, 2004 |
| <i>M. felicinum</i> | Nigeria | 132 | M | 0.372 | 0.009 | 3.23 | 0.92 | 1.23 | A+ | Kingdom, Hart, Erundu & Kwen, 2014 |
| <i>M. felicinum</i> | Nigeria | 726 | F | 0.308 | 0.010 | 2.12 | 0.75 | 1.23 | A- | Kingdom, Hart, Erundu & Kwen, 2014 |
| <i>M. vollenhovenii</i> | Nigeria | 246 | M | 0.820 | 0.007 | 3.29 | 0.98 | 1.26 | A+ | Kingdom, Hart, Erundu & Kwen, 2014 |
| <i>M. vollenhovenii</i> | Nigeria | 94 | F | 0.605 | 0.007 | 3.25 | 0.98 | 1.26 | A+ | Kingdom, Hart, Erundu & Kwen, 2014 |
| <i>M. vollenhovenii</i> | Nigeria | 1070 | M | ... | 0.000 | 3.483 | 0.993 | 1.24 | A+ | Nwosu & Wolfi, 2006 |
| <i>M. vollenhovenii</i> | Nigeria | 1358 | F | ... | 0.000 | 3.329 | 0.990 | 1.25 | A+ | Nwosu & Wolfi, 2006 |
| <i>M. macrobrachion</i> | Nigeria | 352 | P | 0.299 | 0.006 | 3.280 | 0.968 | 1.13 | A+ | Enin, 1994 |
| <i>M. macrobrachion</i> | Nigeria | 36 | M | 0.240 | 0.014 | 2.71 | 0.84 | 1.06 | A- | Kingdom, Hart, Erundu & Kwen, 2014 |
| <i>M. macrobrachion</i> | Nigeria | 61 | F | 0.420 | 0.011 | 2.96 | 0.98 | 1.06 | A- | Kingdom, Hart, Erundu & Kwen, 2014 |
| <i>M. macrobrachion</i> | Nigeria | 27 | P | ... | -1.797 | 1.443 | 0.730 | 0.68 | A- | Olawusi-Peters Olamide <i>et al.</i> , 2014 |
| <i>M. dux</i> | Nigeria | 138 | M | 0.990 | 0.769 | 0.839 | 0.706 | 2.37 | A- | Arimoro & Meye, 2007 |
| <i>M. dux</i> | Nigeria | 166 | F | 0.971 | 0.708 | 0.861 | 0.739 | 1.43 | A- | Arimoro & Meye, 2007 |
| <i>M. rosenbergii</i> | India | 273 | M | 9.046 | -2.718 | 3.550 | 0.958 | 1.14 | A+ | Lalrinsanga <i>et al.</i> , 2012 |
| <i>M. rosenbergii</i> | India | 460 | F | 11.260 | -2.273 | 3.244 | 0.969 | 1.05 | A+ | Lalrinsanga <i>et al.</i> , 2012 |
| <i>M. lar</i> | India | 110 | M | 2.083 | 0.000 | 2.453 | 0.769 | - | A- | Sethi, Ram & Venkatesan, 2012 |

| | | | | | | | | | | |
|------------------------------|-----------------|------|---|-------|--------|-------|-------|------|----|---|
| <i>M. lar</i> | India | 128 | F | 1.971 | 0.000 | 2.622 | 0.839 | - | A- | Sethi, Ram & Venkatesan, 2012 |
| <i>Penaeus monodon</i> | India | 242 | M | 7.661 | 0.077 | 2.409 | 0.778 | 1.38 | A- | Uddin, Ghosh & Maity, 2016 |
| <i>P. monodon</i> | India | 391 | F | 5.256 | 0.091 | 2.388 | 0.847 | 1.48 | A- | Uddin, Ghosh & Maity, 2016 |
| <i>P. monodon</i> | Sri Langka | 98 | M | 1.202 | 0.006 | 3.016 | 0.958 | - | I | Piratheepa, Edrisinghe & Chitravadivelu, 2013 |
| <i>P. monodon</i> | Sri Langka | 153 | F | 1.317 | 0.005 | 3.075 | 0.963 | - | I | Piratheepa, Edrisinghe & Chitravadivelu, 2013 |
| <i>P. monodon</i> | South China Sea | 210 | M | 2.308 | 0.023 | 2.803 | 0.901 | 2.71 | A- | Li <i>et al.</i> , 2016 |
| <i>P. monodon</i> | South China Sea | 202 | F | 2.313 | 0.026 | 2.764 | 0.892 | 2.42 | A- | Li <i>et al.</i> , 2016 |
| <i>P. monodon</i> | Africa | 190 | M | 2.151 | 0.019 | 2.911 | 0.911 | 2.49 | A- | Li <i>et al.</i> , 2016 |
| <i>P. monodon</i> | Africa | 168 | F | 2.174 | 0.023 | 2.795 | 0.901 | 1.83 | A- | Li <i>et al.</i> , 2016 |
| <i>P. monodon</i> | Nigeria | 376 | P | 2.528 | -0.157 | 1.460 | 0.957 | 0.69 | A- | Komi, Wilson & Francis, 2017 |
| <i>P. notialis</i> | Nigeria | 413 | M | 1.714 | 0.539 | 0.619 | 0.302 | 1.38 | A- | Lawal-Are & Akinjogunla, 2012 |
| <i>P. notialis</i> | Nigeria | 187 | F | 1.788 | 0.397 | 0.817 | 0.394 | 1.34 | A- | Lawal-Are & Akinjogunla, 2012 |
| <i>Pandalus borealis</i> | West Greenland | 1225 | P | 0.313 | 0.000 | 3.057 | 0.972 | - | I | Wieland, 2002 |
| <i>Plesionika ensis</i> | Portugal | 81 | P | 3.679 | 0.053 | 2.609 | 0.950 | - | A- | Sousaa <i>et al.</i> , 2019 |
| <i>Acetes japonicas</i> | Malaysia | 25 | M | 5.486 | 0.002 | 3.153 | 0.659 | - | I | Wong, Ong & Khoo, 2015 |
| <i>A. japonicas</i> | Malaysia | 49 | F | 2.810 | 0.017 | 2.432 | 0.798 | - | A- | Wong, Ong & Khoo, 2015 |
| <i>Acetes indicus</i> | Malaysia | 264 | M | 1.197 | 0.010 | 2.694 | 0.907 | - | A- | Wong, Ong & Khoo, 2015 |
| <i>A. indicus</i> | Malaysia | 340 | F | 0.886 | 0.008 | 2.778 | 0.917 | - | A- | Wong, Ong & Khoo, 2015 |
| <i>Metapenaeus monoceros</i> | India | 190 | M | 1.426 | 0.009 | 2.924 | 0.907 | 1.80 | A- | Dineshbabu, 2006 |
| <i>M. monoceros</i> | India | 173 | F | 5.269 | 0.003 | 3.292 | 0.972 | 2.00 | A+ | Dineshbabu, 2006 |
| <i>Litopenaeus vannamei</i> | India | 157 | M | 0.481 | 0.001 | 3.260 | 0.999 | - | A+ | Gautam <i>et al.</i> , 2014 |
| <i>L. vannamei</i> | India | 156 | F | 2.839 | 0.000 | 3.654 | 0.998 | - | A+ | Gautam <i>et al.</i> , 2014 |
| <i>L. vannamei</i> | Iran | 634 | P | 0.758 | 0.011 | 2.694 | 0.889 | 0.56 | A- | Khademzadeh & Haghi, 2017 |

a = constant, *b* = exponent, R^2 = coefficient of determination, A+ = positive allometric, A- = negative allometric, I = isometric, K = Fulton's condition factor




Figure 1. *Macrobrachium* species from Sungai Batang River, Indonesia
CL = carapace length, AL = abdominal length, TsL = telson length,
ChL = chelae length.




Figure 2. Map showing the location of sampling site in Sungai Batang River, Indonesia





Figure 3. The length-weight relationship [A] and the length-ratio of W/TL [B] of male and female *Macrobrachium* species



27
Figure 4. The relationship between the body sizes and total length of male and female *Macrobrachium* species




Figure 5. Number of catches, length size [A] and weight size [B] distribution of male and female *Macrobrachium* species

Growth Patterns and Relative Condition Factor of Macrobrachium Species from Sungai Batang River, Indonesia

ORIGINALITY REPORT

18%

SIMILARITY INDEX

14%

INTERNET SOURCES

13%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to Lambung Mangkurat University **2%**
Student Paper
- 2** eprints.cmfri.org.in **1%**
Internet Source
- 3** Gopalakrishnan, A., M. Rajkumar, M. M. Rahman, J. Sun, P. J. Antony, B. A. Venmathi maran, and J. P. Trilles. "Length-weight relationship and condition factor of wild, grow-out and 'loose-shell affected' giant tiger shrimp, *Penaeus monodon* (Fabricius, 1798) (Decapoda: Penaeidae)", *Journal of Applied Ichthyology*, 2014. **1%**
Publication
- 4** journal.bdfish.org **1%**
Internet Source
- 5** Ahmadi .. "An Introduction of Light Traps for Sampling Freshwater Shrimp and Fish in the Barito River, South Kalimantan", *Journal of Fisheries and Aquatic Science*, 2012 **1%**
Publication

| | | |
|----|--|------|
| 6 | hdl.handle.net Internet Source | 1 % |
| 7 | asianfisheriessociety.org Internet Source | 1 % |
| 8 | www.fisheriesjournal.com Internet Source | 1 % |
| 9 | Jurniati, D Arfiati, A Maizar, A Kurniawan. " Length weight relationships and condition factor of sweet river prawn, (Thalwitss, 1891) in the downstream Rongkong watershed ", IOP Conference Series: Earth and Environmental Science, 2020 Publication | 1 % |
| 10 | www.researchsquare.com Internet Source | <1 % |
| 11 | nopr.niscpr.res.in Internet Source | <1 % |
| 12 | www.ncbi.nlm.nih.gov Internet Source | <1 % |
| 13 | Leticia Félix-Cuencas, Juan Fernando García-Trejo, Samuel López-Tejeida, Jesús Josafat de León-Ramírez et al. "Nitrogen and phosphorus flux in wastewater from three productive stages in a hyperintensive tilapia culture", Journal of Water Reuse and Desalination, 2021 | <1 % |

-
- 14 aquaticcommons.org Internet Source <1 %
-
- 15 topsecretapiaccess.dovepress.com Internet Source <1 %
-
- 16 www.journalijar.com Internet Source <1 %
-
- 17 Michael B. New. "Prawn Capture Fisheries and Enhancement", *Freshwater Prawn Culture*, 12/12/2000
Publication <1 %
-
- 18 eprints.nottingham.ac.uk Internet Source <1 %
-
- 19 www.bioflux.com.ro Internet Source <1 %
-
- 20 www.jsmcentral.org Internet Source <1 %
-
- 21 Ambika Dhakal, Bharat Raj Subba. "Length-Weight Relationship of *Lepidocephalichthys guntea* of Pathri Khola, Morang District", *Our Nature*, 1970
Publication <1 %
-
- 22 Elsayed F. E. Farrag, Ahmed M. Alzaabi. "Length-weight, condition factor and reproductive aspects of the cobia, *Rachycentron canadum* (Linnaeus, 1766) in

the Arabian Gulf off the United Arab Emirates", Egyptian Journal of Aquatic Biology and Fisheries, 2020

Publication

23

Halimeh Zare, Neda Zareei, Leila Kohan, Negar Azarpira. "Genetic polymorphisms of MiR-499a (rs3746444) and MiR-196a2 (rs11614913) in ischemic stroke and correlation with risk factors", Gene Reports, 2022

Publication

<1 %

24

R. Kindong, X.-J. Dai, S.-Q. Tian, C.-X. Gao. "Ichthyofaunal Assemblage of the Dianshan Lake, Shanghai, China: Implications for Management", Proceedings of the Zoological Society, 2017

Publication

<1 %

25

Submitted to University of Stirling

Student Paper

<1 %

26

irgu.unigoa.ac.in

Internet Source

<1 %

27

macrothink.org

Internet Source

<1 %

28

www.allresearchjournal.com

Internet Source

<1 %

29

www.blackmeditjournal.org

Internet Source

<1 %

| | | |
|----|---|------|
| 30 | Submitted to Ateneo de Manila University Student Paper | <1 % |
| 31 | journalarticle.ukm.my Internet Source | <1 % |
| 32 | mospace.umssystem.edu Internet Source | <1 % |
| 33 | Damian Frank, Sue Poole, Stephanie Kirchhoff, Ciarán Forde. " Investigation of Sensory and Volatile Characteristics of Farmed and Wild Barramundi () using Gas Chromatography–Olfactometry Mass Spectrometry and Descriptive Sensory Analysis ", Journal of Agricultural and Food Chemistry, 2009 Publication | <1 % |
| 34 | nottingham-repository.worktribe.com Internet Source | <1 % |
| 35 | repositorio.unal.edu.co Internet Source | <1 % |
| 36 | zoologyandecology.gamtc.lt Internet Source | <1 % |
| 37 | Lazola Bangani, Hlekani Muchazotida Kabiti, Oseni Amoo, Motebang D.V. Nakin, Zendy Magayiyana. "Impacts of illegal solid waste dumping on the water quality of the Mthatha River", Water Practice & Technology, 2023 | <1 % |

38

www.nbfgr.res.in

Internet Source

<1 %

39

www.trjfas.org

Internet Source

<1 %

40

Lota A. Creencia, Lutgardo B. Alcantara, Riza G. San Juan. "Bamboo as Potential Substrate for Sea Cage Culture of Abalone *Haliotis asinina*", Research Square Platform LLC, 2023

Publication

<1 %

41

Nazek Jisr, Ghassan Younes, Carol Sukhn, Mohammad H. El-Dakdouki. "Length-weight relationships and relative condition factor of fish inhabiting the marine area of the Eastern Mediterranean city, Tripoli-Lebanon", The Egyptian Journal of Aquatic Research, 2018

Publication

<1 %

42

assets.nrdc.org

Internet Source

<1 %

43

fison.org.ng

Internet Source

<1 %

44

pdfs.semanticscholar.org

Internet Source

<1 %

45

Ayza, O.. "Estimation of size at onset of sexual maturity and growth parameters in Norway

<1 %

lobster (*Nephrops norvegicus*) off the Portuguese coast", Fisheries Research, 201102

Publication

46

San San Aye. " Population structure and growth characteristics of oriental river prawn, (De Haan, 1849) in Soon Lun Reservoir, Mandalay Region, Myanmar ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

47

jurnal.usk.ac.id

Internet Source

<1 %

48

zsp.com.pk

Internet Source

<1 %

49

A. A. Atici, A. Sepil, F. Sen. "Biological Characteristics of the Invasive *Gambusia holbrooki* in the Wetland, the Lake Van Basin, Turkey", Journal of Ichthyology, 2022

Publication

<1 %

50

Koshal Kumar, Chandra Bhanu Kotnala, Anita Rawat Rana. " Fecundity and egg dimension of freshwater prawn (Tiwari, 1958) from Rawasan stream of Garhwal, Central Himalaya, India ", Aquaculture Research, 2019

Publication

<1 %

51

T Bratel, G Hedenstierna, O Nyquist, E Ripe. "The use of a vasodilator, felodipine, as an adjuvant to long-term oxygen treatment in

<1 %

COLD patients", European Respiratory Journal, 1990

Publication

52

akwafoto.aip.pl

Internet Source

<1 %

53

banglajol.info

Internet Source

<1 %

54

biodiversitas.mipa.uns.ac.id

Internet Source

<1 %

55

drs.cift.res.in

Internet Source

<1 %

56

wajfas.unilag.edu.ng

Internet Source

<1 %

57

www.ajol.info

Internet Source

<1 %

58

www.sapub.org

Internet Source

<1 %

59

www.zsp.com.pk

Internet Source

<1 %

60

A. Arshad, S.M. Nurul Amin, Y.L.Z. Nuradiella, Z.C. Cob, R. Ara, D. Aziz. "Population Characteristics of *A. japonicus* from the Kedah Coastal Waters of Peninsular Malaysia", *Journal of Fisheries and Aquatic Science*, 2012

Publication

<1 %

61

Md. Ashekur Rahman, Md. Yeamin Hossain, Mostafa A. R. Hossain, Zoarder Faruque Ahmed, Ahmed Jaman, Jun Ohtomi. "Are there any effects of eco-climatic factors on freshwater prawn reproduction: a study on monsoon river prawn, *Macrobrachium malcolmsonii*", *Environmental Science and Pollution Research*, 2022

Publication

<1 %

62

Hiram Herrera-Barquín, Antonio Leija Tristán, Carlos Javier Aguilera-González. "Crecimiento, factor de condición y condición relativa del camarón *Tozeuma carolinense* (Decapoda: Caridea: Hippolytidae) en la región meridional de la laguna Madre, Tamaulipas, México", *Revista Mexicana de Biodiversidad*, 2022

Publication

<1 %

63

Hui Yu Li, Lin Lin Yang, Jia Hua Cheng, Jian Zhong Ling. "Length-weight relationships of thirteen penaeoid shrimps in the East China Sea", *Crustaceana*, 2018

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On