

2022-
Antioxidative_response.pdf
by


Submission date: 05-Jun-2023 08:21PM (UTC+0700)

Submission ID: 2109499360

File name: 2022-Antioxidative_response.pdf (718.46K)

Word count: 6081

Character count: 33475



Antioxidative responses in the skin mucus of *Periophthalmodon schlosseri* as biomarkers for the assessment of heavy metal pollution in the coastal wetlands of Kuala Lupak estuary of the Barito River, Indonesia

¹Heri B. Santoso, ¹Hidayaturrahmah, ²Eko Suhartono, ³Rizmi Yunita, ⁴Danang Biyatmoko, ⁵Dewi Anggraini

¹ Department of Biology, Faculty of Mathematics and Natural Sciences, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia; ² Department of Medical Chemistry/Biochemistry, Faculty of Medicine, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia; ³ Department of Aquatic Resources Management, Faculty of Fisheries and Marine Sciences, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia; ⁴ Department of Animal Science, Faculty of Agriculture, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia; ⁵ Department of Statistics, Faculty of Mathematics and Natural Sciences, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia. Corresponding author: E. Suhartono, ekoantioxidant@gmail.com

Abstract. One of the main factors that concern the well-being of estuarine water ecosystems is heavy metal pollution by anthropogenic activities. Heavy metals can cause oxidative stress in fish by increasing the formation of reactive oxygen species (ROS) and, thus, directly disrupt the antioxidant defense system. This article aimed to determine the responses of antioxidant enzymes in the skin mucus of *Periophthalmodon schlosseri* as a biomarker of oxidative stress. *P. schlosseri* or the giant mudskipper is a typical fauna of the contaminated Barito River estuary. The study showed that the activity of the antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GSH-Px) was higher in the skin mucus of the mudskipper from estuary waters compared to that from coastal waters, while the level of H₂O₂ was lower. According to *in silico* studies, heavy metals can act as activators for binding Nrf2-KEAP1, causing antioxidant expression to be disrupted. The activity of antioxidant enzymes such as SOD, CAT, and GSH-Px in *P. schlosseri* skin mucus is the potential to be used as an oxidative stress biomarker in non-invasive biomonitoring assessment of heavy metal pollution in estuaries. The giant mudskipper has the potential to be a sentinel species and a sensitive bioindicator in monitoring/assessing heavy metal pollution in estuaries for ecotoxicological studies in polluted water bodies.

Key Words: antioxidative defense system, biomonitoring, heavy metal pollution, *in silico*, oxidative stress biomarkers, skin mucus.

Introduction. Estuaries and other coastal environments have been subjected to heavy metal pollutants and their derivatives. However, the toxicity and effects of these pollutants are still unknown (Salgado et al 2019). Heavy metal pollution is a major environmental issue that harms the health of the biota and the aquatic environment. In aquatic organisms, the presence and exposure to non-essential heavy metals (Hg, Pb, Cd) and/or essential heavy metals (Zn, Cu, Fe) produces oxidative stress and adverse reactions. Anthropogenic activities primarily cause heavy metal pollution in the aquatic environment (Adeogun et al 2020). Settlements, industrial activities, fisheries, agriculture, and coal transportation on land along the Barito River have the potential to produce heavy metal waste harmful to the aquatic environment. Therefore, biomonitoring measures are required for early detection of the detrimental effects of pollution.

There is insufficient data on biomonitoring heavy metal pollution at the Barito river estuary. Invasive monitoring of the Barito River estuary quality reveals that heavy metals Cd, Pb, Hg, Fe, Zn, and Cu in the water bodies, sediments, liver, and kidneys of *Periophthalmodon schlosseri* surpasses the prescribed quality standards. Non-invasive biomonitoring using fish mucus skin as a biological matrix produced the same results (Santoso et al 2021a). Heavy metal accumulation in fish tissues might be viewed as an indirect marker of heavy metal exposure in aquatic ecosystems. Monitoring/evaluating fish tissue exposure to heavy metals acts as an early warning indicator of poor water quality and heavy metal pollution of sediment, allowing necessary actions to be carried out to protect public health and the environment (El-Shenawy et al 2021).

Fish are extremely vulnerable to heavy metal-induced oxidative stress. Heavy metal-mediated oxidative damage is a complex process, in which redox-active metals including Fe, Cu, Cr, and V generate ROS through the redox cycle. Metals with no redox potential, such as Hg, Ni, Pb, and Cd, can interfere with antioxidant defenses (Sevcikova et al 2011). These heavy metals can inhibit antioxidant gene expression through the NRF2 (the nuclear factor erythroid 2-related factor 2), KEAP1 (Kelch-like ECH-associated protein1), and ARE (the antioxidant responsive element) pathways, resulting in decreased antioxidant activity. As a result, ROS levels increase and the balance of cellular antioxidant activity shifts (Adeogun et al 2020).

Fish have complex antioxidant enzyme defense mechanisms that prevent excessive ROS and oxidative damage, including superoxide dismutase (SOD), catalase (CAT), glutathione S-transferase (GST), and glutathione peroxidase (GPx) (Soliman et al 2019). This defensive mechanisms assist in mitigating the harmful effects of ROS produced by heavy metal metabolism. Antioxidant enzymes such as SOD, CAT, and GPx are useful as heavy metal pollution biomarkers in water (El-Shenawy et al 2021). Antioxidants of low molecular weight, such as glutathione (GSH) and vitamin A, have also been shown to neutralize ROS. Endogenous enzymatic and non-enzymatic antioxidants are required for the conversion of ROS to innocuous metabolites to restore metabolism and cellular function (Orbea et al 2002; Nnamdi et al 2015; Pastorino et al 2020).

Prior studies on *P. schlosseri* exposed to heavy metals revealed damage to liver and kidney cells (Santoso et al 2021b), indicating a failure of antioxidant enzyme activity to reduce excessive quantities of ROS or free radicals. Excess ROS or free radicals produce oxidative stress, which leads to the oxidation of biological membranes, decreased cell stability, cell damage, disease, and death (Magara et al 2021). The appearance of antioxidant enzymes can show cell damage produced by the oxidation of biological membranes, allowing researchers to establish the mechanism of heavy metals contamination in *P. schlosseri*. Given the information provided above, further studies are needed to establish the potential of *P. schlosseri* as a bioindicator species of heavy metal contamination in the Indonesian Barito river estuary. The concentrations of heavy metals Fe, Zn, Cu, Hg, Cd, and Pb in water bodies, sediments, and skin mucus of mudskipper have been previously studied and published by Santoso et al (2021a), but the interaction description between heavy metals and antioxidant enzymes is not defined. The purpose of this article was to assess the activity of antioxidant enzymes in *P. schlosseri* skin mucus in response to oxidative stress due to exposure to heavy metals and to examine the visualization of interactions between heavy metals and antioxidant enzyme activity *in silico*. Changes in the parameters of antioxidant enzyme activity in fish skin mucus can be used as biomarkers of oxidative stress for non-invasive biomonitoring of heavy metal pollution in estuaries.

Material and Method

Description of the study sites. The longest and largest river in South Kalimantan is the Barito River. The Barito River starts from the Schwaner Mountains and flows approximately 1000 km from the Central Kalimantan region in the northern part of Borneo to the river's estuary in the Java Sea. The river is 650 to 800 m wide and 8 m deep. The width of the river in the funnel-shaped estuary reaches 1000 m, making the Barito River Indonesia's widest river. Starting from the upper parts of the river, the

longest section of the Barito River is in Central Kalimantan, while the remainder, up to the river mouth, is in South Kalimantan. Before reaching the mouth of the Barito River in the Barito Kuala region, this river meets the estuary of the Negara River. This study was carried out in the estuary waters (station 1) and the coastal waters of Kuala Lupak (station 2), which are both located along the west coast of the Barito River estuary (Figure 1). Table 1 shows a description of the research locations for each station.




Figure 1. Map of research locations and stations.

Description of stations 1 and 2

Table 1

No	Station	Description
1	Estuary/River estuary	Kuala Lupak is a Wildlife Reserve Area with a mangrove (swamp forest) environment. There are residential areas around the mouth of the Kuala Lupak River. The Kuala Lupak River flows into the Java Sea. Samples were collected in the estuary waters at 3°27'28.223"S; 114°22'04.780"E.
2	Coastal waters	Kuala Lupak's coastal waters are about 8 km west of the mouth of the Barito River, located on the coast of the Java Sea with a coastline of 30 km. Mangrove swamp forests can be found along the waters of Kuala Lupak's coastline. The samples were collected in coastal waters near the river's mouth at 3°28'02.383"S; 114°21'31.282"E.

Water, sediment, and *P. Schlosseri* were collected between February 2-3, 2021, using the "purposive sampling" method. The stations were selected based on the source of activities suspected of producing pollution loads. The settlements in the area justify sampling estuaries and coastal waters, since it is believed that human activity has contaminated the Barito River's water flow. Landslides, sedimentation, flooding, and other natural occurrences are also believed to play a role in polluting. As a result of

anthropogenic activities in the river's upstream and intermediate watersheds, which will have an impact on the estuary area, the samplings further consider that estuaries and coastal locations are the most vulnerable to biodegradation.

Sampling. At each site, 10 *P. schlosseri* were caught. Their average body length was 15.47 ± 0.84 cm, and the average mass was 150.95 ± 3.41 g. Skilled fishermen manually caught the fish by inserting their hands into the fish burrows. To remove contaminants, water was used to clean the fish from each location. The skin mucus of *P. schlosseri* was prepared using the method described by Fernández-Alacid et al (2018). Clean fish were placed on a tray, and the skin mucus was carefully collected using a sterile glass slide by gently rubbing from the region behind the gills down to the caudal fin. A sterile slide was softly slid down on the animal's flanks two or three times, and the skin mucus was then delicately pushed and gathered in a sterile tube (2 mL). A large portion of mucus might be collected by repeatedly touching the skin, but doing so is not recommended because it could lead to epidermal lesions and contaminate the samples with blood or other cells. This method must be carried out precisely, without rewetting the animal and without any contact with the operculum, ventral, anal, or caudal fins, to prevent mucus dilution by the water. For further analysis, the mucus was immediately frozen in liquid nitrogen and preserved at -80°C .

All procedures were carried out following the ethical standards given by the Health Research Ethics Committee No. 549/KPEK-FK ULM/EC/III/2021 Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia, which was formed to control and regulate the animal experiments.

Evaluation of antioxidative stress in the skin mucus of *P. schlosseri*. Antioxidant levels were determined by measuring the activities of SOD, CAT, and GSH-Px. The supernatant was prepared from skin mucus samples in 10 volumes of 0.1 M Tris-EDTA buffer (pH 7.4), centrifuged at $1000 \times g$ at 4°C for 30 min. The kit's instructions were then followed to determine the various oxidative stress/antioxidant parameters. The supernatant was collected for enzyme activity analysis. SOD activity was measured spectrophotometrically at 560 nm, referencing Misra & Fridovich (1977). One unit of SOD is defined as the amount of enzyme required to prevent 50% of epinephrine autooxidation. SOD activity is measured in units of SOD mg^{-1} protein. CAT activity was determined in the samples using the approach of Aebi (1984), by monitoring the enzyme's decrease in H_2O_2 absorbance at 240 nm. For 3 min, the decrease in absorbance was recorded, and the change was calculated in nmol of H_2O_2 decomposed/min/mg protein (Lakra et al 2021). H_2O_2 levels were measured using ammonium molybdate according to Suhartono et al (2013), who used hydrogen peroxide as a standard. The 100 μL of the supernatant sample was stirred with 100 μL of 32.4 mM ammonium molybdate. After 10 min incubation, the absorbance was measured at 405 nm wavelength with a UV-Visible spectrophotometer. The activities of GSH-Px were detected using the GSH-Px Activity Assay Kit (Elabscience®) according to the manufacturer's instructions. All antioxidant enzyme activities were measured at 37°C with a Shimadzu UV-2100 spectrophotometer.

Ligand and protein preparation. Through *in silico* studies, the interaction between proteins involved in antioxidant activity and ligands in the form of heavy metals can be described. Fe^{2+} , Zn^{2+} , Cu^{2+} , Hg^{2+} , and Cd^{2+} ligands were obtained from MIB: Metal Ion-Binding site prediction and server docking (<http://bioinformo.cmu.edu.tw/MIB/>) (Pratidina et al 2022; Lahdimawan et al 2022). The previous study of Santoso et al (2021a) provided the basis for using these heavy metals as ligands. Meanwhile, the protein used as a target was KEAP1 obtained from the RCSB Protein Data Bank (<https://www.rcsb.org/search>) with PDB code: 5cgj (Sari & Bare 2021). The protein was prepared by removing the natural ligand residue present in the protein. The Chimera 1.15 program (<https://www.cgl.ucsf.edu/chimera/download.html>) was used to prepare the ligands and the protein.

Analysis methods. Mean and standard deviation (SD) were used to analyze the distribution and variability of antioxidant enzyme activity and hydrogen peroxide level. The Kruskal Wallis test, with a 5% significance value, was performed to assess whether there is a significant median difference between the activity of antioxidant enzymes and the levels of hydrogen peroxide in two stations (estuary and coastal waters). Docking data are processed and visualized using the Chimera 1.15 program. Visualization is used to explain the interaction between ligands and receptor protein residues, specifically the interacting amino acids and the distance between the receptor protein ligands (Pratidina et al 2022; Lahdimawan et al 2022).

Results and Discussion

Antioxidants biomarkers in skin mucus *P. schlosseri*. Antioxidant enzyme activity was recorded in fish skin mucus in response to the accumulation of heavy metals such as Fe, Zn, Cu, Hg, Cd, and Pb in water bodies, sediments, and fish skin mucus. The heavy metal accumulation data has been previously published (Santoso et al 2021a). Fish skin mucus from estuary waters had an average activity of the SOD enzyme (39.3 ± 1.6 unit of SOD mg^{-1} protein) higher than that of fish from coastal waters (20.46 ± 5.4 unit of SOD mg^{-1} protein). The average CAT enzyme activity in fish skin mucus from estuary waters (18.59 ± 4.82 $\mu\text{mol min}^{-1} \text{mg protein}^{-1}$) was higher than that from fish from coastal waters (8.8 ± 2.94 $\mu\text{mol min}^{-1} \text{mg protein}^{-1}$). The average activity of the GSH-Px enzyme in fish skin mucus from estuary waters (42.7 ± 4.65 $\text{nmol min}^{-1} \text{mg protein}^{-1}$) was higher than that of fish from coastal waters (28.43 ± 5.53 $\text{nmol min}^{-1} \text{mg protein}^{-1}$). Meanwhile, the average level of H_2O_2 in fish skin mucus from estuary waters (11.95 ± 2.38 mmol g^{-1}) was lower than that of fish from coastal waters (17.83 ± 6.21 mmol g^{-1}) (Table 2). The findings of nonparametric statistical tests (Kruskal Wallis) revealed that antioxidant enzyme activity and H_2O_2 levels in fish skin mucus from the estuary and coastal waters differed significantly (Figure 2). Table 2 reveals that the CAT and H_2O_2 enzymes have statistically insignificantly different levels of activity.

Table 2
Antioxidant enzyme activity and H_2O_2 level in *Periophthalmodon schlosseri* skin mucus

Antioxidant biomarkers	Station	
	Estuary (Station 1)	Coastal waters (Station 2)
SOD	39.3 ± 1.6^b	20.46 ± 5.4^b
CAT	18.59 ± 4.82^c	8.8 ± 2.94^c
GSH-Px	42.7 ± 4.65^a	28.43 ± 5.53^a
H_2O_2	11.95 ± 2.38^c	17.83 ± 6.21^c

Note: data are presented as mean \pm SD of three replicates; n=10 samples of each station; different superscripts show significant differences at $p < 0.05$; SOD (unit of SOD mg^{-1} protein) - superoxide dismutase; CAT ($\mu\text{mol min}^{-1} \text{mg protein}^{-1}$) - catalase; GSH-Px ($\text{nmol min}^{-1} \text{mg protein}^{-1}$) - glutathione peroxidase; H_2O_2 level (mmol g^{-1}).

Heavy metals are one of the main sources of pollution in estuary water ecosystems. Several heavy metals, such as Cu, Fe, Mn, and Zn, are essential to biological processes, they can also be hazardous in excessive doses. Harmful effects from Cd, Cr, Pb, and Hg can occur even at very low amounts (Akinsanya et al 2020). Heavy metals significantly increase ROS like superoxide anions, hydroxyl radicals, singlet oxygen, and hydrogen peroxide in the fish body. This may result in cell degeneration due to damage to the cell membrane and genotoxicity. Fish can accumulate these heavy metals and experience severe oxidative stress as a result of their harmful effects, making them an effective bioindicator (Kumar et al 2021). Oxidative stress occurs due to an imbalance between pro-oxidants such as ROS and defense cellular antioxidant enzyme systems (Pastorino et al 2020). Antioxidant biomarkers are used for early detection of the harmful effects of heavy metals on several aquatic organisms (Hamed et al 2020). The first line of defense against excessive ROS brought on by heavy metals is provided by antioxidant enzymes such as SOD and CAT (Kumar et al 2021). Additionally, fish tissues and blood experience

modifications in their physiological and biochemical processes due to heavy metals. Its harmful effects result in oxidative phosphorylation blockade, glutathione enzyme decrease, inhibition of antioxidant enzymatic activity, excessive ROS production, and inhibition of cell repair mechanisms (El-Shenawy et al 2021).




Figure 2. Antioxidant enzyme activity in *Periophthalmodon schlosseri* skin mucus: (a) SOD (unit of SOD mg^{-1} protein); (b) CAT ($\mu\text{mol min}^{-1} \text{mg protein}^{-1}$); (c) GSH-Px ($\text{nmol min}^{-1} \text{mg protein}^{-1}$); (d) hydrogen peroxide (mmol g^{-1}).

Due to the presence of antioxidant enzymes like SOD, CAT, and GPx in the skin mucus of the giant mudskipper, it is possible to use this species as a bioindicator of heavy metal pollution. El-Shenawy et al (2021) explained that the function of enzymatic activity and scavengers is to eliminate excessive reactive oxygen species, such as hydrogen peroxide (H_2O_2) to prevent oxidative damage to cells. This species is a sentinel organism, able to identify the effects of heavy metals in the water and soil of estuarine ecosystems. The findings of this study suggest that environmental risk assessment studies could use oxidative stress biomarkers in the form of cellular antioxidant enzyme expression in *P. schlosseri* skin mucus. This is supported by Hamed et al (2020), who stated that fish exposed to heavy metal pollution develop antioxidant enzymes, which could serve as a biomarker of oxidative stress. Faheem & Lone (2018) confirmed that the antioxidant enzyme activity indices including SOD, CAT, and GPx are frequently used as screening tools to determine the effects of environmental stress.

In this study, the mucus of *P. schlosseri* from estuary waters had stronger antioxidant enzyme activity than that of fish from coastal waters. This is due to increased concentrations of the heavy metals Fe, Cu, Zn, Hg, Pd, and Cd in the water bodies,

sediments, and skin mucus of the mudskipper fish in the Kuala Lupak estuary waters (Santoso et al 2021a). Additionally, there was more anthropogenic influence on this site. Heavy metal accumulation in fish skin mucus results in increased SOD, CAT, and GPx antioxidant activity. Antioxidant defense, according to Prokić et al (2016), functions as a balanced and coordinated system. Heavy metals can induce oxidative stress and disrupt the balance of the antioxidant defense system. SOD includes metalloenzymes that will catalyze the removal of $O_2^{\bullet-}$ into O_2 and H_2O_2 . Additionally, CAT and/or GSH-Px efficiently convert H_2O_2 into H_2O and O_2 . After oxidative stress, SOD and CAT are the first active enzymes to exhibit alteration (Mahamood et al 2021). GSH-Px is a crucial enzyme in preventing hydrogen peroxide damage to cells (Firat et al 2021). SOD, CAT, GSH, and GPx are important antioxidant enzymes that help shield fish from the harmful effects of ROS. Superoxide dismutase is regarded as the body's first line of defense against oxidative stress, which causes cells to produce hydrogen peroxide from superoxide radicals. CAT and GPx then eliminate the hydrogen peroxide and lessen its damaging effects (Gao et al 2021). The findings of this study are consistent with Lakra et al (2021), who found that all fish tissues exposed to coal mine effluent, and coal mine waste had higher levels of SOD and CAT. Antioxidant enzymes appear in *Clarias batrachus* as an early reaction to heavy metals in coal mine effluent. Farombi et al (2007) found a significant increase in SOD and CAT activity in *Clarias gariepinus* when exposed to heavy metals Pb, Zn, Cu, Cd, and As, whereas Yogeshwaran et al (2020) found that heavy metal exposure to Zn, Cu, Cd, and Pb increased antioxidant enzyme activity and hydrogen peroxide (H_2O_2) in *Scylla serrata*. Chang et al (2021) reported that prolonged Cd exposure increases ROS synthesis, resulting in oxidative stress in fish. These exposures to Cd significantly reduced CAT, GSH-Px, and SOD activities, while increasing DA and ROS concentrations in *Cyprinus carpio*. Biomarker activity can also be influenced by abiotic factors such as pH, dissolved oxygen content, and water temperature. Aquatic species' heavy metal concentrations and antioxidant enzyme activity may fluctuate seasonally and in response to biological and environmental stressors (Pastorino et al 2020).

Heavy metal interaction with KEAP1 protein. The mechanism of heavy metals in inducing ROS formation is thought to be through the Nrf2-KEAP1 binding pathway. Heavy metals work as in the Fenton reaction or the Haber-Weiss reaction, which oxidizes H_2O_2 to $\bullet OH$ causing oxidative stress. Oxidative stress also activate various transcription factors including nuclear factor kappa-B (NF κB), p53, HIF-hypoxia-inducible factor 1 α , peroxisome proliferator-activated receptor γ (PPAR- γ), β -catenin/Wnt, and Nrf2. Protein nuclear factor erythroid 2-related (Nrf2) is a key protein in the antioxidant response through the Nrf2/KEAP1 pathway. In inactive conditions, Nrf2 binds to the Kelch protein ECH Associating Protein 1 (KEAP1). In contrast, free Nrf2 will translocate into the cell nucleus and activate the expression of antioxidant genes such as genes encoding SOD, CAT, and peroxidase (Canning et al 2015; Baird & Yamamoto 2020).

The involvement of heavy metals in reducing the expression of antioxidant enzymes is still an interesting study and the mechanism is still unclear. In this study, the interaction of heavy metals with KEAP1 protein is presented in Table 3. The interaction visualization of heavy metals with KEAP1 is presented in Figure 3.

KEAP1 is a protein that has a protein molecular weight of 69-kDa. This protein has a physiological function with the Kelch protein as an actin binder and acts as a negative regulator of Nrf2. Nrf2 further binds to KEAP1 in the cytoplasm. KEAP1 acts as a complex substrate adapter protein and targets E-ligase containing 3 cullin Nrf2. Furthermore, Nrf2 leads to degradation mediated by the proteasome (Sari & Bare 2021).

Electrophilic and oxidative stressors of Nrf2 signaling (CDDO-Im and sulforaphane) interact with residues from KEAP1 to disrupt Nrf2 and proteolytic sequestration (Sari & Bare 2021; Oktavianti et al 2022). Heavy metals are predicted to have the potential to be activators of the KEAP1 protein. The five heavy metals, namely Fe^{3+} , Zn^{2+} , Cu^{2+} , Hg^{2+} and Cd^{2+} , that interact with KEAP1 protein have the potential to activate the Nrf2 protein bound to its active side. The Nrf2 binding of the Nrf2-KEAP1

complex stimulates an increase in ROS in the cells, so the expression of antioxidant formation can be disrupted.

Table 3
Heavy metals interaction with KEAP1 residues

<i>Metals</i>	<i>Residues</i>	<i>Lengths (Å)</i>
Fe ³⁺	Ser431	2.01
Zn ²⁺	Asp538	2.60
Cu ²⁺	His553	2.25
	Cis368;	2.712
Hg ²⁺	Ile425	2.682
	Val370	2.371
	Val420	2.566
Cd ²⁺	ASP585	2.18




Figure 3. Interaction of KEAP1 to heavy metals; (a) Fe³⁺; (b) Zn²⁺; (c) Cu²⁺; (d) Hg²⁺; (e) Cd²⁺.

In Table 3, it can be seen that Hg^{2+} is chelated with 4 residues, namely Cis368, Ile425, Val370, and Val420. This binding causes Hg^{2+} to accumulate in the KEAP1 protein, thus strengthening the Nrf2-KEAP1 complex. In addition, in Table 3, it can also be seen that Fe^{3+} has the shortest bonding length. A shorter bond will increase the strength of the bond energy between Fe^{3+} -Ser431. This also allows Fe^{3+} to accumulate in KEAP1, so that antioxidant expression in the cells will be disrupted. Similarly, Zn^{2+} , Cu^{2+} , and Cd^{2+} all have the same mechanism. The results of this study are consistent with the research Komari & Suhartono (2020), who note that Cd^{2+} can interact with antioxidants SOD, CAT, and glutathione reductase, disrupting its activity.

Conclusions. Heavy metal exposure has a significant negative impact on estuarine ecosystems. Mudskippers are endangered due to heavy metal bioaccumulation. Reactive oxygen species (ROS) are produced when heavy metal accumulation is present, which can lead to an imbalance in the prooxidant or antioxidant homeostasis of the fish. This is referred to as oxidative stress, and it can be measured using a variety of biomarkers, such as alterations in the activity of antioxidant enzymes. The study's findings showed that oxidative stress caused by the bioaccumulation of heavy metals in the skin mucus of *P. schlosseri* is responded to by the appearance of cellular antioxidant enzyme activity, notably SOD, CAT, and GSH-Px as oxidative stress biomarkers. Heavy metals significantly increase the production of ROS and modify the activity of antioxidant enzymes in the mudskipper skin mucus. According to *in silico* studies, heavy metals can activate Nrf2-KEAP1 binding, which disrupts antioxidant expression. The responses of mudskipper fish skin mucus to changes in antioxidant activity are valuable for assessing the quality of estuary waters. The use of antioxidant activity biomarkers in the biomonitoring program serves as a measure to monitor the health of aquatic fauna and assess heavy metal pollution in estuary waters. Therefore, the concentration of heavy metals in water bodies, sediments, and tissues of mudskipper fish should be monitored continuously to control pollution. To maintain the biodiversity of coastal wetland ecosystems, all types of wastewater must be treated before being released into aquatic environments. The findings of this study can be used as a starting point for stakeholders to discuss and explore the appropriate method for evaluating heavy metal concentrations and developing an integrated biomonitoring program.

Acknowledgements. The authors thankfully acknowledge the Rector of Lambung Mangkurat University for providing research funding, through the PDWM program.

Conflict of Interest. The authors declare that there is no conflict of interest.

References

- Adeogun A. O., Ibor O. R., Omiwole R., Chukwuka A. V., Adewale A. H., Kumuyi O., Arukwe A., 2020 Sex-differences in physiological and oxidative stress responses and heavy metals burden in the black jaw tilapia, *Sarotherodon melanotheron* from a tropical freshwater dam (Nigeria). *Comparative Biochemistry and Physiology Part-C: Toxicology and Pharmacology* 229:108676.
- Aebi H., 1984 Catalase *in vitro*. *Methods in Enzymology* 105:121-126.
- Akinsanya B., Ayanda I. O., Fadipe A. O., Onwuka B., Saliu J. K., 2020 Heavy metals, parasitologic and oxidative stress biomarker investigations in *Heterotis niloticus* from Lekki Lagoon, Lagos, Nigeria. *Toxicology Reports* 7:1075-1082.
- Baird L., Yamamoto M., 2020 The molecular mechanisms regulating the KEAP1- NRF2 pathway. *Molecular and Cellular Biology* 40(13):e00099-20.
- Canning P., Sorrell F. J., Bullock A. N., 2015 Structural basis of Keap1 interactions with Nrf2. *Free Radical Biology and Medicine* 88(Part B):101-107.
- Chang X., Chen Y., Feng J., Huang M., Zhang J., 2021 Amelioration of Cd-induced bioaccumulation, oxidative stress, and immune damage by probiotic *Bacillus coagulans* in common carp (*Cyprinus carpio* L.). *Aquaculture Reports* 20:100678.

- El-Shenawy N. S., Gad EL-Hak H. N., Ghobashy M. A., Mansour F. A., Soliman M. F. M., 2021 Using antioxidant changes in liver and gonads of *Oreochromis niloticus* as biomarkers for the assessment of heavy metals pollution at Sharkia Province, Egypt. *Regional Studies in Marine Science* 46:101863.
- Faheem M., Lone K. P., 2018 Oxidative stress and histopathologic biomarkers of exposure to bisphenol-A in the freshwater fish, *Ctenopharyngodon idella*. *Brazilian Journal of Pharmaceutical Sciences* 53(3):e17003.
- Farombi E. O., Adelowo O. A., Ajimoko Y. R., 2007 Biomarkers of oxidative stress and heavy metal levels as indicators of environmental pollution in African catfish (*Clarias gariepinus*) from Nigeria Ogun River. *International Journal of Environmental Research and Public Health* 4(2):158-165.
- Fernández-Alacid L., Sanahuja I., Ordóñez-Grande B., Sánchez-Nuño S., Viscor G., Gisbert E., Herrera M., Ibarz A., 2018 Skin mucus metabolites in response to physiological challenges: a valuable non-invasive method to study teleost marine species. *Science of the Total Environment* 644:1323-1335.
- Firat Ö., Erol R., Firat Ö., 2021 An investigation on freshwater fish *Oreochromis niloticus* (Linnaeus, 1758): Assessing hemotoxic effects of different copper compounds used as nanomaterial or pesticide. *Bulletin of Environmental Contamination and Toxicology* 108:549-554.
- Gao X. Q., Fei F., Huang B., Meng X. S., Zhang T., Zhao K. F., Chen H. B., Xing R., Liu B. L., 2021 Alterations in hematological and biochemical parameters, oxidative stress, and immune response in *Takifugu rubripes* under acute ammonia exposure. *Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology* 243:108978.
- Hamed M., Soliman H. A. M., Osman A. G. M., Sayed A. E. D. H., 2020 Antioxidants and molecular damage in Nile tilapia (*Oreochromis niloticus*) after exposure to microplastics. *Environmental Science and Pollution Research* 27(13):14581-14588.
- Kumar E. K., Midhun S. J., Vysakh A., James T. J., 2021 Antagonistic effects of dietary *Moringa oleifera* on hemato-biochemical and oxidative stress of lead nitrate intoxicated Nile tilapia, *Oreochromis niloticus*. *Aquaculture Research* 52(12):6164-6178.
- Komari N., Suhartono E., 2020 Cadmium binding to antioxidant enzymes: in silico study. *IOP Conference Series: Materials Science and Engineering* 980:012038.
- Lahdimawan A., Bulan S. A., Suhartono E., Setiawan B., 2022 [Impact of cadmium and mercury on carbohydrate metabolism: in silico study on glycogen synthesis and phosphofructokinase enzymes]. *Jurnal Ilmiah Ibnu Sina* 7(1):109-115. [In Indonesian].
- Lakra K. C., Banerjee T. K., Lal B., 2021 Coal mine effluent-induced metal bioaccumulation, biochemical, oxidative stress, metallothionein, and histopathological alterations in vital tissues of the catfish, *Clarias batrachus*. *Environmental Science and Pollution Research* 28(20):25300-25315.
- Magara G., Elia A. C., Dörr A. J. M., Abete M. C., Brizio P., Caldaroni B., Righetti M., Pastorino P., Scoparo M., Prearo M., 2021 Metal load and oxidative stress driven by organotin compounds on rainbow trout. *Environmental Science and Pollution Research International* 28(26):35012-35022.
- Mahamood M., Javed M., Alhewairini S. S., Zahir F., Sah A. K., Ahmad M. I., 2021 *Labeo rohita*, a bioindicator for water quality and associated biomarkers of heavy metal toxicity. *Npj Clean Water* 4(1):17.
- Misra H. P., Fridovich I., 1977 Superoxide dismutase and peroxidase: a positive activity stain applicable to polyacrylamide gel electropherograms. *Archives of Biochemistry and Biophysics* 183(2):511-515.
- Nnamdi A. H., Olumide A. A., Adeladun A. E., Oyenike K., Rosemary E. I., 2015 The use of antioxidant stress enzymes, lipid peroxidation, and red blood cell abnormalities as biomarkers of stress in *Periphalmus papilio* of the polluted coastal Lagos lagoon. *Environmental Monitoring and Assessment* 187(3):103.
- Oktaviyanti I. K., Nurrasyidah I., Muthmainah N., Diany H., Komari N., Suhartono E., 2022 Supplementation of *Nigella sativa* as antioxidant in COVID-19 patients: *In silico*

- study via the Nrf2-Keap1 pathway. *International Journal of Drug Delivery Technology* 12(3):1028-1032.
- Orbea A., Ortiz-Zarragoitia M., Solé M., Porte C., Cajaraville M. P., 2002 Antioxidant enzymes and peroxisome proliferation in relation to contaminant body burdens of PAHs and PCBs in bivalve molluscs, crabs and fish from the Urdaibai and Plentzia estuaries (Bay of Biscay). *Aquatic Toxicology* 58(1-2):75-98.
- Pastorino P., Elia A. C., Caldaroni B., Menconi V., Abete M. C., Brizio P., Bertoli M., Zaccaroni M., Gabriele M., Dörr A. J. M., Pizzul M., Prearo M., 2020 Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake (Cottian Alps). *Science of the Total Environment* 715:136946.
- Pratidina E. A., Suhartono E., Setiawan B., 2022 Impact of heavy metals on hexokinase isoforms: an in silico study. *Berkala Kedokteran* 18 (1):29-35.
- Prokić M. D., Borković-Mitić S. S., Krizmanić I. I., Mutić J. J., Vukojević V., Nasia M., Gavrić J. P., Despotović J. P., Gavrilović B. R., Radovanović T. B., Pavlović S. Z., Saičić Z. S., 2016 Antioxidative responses of the tissues of two wild populations of *Pelophylax kl. esculentus* frogs to heavy metal pollution. *Ecotoxicology and Environmental Safety* 128:21-29.
- Salgado L. D., Marques A. E. M. L., Kramer R. D., de Oliveira F. G., Moretto S. L., de Lima B. A., Prodocimo M. M., Cestari M. M., de Azevedo J. C. R, de Assis H. C. S., 2019 Integrated assessment of sediment contaminant levels and biological responses in sentinel fish species *Atherinella brasiliensis* from a sub-tropical estuary in south Atlantic. *Chemosphere* 219:15-27.
- Santoso H. B., Hidayaturrehman, Suhartono E., Yunita R., Biyatmoko D., Anggraini D., 2021a Heavy metal concentrations in water, sediment and giant mudskipper (*Periophthalmodon schlosseri*) in the coastal wetlands of Kuala Lupak estuary of the Barito River, Indonesia. *AACL Bioflux* 14(5):2878-2893.
- Santoso H. B., Hidayaturrehman, Sihananto B. S., 2021b [Application of liver and kidney hispathological biomarkers of mudskipper fish (*Periophthalmodon schlosseri*) as an early warning of heavy metal lead (Pb) toxicity in the Barito River estuary]. *Proceedings of the National Wetland Environment Seminar* 6(3):1-8. [In Indonesian].
- Sari D. R. T., Bare Y., 2021 [In silico study of antioxidant activity of bioactive compounds in lemongrass oil (*Cymbopogon citratus*)]. *Al-Kimia* 9(1):61-69. [In Indonesian].
- Sevcikova M., Modra H., Slaninova A., Svobodová Z., 2011 Metals as a cause of oxidative stress in fish: a review. *Veterinarni Medicina* 56(11):537-546.
- Soliman H. A. M., Hamed M., Lee J. S., Sayed A. E. D. H., 2019 Protective effects of a novel pyrazolecarboxamide derivative against lead nitrate induced oxidative stress and DNA damage in *Clarias gariepinus*. *Environmental Pollution* 247:678-684.
- Suhartono E., Triawanti, Yunanto A., Firdaus R. T., Iskandar, 2013 Chronic cadmium hepatooxidative in rats: treatment with haruan fish (*Channa striata*) extract. *APCBEE Procedia* 5:441-445.
- Yogeshwaran A., Gayathiri K., Muralisankar T., Gayathri V., Monica J. I., Rajaram R., Marimuthu K., Bhavan P. S., 2020 Bioaccumulation of heavy metals, antioxidants, and metabolic enzymes in the crab *Scylla serrata* from different regions of Tuticorin, Southeast Coast of India. *Marine Pollution Bulletin* 158:111443.
- *** <http://bioinmfo.cmu.edu.tw/MIB/>
- *** <https://www.cgl.ucsf.edu/chimera/download.html>
- *** <https://www.rcsb.org/search>

Received: 25 June 2022. Accepted: 14 August 2022. Published online: 23 October 2022.

Authors:

Heri Budi Santoso, Department of Biology, Faculty of Mathematics and Natural Sciences, Lambung Mangkurat University, Banjarbaru, 70123 South Kalimantan, Indonesia, e-mail: heribudisantoso@ulm.ac.id

Hidatarramah, Department of Biology, Faculty of Mathematics and Natural Sciences, Lambung Mangkurat University, Banjarbaru, 70123 South Kalimantan, Indonesia, e-mail: hidatarramah@ulm.ac.id

Eko Suhartono, Department of Medical Chemistry/Biochemistry, Faculty of Medicine, Lambung Mangkurat University, Banjarbaru, 70123 South Kalimantan, Indonesia, e-mail: ekoantioxidant@gmail.com

Rizmi Yunita, Department of Aquatic Resources Management, Faculty of Fisheries and Marine Sciences, Lambung Mangkurat University, Banjarbaru, 70123 South Kalimantan, Indonesia, e-mail: rizmiyunita@ulm.ac.id

Danang Biyatmoko, Department of Animal Science, Faculty of Agriculture, Lambung Mangkurat University, Banjarbaru, 70123 South Kalimantan, Indonesia, e-mail: danangbiyatmoko@ulm.ac.id

Dewi Anggraini, Department of Statistics, Faculty of Mathematics and Natural Sciences, Lambung Mangkurat University, Banjarbaru, 70123 South Kalimantan, Indonesia, e-mail: dewi.anggraini@ulm.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Santoso H. B., Hidayatarramah, Suhartono E., Yunita R., Biyatmoko D., Anggraini D., 2022 Antioxidative responses in the skin mucus of *Periophthalmodon schlosseri* as biomarkers for the assessment of heavy metal pollution in the coastal wetlands of Kuala Lupak estuary of the Barito River, Indonesia. *AAFL Bioflux* 15(5):2500-2511.

ORIGINALITY REPORT

19%

SIMILARITY INDEX

%

INTERNET SOURCES

19%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

Nahla S. El-Shenawy, Heba N. Gad EL-Hak, Mahi A. Ghobashy, Farida A. Mansour, Maha F.M. Soliman. "Using antioxidant changes in liver and gonads of Oreochromis niloticus as biomarkers for the assessment of heavy metals pollution at Sharkia province, Egypt", *Regional Studies in Marine Science*, 2021

Publication

3%

2

Nurudeen A. Azeez, Seun Oyelami, Abideen A. Adekanmi, Opeyemi B. Ologunye et al. "Biodiesel potentials of microalgal strains isolated from fresh water environment", *Environmental Challenges*, 2021

Publication

1%

3

Marko D. Prokić, Slavica S. Borković-Mitić, Imre I. Krizmanić, Jelena J. Mutić et al. "Antioxidative responses of the tissues of two wild populations of *Pelophylax kl. esculentus* frogs to heavy metal pollution", *Ecotoxicology and Environmental Safety*, 2016

Publication

1%

4

Laura Fernández-Alacid, Ignasi Sanahuja, Borja Ordóñez-Grande, Sergio Sánchez-Nuño et al. "Skin mucus metabolites in response to physiological challenges: A valuable non-invasive method to study teleost marine species", *Science of The Total Environment*, 2018

Publication

1 %

5

Thitaporn Thitiyan, Paweena Pongdontri, Bundit Tengjaroenkul, Lamyai Neeratanaphan. " Bioaccumulation and oxidative stress in affected by heavy metals and metalloid in municipal landfill reservoir ", *International Journal of Environmental Studies*, 2021

Publication

1 %

6

Aina O. Adeogun, Oju R. Ibor, Roseline Omiwole, Azubuike V. Chukwuka, Adeola H. Adewale, Omobolanle Kumuyi, Augustine Arukwe. "Sex-differences in physiological and oxidative stress responses and heavy metals burden in the black jaw tilapia, *Sarotherodon melanocheilus* from a tropical freshwater dam (Nigeria)", *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 2020

Publication

1 %

7

Paolo Pastorino, Antonia Concetta Elia, Barbara Caldaroni, Vasco Menconi et al. "Oxidative stress ecology in brook trout (*Salvelinus fontinalis*) from a high-mountain lake (Cottian Alps)", *Science of The Total Environment*, 2020

Publication

8

Kalpana Chhaya Lakra, Tarun Kumar Banerjee, Bechan Lal. "Coal mine effluent-induced metal bioaccumulation, biochemical, oxidative stress, metallothionein, and histopathological alterations in vital tissues of the catfish, *Clarias batrachus*", *Environmental Science and Pollution Research*, 2021

Publication

9

Wenjun Tu, Hong Wang, Song Li, Qiang Liu, Hong Sha. "The Anti-Inflammatory and Anti-Oxidant Mechanisms of the Keap1/Nrf2/ARE Signaling Pathway in Chronic Diseases", *Aging and disease*, 2019

Publication

10

Jun-Hwan Kim, Ju-Chan Kang. "Effects of sub-chronic exposure to lead (Pb) and ascorbic acid in juvenile rockfish: Antioxidant responses, MT gene expression, and neurotransmitters", *Chemosphere*, 2017

Publication

1 %

1 %

1 %

<1 %

11

Fahri Karayakar, Özge Yurt, Bedii Cicik, Mustafa Canli. "Accumulation and Elimination of Cadmium by the Nile Tilapia (*Oreochromis niloticus*) in differing Temperatures and Responses of Oxidative Stress Biomarkers", *Bulletin of Environmental Contamination and Toxicology*, 2022

Publication

<1 %

12

Slavka Stankovic, Predrag Kalaba, Ana R. Stankovic. "Biota as toxic metal indicators", *Environmental Chemistry Letters*, 2013

Publication

<1 %

13

Qian Qi, Chunnuan Zhang, Shun Shi, Xiaolin Liu, Yang Liu, Lin Fu, Yong Liu, Peiyi Meng, Yiping Zhang, Pinpin Lv. " Effects of ammonia stress on liver microstructure, antioxidant capability and inflammation - related genes and post - exposure recovery in the hybrid sturgeon (♀ × ♂) ", *Aquaculture Research*, 2021

Publication

<1 %

14

Yuying Pan, Lina Tian, Qiaoling Zhao, Zhen Tao, Jinsheng Yang, Youlin Zhou, Rui Cao, Guangxu Zhang, Wenyu Wu. "Evaluation of the acute toxic effects of crude oil on intertidal mudskipper (*Boleophthalmus pectinirostris*) based on antioxidant enzyme

<1 %

activity and the integrated biomarker response", Environmental Pollution, 2021

Publication

15

Cristina Alarcon, Barton Wicksteed, Marc Prentki, Barbara E. Corkey, Christopher J. Rhodes. "Succinate Is a Preferential Metabolic Stimulus-Coupling Signal for Glucose-Induced Proinsulin Biosynthesis Translation", Diabetes, 2002

Publication

<1 %

16

Prabhat Kumar Rai. "Heavy Metal Phytoremediation from Aquatic Ecosystems with Special Reference to Macrophytes", Critical Reviews in Environmental Science and Technology, 2009

Publication

<1 %

17

Caimi, Gasco, Biasato, Malfatto, Varello, Prearo, Pastorino, Bona, Francese, Schiavone, Elia, Dörr, Gai. "Could Dietary Black Soldier Fly Meal Inclusion Affect the Liver and Intestinal Histological Traits and the Oxidative Stress Biomarkers of Siberian Sturgeon (*Acipenser baerii*) Juveniles?", Animals, 2020

Publication

<1 %

18

Davi Vieira Teixeira da Silva, Diego dos Santos Baião, Vitor Francisco Ferreira, Vania Margaret Flosi Paschoalin. "Betanin as a multipath oxidative stress and inflammation

<1 %

modulator: a beetroot pigment with protective effects on cardiovascular disease pathogenesis", *Critical Reviews in Food Science and Nutrition*, 2020

Publication

19

Casalino, E.. "Acute cadmium intoxication induces alpha-class glutathione S-transferase protein synthesis and enzyme activity in rat liver", *Toxicology*, 20060116

Publication

20

E. Chris Kazala. "Fatty acid composition of muscle fat and enzymes of storage lipid synthesis in whole muscle from beef cattle", *Lipids*, 11/2006

Publication

21

Mahmoud Mohseni, Najjar Lashgari Saltanat, Mohammad Esmail Rastravan, Younes Golalipour. " Effects of betaine supplementation in plant - protein - based diets on growth performance, haemato - immunological parameters, antioxidant status and digestive enzyme activities of juvenile Caspian trout (, Kessler, 1877) ", *Aquaculture Nutrition*, 2021

Publication

22

Irwin, Mary Elizabeth, Nilsa Rivera-Del Valle, and Joya Chandra. "Redox Control of Leukemia: From Molecular Mechanisms to

<1 %

<1 %

<1 %

<1 %

Therapeutic Opportunities", Antioxidants & Redox Signaling, 2012.

Publication

23

Lucilene Finoto Viana, Debora Cristina Damasceno de Souza, Edmilda Batista da Silva, Fábio Kummrow et al. "Bioaccumulation of metals and genotoxic effects in females of *Colomesus asellus* collected in an Amazon River estuary, Amapá, Brazil", *Limnetica*, 2023

Publication

24

Madiha Zaynab, Rashid Al-Yahyai, Ayesha Ameen, Yasir Sharif, Liaqat Ali, Mahpara Fatima, Khalid Ali Khan, Shuangfei Li. "Health and environmental effects of Heavy metals", *Journal of King Saud University - Science*, 2021

Publication

25

Yu-Feng Lin, Chih-Wen Cheng, Chung-Shiuan Shih, Jenn-Kang Hwang, Chin-Sheng Yu, Chih-Hao Lu. "MIB: Metal Ion-Binding Site Prediction and Docking Server", *Journal of Chemical Information and Modeling*, 2016

Publication

26

David U. Hernández-Becerril. "Toxic and harmful marine phytoplankton and microalgae (HABs) in Mexican Coasts", *Journal of Environmental Science and Health Part A*, 8/2007

Publication

<1 %

<1 %

<1 %

<1 %

27

Ming-Jian Liu, Hua-Yang Guo, Bo Liu, Ke-Cheng Zhu et al. "Gill oxidative damage caused by acute ammonia stress was reduced through the HIF-1 α /NF- κ B signaling pathway in golden pompano (*Trachinotus ovatus*)", *Ecotoxicology and Environmental Safety*, 2021

Publication

<1 %

28

Srinibas Das, Indrajit Kar, Amlan Kumar Patra. "Cadmium induced bioaccumulation, histopathology, gene regulation in fish and its amelioration – a review", *Journal of Trace Elements in Medicine and Biology*, 2023

Publication

<1 %

29

Anna Carolina Resende, Diego Mauro Carneiro Pereira, Ieda Cristina Schleger, Maria Rosa Dmengenon Pedreiro de Souza et al. "Effects of heat shock on energy metabolism and antioxidant defense in a tropical fish species", *Journal of Fish Biology*, 2022

Publication

<1 %

30

Bee Ooi, Bey Goh, Wei Yap. "Oxidative Stress in Cardiovascular Diseases: Involvement of Nrf2 Antioxidant Redox Signaling in Macrophage Foam Cells Formation", *International Journal of Molecular Sciences*, 2017

Publication

<1 %

- | | | |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 31 | Chen, F.. "Toxicity assessment of simulated urban runoff containing polycyclic musks and cadmium in <i>Carassius auratus</i> using oxidative stress biomarkers", <i>Environmental Pollution</i> , 201203
Publication | <1 % |
| 32 | Luísa A. Barreira. "Oxidative stress in the clam <i>Ruditapes decussatus</i> (Linnaeus, 1758) in relation to polycyclic aromatic hydrocarbon body burden", <i>Environmental Toxicology</i> , 04/2007
Publication | <1 % |
| 33 | Peter F. Surai. "Vitagenes in avian biology and poultry health", Wageningen Academic Publishers, 2020
Publication | <1 % |
| 34 | Zheng-Yu Jiang, Meng-Chen Lu, Qi-Dong You. "Nuclear Factor Erythroid 2-Related Factor 2 (Nrf2) Inhibition: An Emerging Strategy in Cancer Therapy", <i>Journal of Medicinal Chemistry</i> , 2018
Publication | <1 % |
| 35 | "Tropical Peatland Ecosystems", Springer Nature, 2016
Publication | <1 % |
| 36 | A. Eroglu, Z. Dogan, E. G. Kanak, G. Atli, M. Canli. "Effects of heavy metals (Cd, Cu, Cr, Pb, | <1 % |

Zn) on fish glutathione metabolism",
Environmental Science and Pollution
Research, 2014

Publication

37

Ashish Kumar Srivastava, Poonam Bhargava,
Lal Chand Rai. "Salinity and Copper-Induced
Oxidative Damage and Changes in the
Antioxidative Defence Systems of *Anabaena
doliolum*", World Journal of Microbiology and
Biotechnology, 2005

Publication

<1 %

38

Gideon Aina Idowu. "Heavy metals research in
Nigeria: a review of studies and prioritization
of research needs", Environmental Science
and Pollution Research, 2022

Publication

<1 %

39

Jochen A. Müller, Bernhard Schink. "Initial
steps in the fermentation of 3-
hydroxybenzoate by *Sporotomaculum
hydroxybenzoicum*", Archives of Microbiology,
2000

Publication

<1 %

40

Laura Fernández-Alacid, Ignasi Sanahuja,
Borja Ordóñez-Grande, Sergio Sánchez-Nuño,
Marcelino Herrera, Antoni Ibarz. "Comparison
between properties of dorsal and ventral skin
mucus in Senegalese sole: Response to an
acute stress", Aquaculture, 2019

Publication

<1 %

41

Lin Han, Qing Yang, Wenfang Ma, Jia Li, Liuzhu Qu, Min Wang. "Protocatechuic Acid Ameliorated Palmitic-Acid-Induced Oxidative Damage in Endothelial Cells through Activating Endogenous Antioxidant Enzymes via an Adenosine-Monophosphate-Activated-Protein-Kinase-Dependent Pathway", *Journal of Agricultural and Food Chemistry*, 2018

Publication

<1 %

42

Natalia Kurhalyuk, Halyna Tkachenko, Katarzyna Pałczyńska. "Lipid peroxidation and antioxidant defense system in spawn of brown trout (*Salmo trutta m. trutta* L.) affected by ulcerative dermal necrosis", *Archives of Polish Fisheries*, 2010

Publication

<1 %

43

Orhan Bahar, Gökhan Eraslan. "Investigation of the efficacy of diosmin against organ damage caused by bendiocarb in male Wistar albino rats", *Environmental Science and Pollution Research*, 2023

Publication

<1 %

44

T.F. Amaral, F.H. Miyasaki, E.S. Braga, J.S. Azevedo. "Temporal and spatial toxicogenetic damage in estuarine catfish *Cathorops spixii* from a marine protected area with evidence of anthropogenic influences", *Science of The Total Environment*, 2021

<1 %

45 Xiao-Qiang Gao, Fan Fei, Huan Huan Huo, Bin Huang, Xue Song Meng, Tao Zhang, Bao-Liang Liu. "Effect of acute exposure to nitrite on physiological parameters, oxidative stress, and apoptosis in *Takifugu rubripes*", *Ecotoxicology and Environmental Safety*, 2020

Publication

46 Giuseppe Sciortino, Eugenio Garribba, Jaime Rodríguez-Guerra Pedregal, Jean-Didier Maréchal. "Simple Coordination Geometry Descriptors Allow to Accurately Predict Metal-Binding Sites in Proteins", *ACS Omega*, 2019

Publication

47 Margaret G. Mills, Richard Ramsden, Eva Y. Ma, Jone Corrales et al. "CRISPR-Generated *Nrf2a* Loss- and Gain-of-Function Mutants Facilitate Mechanistic Analysis of Chemical Oxidative Stress-Mediated Toxicity in Zebrafish", *Chemical Research in Toxicology*, 2019

Publication

48 Matos, P.. "Biochemical and histological hepatic changes of Nile tilapia *Oreochromis niloticus* exposed to carbaryl", *Pesticide Biochemistry and Physiology*, 200709

Publication

49

Cheng Chen, Yankui Tang, Yuyang Liu, Yuanzhang Liang, Kaixuan Zhang, Shengye Wang, Yan Liang. "Effect of competitive adsorption on zinc removal from aqueous solution and zinc smelting effluent by eucalyptus leaf-based magnetic biosorbent", *Journal of Environmental Science and Health, Part A*, 2017

Publication

<1 %

50

Lozenka Ivanova, Katerina Rebok, Maja Jordanova, Zrinka Dragun et al. " The effect of different pollutants exposure on the pigment content of pigmented macrophage aggregates in the spleen of Vardar chub (Karaman, 1928) ", *Microscopy Research and Technique*, 2020

Publication

<1 %

51

Mostafa F. Abdelzaher et al.. "Effects of Lead and Cadmium Accumulation on Survival and Growth of the Nile Tilapia (*Oreochromis niloticus*)", *Egyptian Journal of Aquatic Biology and Fisheries*, 2022

Publication

<1 %

52

Natalija Topić Popović, Lara Čižmek, Sanja Babić, Ivančica Strunjak-Perović, Rozelindra Čož-Rakovac. "Fish liver damage related to the wastewater treatment plant effluents",

<1 %

Environmental Science and Pollution Research, 2023

Publication

Exclude quotes Off

Exclude matches < 5 words

Exclude bibliography On