

Oodev Injection Frequency and Time Period in Advancing Gonad Rematuration of Snakehead (*Channa striata* Blkr) in Hapa System *by Herliwati .*

Submission date: 28-May-2023 06:08PM (UTC+0700)

Submission ID: 2103576427

File name: Oodev_Injection_Frequency_and_Time_Period.pdf (204.75K)

Word count: 7152

Character count: 37443

Oodev Injection Frequency and Time Period in Advancing Gonad Rematuration of Snakehead (*Channa striata* Blkr) in Hapa System

Kaspul Anwar¹, Untung Bijaksana², Herliwati², Ahmadi^{2*}

¹Postgraduate Program of Fishery Science, ²Faculty of Marine and Fisheries
Universitas Lambung Mangkurat, Banjarbaru 70714 Indonesia

*Corresponding author: ahmadi@unlam.ac.id

Abstract—This study provides meaningfully scientific information on the attempt to accelerate the gonad rematuration of snakehead (*Channa striata*) to support the quality seed production for both commercial and restocking purposes in South Kalimantan, Indonesia. A total of 48 snakehead fish broods (296-312 mm total length and 234-298 g weight) were subjected to different treatment levels using oodev dose of 0.5 ml kg⁻¹ fish weight to determine the best frequency and time period for gonad rematuration after spawning. Treatment-A: 3 times/9 days, B: 2 times/6 days, C: 1 time/3 days, and D: no oodev injection (control). The 12 hapas were used comprising 4 individuals per hapa (0.4×0.4×1 m) under the controlled condition. After all, ten snakehead brood samples were dissected to find out the gonad maturity level of fish. The treatment-A showed the best performance in term of mean fecundity (19600±450.00 granules/individuals), egg diameter (1.10±0.15 mm), and gonad somatic index/GSI (3.41±0.90 %) among other treatments. Dealing with hepatosomatic index (HSI), the treatment-C was significantly higher than treatment-A and D, but not differ from treatment-B. The mean HSI was varied from 0.89±0.03 % to 1.53±0.23 %. Temperature, pH and DO during sampling period are considered comfortable for snakehead fish broods.

Keywords— *Channa striata*, gonad rematuration, fecundity, egg diameter, GSI, HSI.

INTRODUCTION

In the world, the snakeheads comprise two extant genera, namely *Channa* (36 species native to Asia) and *Parachanna* (four African species), and most recently, barcoding snakehead is revisited to solve perpetuated taxonomic confusions (Conte-Grand *et al.*, 2017). Snakehead (*Channa striata* Blkr) of family Channidae, is considered as valuable food fish not only in Indonesia (Widodo *et al.*, 2013; Irhamsyah *et al.*, 2017), but also other countries such as Thailand (Khomsab and Wannasri, 2017), Philippines (Jumawan and Seronay, 2017), Vietnam

(Quyen *et al.*, 2016), Malaysia (Song *et al.*, 2013), Cambodia (Sinh, 2014), Sri Lanka (Wijeyaratne, 1994), Nigeria (Ama-Abasi and Ogar, 2013), Bangladesh (Islam *et al.*, 2013), India (Kashyap *et al.*, 2014), Pakistan (Najero *et al.*, 2015) and China (Guet *et al.*, 2015) due to delicious, high-quality meat fish and availability throughout the year. Snakehead can be commercially cultured in fish farming, earthen ponds, or hapa system (Kumar *et al.*, 2011; Quyen *et al.*, 2016), and culture strategies of them are currently being developed (Xie *et al.*, 2002; Xie *et al.*, 2017; He *et al.*, 2015; Istiyanto and Diana, 2016). They inhabit freshwater such as swamps, rivers, streams, lakes, reservoirs, irrigation canals and paddy fields (Saikia *et al.*, 2012; Muthmainnah, 2013; Kashyap *et al.*, 2014; Saare, 2015; Singh and Serajuddin, 2017). Snakehead locally known as “Haruan”, is able to tolerate to adverse environments due to its hardiness and air-breathing capabilities assisted with a suprabranchial chamber, an air-breathing organ (Chandra and Banerjee, 2004). They are carnivorous feeders that consume plankton, aquatic insect, mollusks, fish or frogs. Snakeheads from river or swamps are caught by using hooks, gillnet, castnet or fish pot (Song *et al.*, 2013; Irhamsyah *et al.*, 2017). Overfishing, pollution, habitat disruption, disease, and growing human intervention on wetlands is very likely threat to this species (Balkhis *et al.*, 2011; Uthayakumar *et al.*, 2014; Rao *et al.*, 2015).

Seasonal reproduction is a strategy adapted by fishes to guarantee better chances of their offspring’s survival by timing reproductive activity with favorable conditions during certain times of the year (Bernal *et al.*, 2015). An understanding of the reproductive biology of fishes in relation to external factors (e.g. photoperiod, temperature, rainfall, salinity, and food supply) in their habitat is useful in the propagation of captive fish broodstock (Bromage *et al.*, 2001). Snakehead in swamp waters spawning in the early or mid-rainy season. It takes longer time for the gonad growth and reproductive process, meanwhile the culture development is highly depend on the availability of

fish seeds that meet the timeliness, quality and quantity. Seeds can be continuously produced if it is supported by the availability of matured broodstock with good egg quality. One of efforts to meet the need of seeds production is by increasing the spawning frequency of Nilem fish broodstock through the acceleration of gonad rematuration period after spawning. The gonad rematuration period usually takes longer time compared to the maturation process. It is clearly found in *Osteochilus hasselti* in which the gonad maturation period takes 10 days (Cholifah, 2016), while its rematuration period takes 17 days (Fitriatin *et al.*, 2018) by using oodev hormone 1 ml kg⁻¹.

Understanding the proper function of follicle stimulating hormone (FSH) of oodev content will reduce the artificial spawning failure of the snakehead, because it is directly involved in the acceleration of the vitelogenesis process (Moore and Ward, 1980). The previous studies confirmed that oodev hormone beneficially supports for gonad maturation and spawning in *Pangasius hypophthalmus* (Ernawati, 1999), *Clarias batrachus* (Zairin *et al.*, 2000), *Hemibagrus nemurus* (Supriyadi, 2005), and *Osteochilus hasselti* (Cholifah, 2016), as well as for gonad rematuration in *P. hypophthalmus* (Agustinus, 2013), *Anabas testudineus* (Sari, 2015), and *O. hasselti* (Fitriatin *et al.*, 2018), but lack of available information on most snakehead species. Therefore, the present study is performed to determine the best frequency and time period (day) for advancing the snakehead gonad maturation through oodev hormone intervention. From all stages of this research, we endeavor to provide a simple technology package of snakehead gonad rematuration to supply the needs of quality seeds for both commercial and restocking purposes.

II. MATERIALS AND METHODS

Study site and experimental condition

The research was carried out in the Wet Laboratory at the Faculty of Marine and Fisheries, University Lambung Mangkurat for three months. A set of tools, equipment and materials are well-prepared as described in Table 1. About one hundred snakehead broodstocks were collected from local fishers in the monotonous swamp and were kept in the Wet Laboratory. The only forty-eight fish samples (296-312 mm total length and 234-298 g weight) were used in this study, while the remaining fish was kept in a large hapa (1.5×1.5×1.0 m) as stock in case of the death during the trial periods. The hapas were placed in a concrete tank pond (7×4×1.2 m). Tap water was precipitated one week before treatment. Water lettuce (*Pistia stratiotes* L) was given for shelter and supplemental nutrients in the water. Fish were fed with the frog as 4% of body weight per day with the frequency of once a

day. The twelve hapas were used in this experiment with density level of 4 individuals per hapa (0.4×0.3×1 m). Water quality parameters recorded during the study were as follows: temperatures from 26.40 to 27.45 °C, pH from 6.90 to 7.23, DO from 4.59 to 6.10 mg l⁻¹ and NH₃ from 0.15 to 0.23 mg l⁻¹. Temperature, pH and DO were measured using Watercheckker U10 Horiba, while NH₃ was determined by Spectofotometer with Spec-Nessler method.

Experimental Procedures

The experimental design used was Completely Randomized Design with four different treatments (A, B, C, and D) and each treatment was repeated three times. For treatment-A, B and C, the Oodev dose was given to the snakehead broodstocks with the same amount of 0.5 ml kg⁻¹ fish weight but vary with frequency and time period (day). For instance, the treatment-A: 3 times/9 days, B: 2 times/6 days, C: 1 time/3 days, and D: no oodev injection (control) and later to be compared to other treatments. The details of experimental treatment are given in Table 2. At the end of experiments, a total of ten snakehead samples were dissected to find out the gonad maturity level of fish, and reproductive indicators such as egg diameter, fecundity, Somatic Gonad Index (SGI) and Hepato Somatic Index (HSI) were measured. The egg diameter was measured using a micrometer. The fecundity, SGI and HSI were calculated using the following formulas:

$$F = n \times G / g \quad (1)$$

F is fecundity; n is the average number of eggs in sub-sample; G is gonad weight (g); and g is the sub-sample weight (g).

$$GSI = Gw / Bw \times 100\% \quad (2)$$

GSI is Gonado Somatic Index; Gw is Gonad weight (g); and Bw is Body weight of fish (g)

$$HSI = Lw / Bw \times 100\% \quad (3)$$

HSI is Hepato Somatic Index; Lw is Liver weight (g); and Bw is Body weight of fish (g)

Statistical Analysis

At the beginning, the normality and homogeneity of experimental data obtained were analysed using Lilliefors test and Bartlett test, respectively. Data transformation should be first done if data were found not normal or not homogeneous. If the assumption was fulfilled, then apply for the Analysis of Variants (ANOVA). The Mean differences test was used if there were significantly

differences among the treatments. All tests were analysed at the 0.05 level of significance using SPSS-16 software.

III. RESULTS AND DISCUSSION

The mean \pm standard deviation of fecundity, egg diameter, somatic gonad index (SGI) and hepatosomatic index (HSI) obtained from experiments was presented in Table 3. It is clearly shown that among different injection frequency, the treatment-A was determined as the best performance in term of generating fecundity, egg diameter and SGI rate i.e. 19600 ± 450.00 granules/individuals, 1.10 ± 0.15 mm and 3.41 ± 0.90 % respectively. It was followed by treatment-B with the respective values of 7300 ± 556.77 granules/individuals, 0.90 ± 0.11 mm and 1.39 ± 0.14 %. Meanwhile, the treatment-C showed better performance as compared to treatment-D (the control) in term of fecundity that is 1367 ± 2367.13 granules/individuals. With regard to the egg production, the mean fecundity obtained ranging from 1033 ± 1789.79 to 19600 ± 450.00 granules/individuals. There were significant differences in the mean fecundity among the treatments ($p < 0.05$). The mean fecundity of treatment-A was considerably higher than other treatments ($p < 0.01$). The treatment-B was significantly higher than treatment-C ($p < 0.05$); whereas treatment-C and D was not significant different ($p > 0.05$). There were significant differences in mean egg diameter among the treatments ($p < 0.05$). The mean egg diameter was ranged from 0.20 ± 0.40 mm to 1.10 ± 0.15 mm. The maximum size of egg diameter for injected fish in the treatment-A or treatment-B was found larger than treatment-D without injection in the control group ($p < 0.05$), indicating that Oodev had positive effect on the egg maturation of snakehead broodstock. No significant difference was observed in the egg diameter between treatment-C and D ($p > 0.05$).

Overall, the result also clearly demonstrated that the treatment-A yielded the highest GSI rate among other treatments ($p < 0.01$). The treatment-B was considerably higher than treatment-D ($p < 0.05$), but not significant different from treatment C ($p > 0.05$). There was no significant difference in the SGI rate between treatment-C and D ($p > 0.05$). The mean SGI rate was ranged from 0.40 ± 0.62 % to 3.41 ± 0.90 mm. The GSI rate gradually increases with increased frequency and time given. The other way the HSI rate decreases with increased frequency and time. The treatment-A showed higher HSI rate as compared to treatment-A and D ($p < 0.05$), but not differ from treatment-B ($p > 0.05$) was observed. No statistical difference in the mean HSI rate was found between treatment-A, B and D ($p > 0.05$). The mean HSI rate was varied from 0.89 ± 0.03 % to 1.53 ± 0.23 %.

Table 4 shows the mean \pm standard deviation of the three estimated weights i.e. body weight, liver weight and gonad

weight, as well as the ratios of these parameters for each treatment. The body weight and the liver weight were ranged from 253.67 ± 21.33 to 266.80 ± 7.26 g and from 2.43 ± 0.61 to 3.68 ± 0.45 g respectively. The ratio of liver weight to body weight obtained from the treatment-C was considerably higher than that of treatment-A and D ($p < 0.05$), but not differ from treatment-B ($p > 0.05$) was detected. There were no statistically significant difference in those ratios between treatment-A, B and D ($p > 0.05$). The ratio of liver weight to body weight was varied from 0.0091 to 0.0143. The results also clearly revealed that the treatment-A yielded the highest gonad weight (9.20 ± 2.50 g) among other treatments ($p < 0.05$). The ratio of gonad weight to body weight was ranged from 0.0050 to 0.0345. Furthermore, the highest ratio of fecundity to body weight was 73.44 (treatment-A), while the lowest ratio was 3.88 (treatment-D). It was similarly demonstrated, the highest ratio of fecundity to body length was 62.88 (treatment-A), while the lowest ratio was 3.32 (treatment-D). Such relationships between fecundity and body weight as well as between fecundity and body length can be seen in Figure 1.

The appropriate hormone preparation should be selected on the basis of the species to be spawned and the availability of the hormones. Many factors which have impact on ability of induced spawning, include: 1) condition of the fish, 2) stage of sexual maturity, 3) size of the fish, 4) previous spawning history, 5) water temperature, 6) season of the year and 7) dosage of hormone use (Rottmann *et al.*, 1991). Hormones for induced spawning have been widely used in air-breathing fishes such as human chorionic gonadotropin (HCG) (Mollah and Tan, 1983; Inyang and Hettiarachchi, 1994), luteinizing hormone releasing hormone analogue (LHRHa) (Fermin, 1992), and ovaprim (Alok *et al.*, 1993; Haniffa *et al.*, 1996). Oodev is one solution when fish can not perform vitellogenesis and spermatogenesis.

The oodev hormonal treatments have been successfully used for advancing the process of gonad rematuration period in *A. testudineus* for 12-24 days (Sari, 2015). Furthermore, Agustinus (2016) reported that *P. hypophthalmus* being injected with oodev 0.7 ml kg^{-1} fish weight every 7 days for 4 weeks showed the best performance in term of growth rate and egg diameter and also can accelerate gonad rematuration of them for 28 days after spawning. Fitriatin *et al.* (2018) affirmed that oodev hormone dose of 1 ml kg^{-1} fish weight was effective for gonad rematuration of *O. hasselti* broodstock within 17 days with the GSI rate of 14.48 %, average egg diameter of 1.035 mm and body weight of 31.25 mm. For the time being, the effectiveness of oodev works in snakehead is still questionable due to lack of available evidence. Dealing with the presence of oodev itself, it is commonly

known that oodev hormone is a combination of Pregnant Mare's Serum Gonadotropin (PMSG) hormone and Antidopamine (AD) compounds. PMSG hormone contains follicle stimulating hormone (FSH) and luteinizing hormone (LH) (Moore and Ward, 1980). The role of FSH increases the chances of egg cell maturation, while the role of LH stimulates ovulation. PMSG chemical structure is similar to the FSH and LH, whereas FSH structure is greater than LH (Reeves, 1987). The PMSG hormone in oodev is thought to have triggered the gonad maturation of brood fish, where vitelogenesis only takes place when a hormone is present associated with this process. The gonadotropin hormone of PMSG injected from the outside will work on the egg by maturing the existing follicle by stimulating the growth of interstitial cells and the formation of luteal cells. The process of gonadal rematuration starts from the synthesis of vitelogenin which is the precursor of egg yolks (Wiegand, 1982). One of the conditions governing vitelogenesis is the availability of hormones associated with vitelogenesis in the body. It is known that there is a positive correlation between gonadal steroid hormone and vitelogenesis (Mackenzie *et al.* 1998). It was clear from our findings that snakehead successively induced with 0.5 ml kg⁻¹ fish weight for three-frequency and time periods in the treatment-A produced more eggs about 3 times higher than twice, or about 14 times higher than once or about 19 times higher than with no injection. It was significantly different from other treatments ($p < 0.05$). The fecundity of fish varied according to the size (length-weight), age, species, food availability and season. Larger fish tends to have greater fecundity than small fish. For a given size, females in better condition exhibit higher fecundity (Kjesbu *et al.*, 1991). The relationship between the age of brood female and the size of the egg is the square. The young mother spawning for the first time produces small eggs (about 0.8 mm diameter); the productive females will produce large eggs (about 1 mm) and females again producing small eggs (about 0.8 mm). A relationship between length, weight, and fecundity for the family Clariidae was well-documented in *C. gachua* (Gaikwad *et al.*, 2009; Widodo *et al.*, 2013), *C. striatus* (Islam *et al.*, 2013; Sakhare, 2015), and *C. Limbata* (Kamsab and Wannasri, 2017). Figure 1 clearly shows that the ratio of fecundity to body weight as well as the ratio of fecundity to body length increases proportionally to the time periods of oodev injections given. The longer time period of oodev injections, the higher the ratio of fecundity to body sizes. It also implies that the larger fish the higher ratio of fecundity to body sizes. Such relationships were expressed in the following equations i.e. $y = 3.578 x^{2.0343}$ and $y = 3.064 x^{2.0343}$ with the coefficient of determination (R^2) was 0.9748, suggesting that approximately 97% of the variation in the fecundity

data could be explained by the body weight of the fish. In practical, when researchers endeavor to evaluate the commercial potential of fish stocks, the fecundity estimation must consider a variety of attributes, including size at first sexual maturity, duration of spawning season, daily spawning behavior and spawning fraction (El-Drawany, 2013).

Snakehead fish brood injected with oodev 0.5 ml kg⁻¹ fish weight in the treatment-A showing the biggest egg diameter and was significantly different from other treatments ($p < 0.05$). However, the average egg diameter of *C. striata* in the present study (1.10 mm) was smaller than *C. striatus* (1.53 mm) in Malaysia and India (Ghaedi *et al.*, 2013; Sakhare, 2015.) or *Ophiocephalus striatus* (1.60 mm) in Mekong River Vietnam (Long *et al.*, 2002). The vitelogenesis process can be seen from the value of the enlarged egg diameter. It is the process of vitelogenin synthesis in the liver which is the main precursor of yolk, resulted in the egg diameter increases. The diameter of the eggs increases as the weight of the gonads increase. The sexual maturity in fish is characterized by the development of egg diameter and through the distribution of its egg size (Kuo *et al.*, 1974). The egg size can be expressed in many ways. Single diameter is commonly used, but the longest diameter (i.e. egg length and egg width) is also occasionally used. Other egg sizes include the volume of eggs, wet weight and dry weight. From an energetic standpoint, the best term for egg size is the equality of egg calorie (energy content per egg or joule per egg), because it shows the energy available to the developing embryo (Ginzburg, 1972). The large egg size is a guarantee of higher survival (Kamler, 1992), this because the egg contains food reserve to be used by fish larvae for survives. Larvae derived from the large egg will have more egg yolk reserves as energy source before getting food from outside. The size of the egg diameter can determine the quality of the yolk content. The egg diameter is the accumulation of the vitelogenesis process that is the absorption of vitelogenin which is going to prospective yolk. Increasing the yolk granules in the number and size resulted in the oocyte volume will be greater until the maximum size then the egg is in "dormant" phase. Egg quality is affected by both internal factors (e.g. brood age, brood size and genetic) and external factors (e.g. feed, temperature, light, density, and population). The first spawning female fish produce small diameter eggs. The egg diameter increases clearly when the second spawning and the rate of increase are slower in subsequent spawning.

The results also confirmed that snakehead consecutively injected with oodev 0.5 ml kg⁻¹ in the treatment-A increasing the GSI rate about 2.5 times higher than treatment-B, or about 8.5 times higher than treatment-C or

about 7 times higher than with no injection in treatment-D. It is clearly revealed that the GSI increases proportionally towards the time periods of oodev interventions. Also, an increase of GSI value is in conjunction with the increased doses of Oodev hormone given (Fitriatin *et al.*, 2018). It is alleged that FSH and LH activity in oodev hormone influences the gonad development as a whole (Moore and Ward, 1980). The GSI of family Channidae varies according to the type of species (Gaikwad *et al.*, 2009; Kapil *et al.*, 2011; Widodo *et al.*, 2013; Tiwari *et al.*, 2014). For example, the GSI of *C. striata* (3.41 %) in the present study was lower than GSI of *C. gachua* (6.00 %) in Godavari River, India (Gaikwad *et al.*, 2009) or *C. marulius* (47.56 %) in Son River Shahdol, India (Tiwari *et al.*, 2014), or *C. striatus* (8.00 %) in Badin Sindh District, Pakistan (Narejo *et al.*, 2015). The IGS value is used to predict when the fish will be ready for spawning. Increased IGS values can be triggered by oocytes, while vitelogenin is main precursor of yolk which is the main component of the growing oocyte. When the process of vitelogenesis is continuing, the yolk granules increase in number and size, so that the volume of the oocyte enlarges and eventually will lead to increased value of GSI. The process of vitelogenin formation starts from the presence of environmental factor cues such as fotoperiode, temperature, eating activity, and other factors that will all stimulate the hypothalamus to secrete gonadotropin releasing hormone (GnRH). GnRH will be secreted into the blood stream will stimulate hypophysis to secrete gonadotropin in hormones (Darwisito *et al.*, 2006). In the reproductive cycle, GSI increases with the maturation process, whereas the HSI in contradicting to GSI (Lodeiros *et al.*, 2001). It was clear from our findings that the treatment-A produced the highest GSI among other treatments (3.41 %), contrariwise it provided the lowest HSI among other treatments (0.89 %) in the same amount of oodev content. In other words, the longer time period of oodev injection given, the lower the ratio of liver weight to body weight gained. This is attributable to the use of energy reserves derived from the liver instead of energy sourced from the body. The HIS of *C. striata* (1.53 %) in the present study was lower than HSI of *C. punctatus* (1.64 %) in Sylhet, Bangladesh (Hossain 2013), but slightly higher than *C. striatus* (1.40 %) in Penang, Malaysia (Ghaedi *et al.*, 2013). In addition, Bijaksana (2006) argued that most of the snakehead caught during fishing season is allocating their somatic growth for reproductive growth. In line with this; temperature, pH and DO during sampling period are considered to be comfortable for snakehead fish broods.

CONCLUSION

Oodev hormone had a significant effect in speeding up the process of gonad rematuration in the snakehead injected. Oodev dose of 0.5 ml kg⁻¹ those given in 3 times/9 days (treatment-A) is the most effective treatment to rematuration the gonad of snakehead female broodstock within 30 days, producing fecundity of 19,600 granules/individuals, egg diameter of 1.10 mm, and GSI rate of 3.41 %. The results may applicable when outdoor environmental quality is improved. Despite these findings, the outdoor environmental quality should be considered into account. Some of the information presented here may be applicable in other geographical areas.

ACKNOWLEDGMENTS

This research was under our own means of funding. The authors are grateful to the Head of Wet Laboratory, Faculty of Marine and Fisheries, Lambung Mangkurat University for facilitating this study. The authors also thank the reviewers for significantly improving the quality of paper to a publishable level.

REFERENCES

- [1] Agustinus. (2016). Kinerja reproduksi dengan induksi oodev dalam vitelogenesis pada rematuration induk ikan patin (*Pangasius hypophthalmus*) di dalam wadah budidaya. Fish Scientiae, 3(5): 1-16
- [2] Ama-abasi, D. and Agar, A. (2013). Proximate analysis of snakehead fish, *Parachanna obscura*, (Gunther 1861) of the cross river, Nigeria. Journal of Fisheries and Aquatic Sciences 8(1): 295-298. <https://doi.org/10.3923/jfas.2013.295.298>
- [3] Alok, D., Krishnan, T., Talwar, G.P. and Garg, L.C. (1993). Induced spawning of cat fish *Heteropneustes fossilis* (Bloch), using D-Lys super (6) salmon gonadotropin releasing hormone analog. Aquaculture 115: 159-167
- [4] Balkhis, A.B.S., Jamsari, A.F.J., Hwai, T.S., Yasin, Z. and Azizah, M.N.S. (2011). Evidence of geographical structuring in the Malaysian snakehead, *Channa striata* based on partial segment of the CO1 Gene. Journal of Genetics and Molecular Biology 34(3): 520-523.
- [5] Bernal, R.A.D., Aya, F.A., Garcia, L.M.B. and De Jesus-Ayson, E.G.T. (2015). Seasonal gonad cycle of the climbing perch *Anabas testudineus* (Teleostei: Anabantidae) in a tropical wetland. Ichthyology Research 62: 389-395. <https://doi.org/10.1007/s10228-014-0454-3>
- [6] Bijaksana, U. (2006). Preliminary study of bio-eco reproduction snakehead in Rawa Bangkai in South Kalimantan Province. National Symposium on Biotechnology in Aquaculture, 5 July 2016.

- Department of Aquaculture, Faculty of Fisheries and Marine Sciences, Bogor Agricultural University and Research Institute of Fisheries Freshwater Aquaculture Research Institute of Marine and Fisheries, Bogor.
- [7] Bromage, N., Porter, M. and Randall, C. (2001). The environmental regulation of maturation in farmed fish with special reference to the role of photoperiod and melatonin. *Aquaculture* 197: 63-98
- [8] Chandra, S. and Banerjee, T.K. (2004). Histopathological analysis of the respiratory organs of *Channa striata* subjected to air exposure. *Veterinarski Arhiv* 74(1): 37-52.
- [9] Cholifah, E.D. (2016). Pengaruh induksi hormon oocyte developer (Oodev) terhadap kematangan gonad calon induk ikan nilam (*Osteochilus hasselti*). Skripsi. Fakultas Perikanan dan Kelautan, Universitas Airlangga. Surabaya. 91 hal
- [10] Conte-Grand, C., Britz, R., Dahanukar, N., Raghavan, R., Pethiyagoda, R., Tan, H.H. et al. (2017). Barcoding snakeheads (Teleostei, Channidae) revisited: Discovering greater species diversity and resolving perpetuated taxonomic confusions. *PLoS ONE* 12(9): e0184017. <https://doi.org/10.1371/journal.pone.0184017>
- [11] Darwisito, S., Junior, M.Z., Syafei, D.S., Manaludan, W. and Sudrajat, A.D. (2006). Kajian performans reproduksi perbaikan pada kualitas telur dan larva ikan Nila (*Oreochromis niloticus*) yang diberi vitamin E dan minyak ikan berbeda dalam pakan. *Prosiding Seminar Nasional Ikan IV*. Jatiluhur, 29-30 Agustus 2006.
- [12] El-Drawany, M.A. (2013). Some biological aspects of the Por's goatfish, (Family: Mullidae) from Tripoli Coast of Libya. *Egyptian Journal of Aquatic Research* 39: 261-266. <http://dx.doi.org/10.1016/j.ejar.2013.11.003>
- [13] Ernawati, Y. (1999). Efisiensi Implantasi Analog LHRH dan 17 α -Metiltestosteron serta Pembekuan Semen dalam upaya peningkatan produksi benih ikan Jambal Siam, *Pangasius hypophthalmus*. [Disertasi]. Program Pascasarjana. Institut Pertanian Bogor, Bogor.
- [14] Fermin, J.D.T. (1992). Induction of oocyte maturation and ovulation in the freshwater Asian catfish, *Clarius macrocephalus* by LHRHa and pimozide. *Journal of Applied Ichthyology* 8: 90-98
- [15] Fitriliyani, I. (2005). Pembesaran larva ikan gabus, *Channa striata* dan efektifitas induksi hormon gonadotropin untuk pemijahan induk. [Tesis]. Program Studi Biologi Reproduksi. Sekolah Pascasarjana. IPB, Bogor.
- [16] Gaikwad, M.V., More, V.R., Shingare, S.M., Hiwarale, D.K. and Khillare, Y.K. (2009). Study on gonadosomatic and fecundity relationship in air-breathing fish *Channa gachua* (Ham.) from Godavari near Aurangabad. *African Journal of Basic and Applied Science* 1(5-6): 93-95.
- [17] Ghaedi, A., Kabir, M.A. and Hashim, R. (2013). Oocyte development and fecundity of snakehead murrel, *Channa striatus* (Bloch 1793) in captivity. *Asian Fisheries Science* 26 (2013): 39-51
- [18] Gu, H., Yu, Z., Wang, G., Ju, Q., Yang, C. and Fan, C. (2015). Impact of climate change on hydrological extremes in the Yangtze River basin, China. *Stochastic Environmental Research and Risk Assessment* 29: 693-707. <http://dx.doi.org/10.1007/s00477-014-0957-5>
- [19] Haniffa, M.A., Shaik, M.J. and Merlin, R. (1996). Induction of ovulation in *Channa striatus*. *Fishing Chimes* 16: 23-24.
- [20] Hossain, M.A. (2013). Ovarian development of spotted snakehead *Channa punctatus*, Bloch, 1853 found in natural waters of Sylhet. Thesis. Faculty of Fisheries, Sylhet Agricultural University. 51 p. <http://dx.doi.org/10.13140/RG.2.2.21339.98083>
- [21] Irahmsyah, Ahmadi, and Rusmilyansari. (2017). Fish and fishing gears of the Bangkai swamp, Indonesia. *Journal of Fisheries* 5(2): 489-496. <http://dx.doi.org/10.17017/jfish.v5i2.2017.223>
- [22] Islam, S.S., Shah, S.M., Akter, R., Biswas, P., Sabbir, W. and Bir, J. (2013). Some aspect of biology of snake head *Channa striatus*. *Khulna University Studies* 12(1&2): 59-66.
- [23] Istiyanto, S. and Diana, R. (2016). Technology engineering of aquaculture snakeheads [*Channa striatus* (Bloch, 1793)] using cross breeding from different waters for determining the genetic variation of superior seeds. *Aquatic Procedia* 7: 136-145. <http://dx.doi.org/10.1016/j.aqpro.2016.07.019>
- [24] Inyang, N.M. and Hettiarachchi, M. (1994). Efficacy of human chorionic gonadotropin (HCG) and crude extract of fish and frog in oocyte maturation and ovulations in African catfish *Clarias gariepinus* Burchell. *Aquaculture and Fisheries Management* 25: 245-258
- [25] Jumawan, J.C. and Seronay, R.A. (2017). Length-weight relationships of fishes in eight floodplain lakes of Agusan marsh, Philippines. *Philippine Journal of Science* 146(1): 95-99.
- [26] Kapil, S., Kulkarni, K.M., Gijare, S.S., Tantarapale, V.T. (2011). Seasonal change of gonadosomatic index observed in the freshwater fish *Channa punctatus*. *The Bioscan* 6(4): 571-573.

- [27] Kashyap, A., Awasthi, M. and Serajuddin, M. (2014). Length-weight and length-length relationship of freshwater murrel, *Channa punctatus* (Bloch, 1793) sampled from river Gomti in Lucknow region (Uttar Pradesh). World journal of Fisheries and Marine Sciences 6(4): 336-339. <http://dx.doi.org/10.5829/idosi.wjfm.2014.06.04.84293>
- [28] Khomsab, K. and Wannasri, S. (2017). Biological aspects of *Channa limbata* (Cuvier, 1831) in Ta Bo - Huai Yai wildlife sanctuary, Phetchabun Province, Thailand. Sains Malaysiana 46(6): 851-858. <http://dx.doi.org/10.17576/jsm-2017-4606-03>
- [29] Kjesbu, O.S., Klungsoyr, J., Kryvi, H., Witthames, P.R. and Walker M.G. (1991). Fecundity, artesia, and egg size of captive Atlantic cod (*Gadus morhua*) in relation to proximate body composition. Canadian Journal of Fisheries and Aquatic Sciences 48: 2333-2343 <http://dx.doi.org/10.1139/f91-274>
- [30] Kumar, K., Eknath, A.E., Sahu, A.K., Mohanty, U.L., Kumar, R., Sahoo, M. and Noor, J. (2011). Snakeheads: challenging fish for diversification of fish farming. Fishing chimes, 31(1): 110-113.
- [31] Lalta, P., Dwivedi, A.K., Duvey, V.K. and Serajuddin, M. (2011). Reproductive biology of freshwater murrel, *Channa punctatus* (Bloch 1793) from river Varuna in India. Journal of Ecophysiology and Occupational Health 11(1/2): 69.
- [32] Long, D.N., Nguyen, V.T. and Lee, S.T. (2002). Technical aspects for artificial propagation of snakehead, *Ophiocephalus striatus* in Mekong Delta. Fisheries Sciences Institut Cantho University, Vietnam.
- [33] Mackenzie, D., Thomas, P. and Farrar, S.M. (1989). Seasonal changes in thyroid and reproductive steroids hormones in female channel catfish, *Ictalurus punctatus* in pond culture. Aquaculture 78: 63-80.
- [34] Marimuthu, K. and Haniffa, M.A. (2006). Studies on fecundity of captive reared spotted snakehead *Channa punctatus*. Journal of fisheries and Aquatic Science 1(3): 291-296
- [35] Mishra, S.K. (1991). Reproductive biology of a freshwater teleost, *Channa gachua* (Ham): Proceedings of the National Symposium on New Horizons in Freshwater Aquaculture. CIFA Bhubaneswar, India. pp. 55-56.
- [36] Mollah, M.F.A. and Tan, E.S.P. (1983). HCG – induced spawning of the catfish, *Clarius macrocephalus* (Gunther). Aquaculture 35: 239-247
- [37] Moore, Jr.W.T. and Ward, D.N. (1980). Pregnant mare serum gonadotropin: rapid chromatographic procedures for the purification of intact hormone and isolation of subunits. The Journal of Biological Chemistry 255 (14): 6923-6929.
- [38] Muthmainnah D. (2013). The length-weight relationship and condition factor of striped snakehead (*Channa striata* Bloch, 1793) grow out in swamp pond, South Sumatra Province. Depik 2(3): 184-190
- [39] Najero, N.T., Jalbani and Dastagir, G. (2015). Breeding biology of snakehead, *Channa striatus* (Bloch) from district Badin Sindh, Pakistan. Biolife 3(2): 434-436. <http://dx.doi.org/10.17812/blj2015.32.10>.
- [40] Quyen, N.T.K., Minh, T.H., Hai, T.N., Hien, T.T.T. and Dinh, T.D. (2016). Technical-economic efficiencies of snakehead seed production under impacts of climate change in the Mekong delta, Vietnam. Animal Review 3(4): 73-82. <http://dx.doi.org/10.18488/journal.ar/2016.3.4/101.4.73.82>
- [41] Rao, K., Podeti and Benarjee, G., (2015). Haematological change in freshwater fish, *Channa striatus* diagnosed with the Epizootic Ulcerative Syndrome (EUS). International Journal of Advanced Biotechnology and Research 6(2): 238-244
- [42] Rottmann, J.V. Shireman and Chapman, F.A. (1991). Hormonal control of reproduction in fish for induced spawning. The Southern Regional Aquaculture Center/SRAC Publication No. 424. p. 1-4
- [43] Saikia A.K., Abujam S.K.S., Biswas S.P. 2012. Food and feeding habit of *Channa punctatus* (Bloch) from the paddy field of Sivsagar District, Assam. Bulletin Environment Pharmacology and Life Sciences 1(5): 10- 15
- [44] Sakhare, V.B. (2015). Fecundity of air-breathing fish *Channa striatus* (Bloch) from waterbodies of Beed District, Maharashtra, India. International Journal of Aquaculture 5(18): 1-3
- [45] Sari, E. (2015). Rekayasa rematurasi ikan betok (*Anabas testudineus*) menggunakan hormon Oodev pada dosis berbeda melalui penyuntikan dengan rentang waktu 6 hari. Skripsi. Departemen Budidaya Perairan. Fakultas Perikanan dan Ilmu Kelautan. Institut Pertanian Bogor. Bogor. 36 hal.
- [46] Singh, M. and Serajuddin, M. (2017). Length-weight, length-length relationship and condition factor of *Channa punctatus* collected from three different rivers of India. Journal of Entomology and Zoology Studies 5(1): 191-197
- [47] Sinh, L.X., Navy, H. and Pomeroy, R. (2014). Value chain analysis of snakehead fish in the lower Mekong Basin of Cambodia and Vietnam. Aquaculture Economics and Management 18: 76-96

- [48] Song, L.M., Munian, K., Abd Rashid, Z. and Bhassu, S. (2013). Characterisation of Asian snakehead murrel *Channa striata* (channidae) in Malaysia: an insight into molecular data and morphological approach. The Scientific World Journal 2013: 1-16. <http://dx.doi.org/10.1155/2013/917506>
- [49] Supriyadi, (2005). Efektivitas pemberian Hormon 17 α -Metiltestosteron dan HCG yang Dienkapsulasi di dalam emulsi terhadap perkembangan gonad ikan Baung, *Hemibagrus nemurus*. [Tesis]. Program Studi Ilmu Perairan. Sekolah Pascasarjana IPB, Bogor.
- [50] Tiwari, K., Singh, B.K., Singh, S. and Tiwari, A. (2014). Study of gonadosomatic index of freshwater fish *Channa marulius*. International Journal of Scientific and Research Publications 4(5): 1-2.
- [51] Uthayakumar, V., Chandirasekar, R., Sreedevi, P.R., Senthilkumar, D., Jayakumar, R. and Ramasubramanian, V. (2014). Immunostimulatory effect and disease resistance induced by *Lawsonia inermis* against *Aphanomyces invadans* in striped murrels (*Channa striatus*). Malaya Journal of Biosciences 1(4): 231-241
- [52] Widodo, M.S., Marsodi, Susilawati, T., Agung Permana, W.M. (2013). Maturity level and somatic index of gonado at dwarf snake-head (*Channa gachua*) during January to December 2009. Journal of Basic applied Science Research 3(3): 387-393
- [53] Wiegand, M.D. (1984). Vitellogenesis in fishes. In: Reproductive physiology of fish. Edited by Richer, G.J. and Goss, H.J. Proceeding of International Symposium on reproductive Physiology in fish, p: 233 -241. Center for Agricultural Publishing and Documentation, Wegeningen.
- [54] Wijeyaratne, M.J.D. (1994). Some aspects of biology of the snakehead, *Ophicephalus striatus* Bloch in Muthurajawela, a peaty swamp in Sri Lanka. Vidyodaya Journal Sciences 5(1): 175-182.
- [55] Xie, Z.M., Li, W.S., Luo, M.L., Lin, G., Zheng, H.F. and Lin, X. (2002). The aquaculture techniques of the Chinese snakehead and the small snakehead, Beijing, China. China Agricultural University Press p. 208.
- [56] Xie, H., Lü, X., Zhou, J., Shi C., Li, Y., Duan, T., Ge, Li G. and Luo, Y. (2017). Effects of acute temperature change and temperature acclimation on the respiratory metabolism of the snakehead. Turkish Journal of Fisheries and Aquatic Sciences 17: 535-542. http://dx.doi.org/10.4194/1303-2712-v17_3_10
- [57] Zairin, M.Jr. (2000). Annual changes in ovarian maturity of female Thai catfish (*Clarias batrachus*) reared in a culture pond. Biotropia 15: 48-57.

Table.1: Tools, instruments and materials used in snakehead gonad rematuration experiment

No.	Tools and Materials	Size/Merk	Utility
1.	Concrete pond	4×7×1.2 m	Placement of hapa
2.	Small hapa 12 units	0.4×0.3×1 m	Placement of fish brood tested
3.	Large hapa 2 units	1.5×1.5 m	For brood fish reserves
4.	Watercheckker U10	Horiba	Measurement of water quality
5.	Scoop net	30 cm	For collecting the fish samples
6.	Name label	15×10 cm	Mark offhapa
7.	Digital balance	ACS	Weighing of fish samples
8.	Ruler	30 cm	Measurement of body length
9.	Washbasin	Ø 50 cm	Container for brood fish
10.	Knife	10 cm	Dissection of the selected brood fish
11.	Rope	10 m	Hapa installation
12.	Injection	1 ml	Oodev injection
13.	Snakehead brood	48 fish	Experimental fish sample
14.	Frogs	4% of body weight/day	Feed/bait served for snakehead
15.	Oodev hormone	0.5 ml kg ⁻¹ fish weight	Dose of oodev to be injected to snakehead fish brood samples.

Table.2: The experimental treatments for investigating the snakehead gonad rematuration

Treatment	Oodev dose/fish weight	Frequency	Time period
A	0.5 ml kg ⁻¹	3 times	Day-3, day-6, day-9
B	0.5 ml kg ⁻¹	2 times	Day-3, day-6
C	0.5 ml kg ⁻¹	1 time	Day-3
D	No injection	0 time	Control

Table 3: The mean \pm standar deviation of the parameters observed for each treatment.

Treatment	Parameters observed			
	Fecundity (granule/individual)	Egg diameter (mm)	GSI (%)	HSI (%)
A	19600 \pm 450.00 **	1.10 \pm 0.15 *	3.41 \pm 0.90 **	0.89 \pm 0.03
B	7300 \pm 556.77 *	0.90 \pm 0.11 *	1.39 \pm 0.14 *	1.11 \pm 0.03
C	1367 \pm 2367.13	0.20 \pm 0.40	0.40 \pm 0.62	1.53 \pm 0.23 *
D	1033 \pm 1789.79	0.20 \pm 0.40	0.50 \pm 0.54	0.93 \pm 0.09

Significance level: ** (p<0.01), and * (p<0.05)

Table 4: The mean \pm standar deviation of the estimated weights and their ratios for each treatment. A = three time injections, B = two time injections, C = once injection, and D = no injection (control).

The estimated weight, and the ratio	Treatment			
	A	B	C	D
Body weight (g)	266.8 \pm 727.78	253.67 \pm 21.33	258.03 \pm 10.13	266.57 \pm 11.60
Liver weight (g)	2.43 \pm 0.61	2.84 \pm 0.38	3.68 \pm 0.45	2.48 \pm 0.16
Gonad weight (g)	9.20 \pm 2.50*	3.83 \pm 0.31	2.10 \pm 1.49	1.33 \pm 1.53
Ratio of liver weight to body weight	0.0091	0.0112	0.0143*	0.0093
Ratio of gonad weight to body weight	0.0345*	0.0151	0.0081	0.0050

Significance level: ** (p<0.01), and * (p<0.05)

Table 5: Comparative fecundity, egg diameter, GSI and HIS of family Channidae from different geographical areas

Species	Fecundity (granule/ind)	Egg diameter (mm)	GSI %	HSI %	Locations	Country	References
<i>Channa striata</i>	1,033 - 19,600	0.20 - 1.10	0.40 - 3.41	0.89 - 1.53	Sungai Batang	Indonesia	Present study
<i>C. striata</i>	-	1.00 - 1.60	-	-	Danau Panggang	Indonesia	Fitriyani, 2005
<i>C. striatus</i>	28,332 - 41,068	1.21 - 1.33	10.4 - 11.5	0.70 - 1.40	Penang	Malaysia	Ghaedi <i>et al.</i> , 2013
<i>C. striatus</i>	3,000 - 12,000	0.70 - 1.30	2.10 - 8.00	-	Badin Sindh	Pakistan	Narejo <i>et al.</i> , 2015
<i>C. striatus</i>	4,900 - 14,028	1.53	-	-	Beed District	India	Sakhare, 2015
<i>C. punctatus</i>	2,538 - 32,987	-	0.53 - 5.34	-	Syilhet ditches	Bangladesh	Mian <i>et al.</i> , 2017
<i>C. punctatus</i>	2,116 - 11,332	-	-	-	Pond	India	Marimuthu and Haniffa, 2006
<i>C. punctatus</i>	3,678 - 27,853	-	-	-	Varuna River	India	Lalta <i>et al.</i> , 2011
<i>C. punctatus</i>	2,654 - 26,294	-	0.19 - 5.64	0.84 - 1.64	Syilhet	Bangladesh	Hossain, 2013
<i>C. gachua</i>	-	-	5.57 - 6.00	-	Godavari River	India	Gaikwad <i>et al.</i> , 2009
<i>C. gachua</i>	2,539 - 7,194	-	-	-	Bhubaneswar	India	Mishra, 1991
<i>C. gachua</i>	-	-	5.31 - 5.63	-	Dinoyo District	Indonesia	Widodo <i>et al.</i> , 2013
<i>C. limbata</i>	956 - 4,652	-	1.96 - 3.74	-	Ta Bo - Huai Yai	Thailand	Khomsab and Wannasri, 2017
<i>C. marulius</i>	-	-	8.21 - 47.56	-	Son River	India	Tiwari <i>et al.</i> , 2014.
<i>Ophiocephalus striatus</i>	-	0.20 - 1.60	-	-	Mekong River	Vietnam	Long <i>et al.</i> , 2002




Fig. 1: The ratio of fecundity to body weight (top) and the ratio of fecundity to body length (bottom) increase proportionally to the time periods of oodev intervention at the same amount of 0.5 ml kg^{-1} fish weight.

The treatment-A showed the highest ratios among other treatments ($p < 0.05$).

A = three time injections, B = two time injections, C = once injection, and D = no injection (control).

Oodev Injection Frequency and Time Period in Advancing Gonad Rematuration of Snakehead (Channa striata Blkr) in Hapa System

ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

9%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	sphinxesai.com Internet Source	3%
2	www.ciba.res.in Internet Source	1%
3	Dao Huy Giap. "Effects of different fertilization and feeding regimes on the production of integrated farming of rice and prawn <i>Macrobrachium rosenbergii</i> (De Man)", <i>Aquaculture Research</i> , 2/2005 Publication	1%
4	www.ukm.my Internet Source	1%
5	core.ac.uk Internet Source	1%
6	eprints.cmfri.org.in Internet Source	1%
7	krishikosh.egranth.ac.in Internet Source	1%

8	www.hindawi.com Internet Source	1 %
9	Submitted to Higher Education Commission Pakistan Student Paper	1 %
10	Submitted to Sogang University Student Paper	1 %
11	garuda.ristekbrin.go.id Internet Source	1 %
12	oceanrep.geomar.de Internet Source	<1 %
13	bioflux.com.ro Internet Source	<1 %
14	repository.ugm.ac.id Internet Source	<1 %
15	Hiroshige Tanaka, Toru Nakagawa, Takashi Yokota, Masayuki Chimura, Yuuho Yamashita, Tetsuichiro Funamoto. "Effects of non-spawning season (yolkless to early yolk formation stage) temperature on growth and reproductive characteristics in hatchery-reared, repeat-spawning walleye pollock <i>Gadus chalcogrammus</i> ", Fisheries Science, 2020 Publication	<1 %

16

Internet Source

<1 %

17

jurnal.ar-raniry.ac.id

Internet Source

<1 %

18

naturepub.org

Internet Source

<1 %

19

CD Ezeonyejiaku, MO Obiakor. "Metal enrichment in water and fish in a semi-urban Nigerian lake, and their associated risks", African Journal of Aquatic Science, 2016

Publication

<1 %

20

Senfan Ke, Zhiying Tu, Zhimin Li, Shoujiang Zhao, Defu Liu, Xiaotao Shi. " Effects of acute temperature changes on the swimming abilities and oxygen consumption of from the Lancang River ", Journal of Applied Ichthyology, 2019

Publication

<1 %

21

bioone.org

Internet Source

<1 %

22

textroad.com

Internet Source

<1 %

23

pdacrsp.oregonstate.edu

Internet Source

<1 %

24

M. E. CASTLE. "Silage and milk production: a comparison between soya, groundnut and

<1 %

single-cell protein as silage supplements",
Grass and Forage Science, 6/1979

Publication

25 Tien Hai Ly, Cheng - Ting Huang, Meng - Chou Lee, Vo Van Thanh, Doan Xuan Diep. " (Cuvier, 1831) Channidae, Perciformes , ", Journal of Fish Biology, 2023

Publication

26 "Culture of Cold - Water Marine Fish", Wiley, 2004

Publication

27 aquafishcrsp.oregonstate.edu

Internet Source

28 fishscientiae.ulm.ac.id

Internet Source

29 idoc.pub

Internet Source

30 jurnaltarbiyah.uinsu.ac.id

Internet Source

31 www.ijsrp.org

Internet Source

32 jifro.ir

Internet Source

33 journal.ipb.ac.id

Internet Source

34

ufdc.ufl.edu

Internet Source

<1 %

35

Evi Tahapari, Jadmiko Darmawan.
"KEBUTUHAN PROTEIN PAKAN UNTUK
PERFORMA OPTIMAL BENIH IKAN PATIN
PASUPATI (PANGASIID)", Jurnal Riset
Akuakultur, 2018

Publication

<1 %

36

Fatimat Adenike Adebisi, Siti Shapor Siraj,
Sharr Azni Harmin, Annie Christianus. "Plasma
sex steroid hormonal profile and gonad
histology during the annual reproductive cycle
of river catfish *Hemibagrus nemurus*
(Valenciennes, 1840) in captivity", Fish
Physiology and Biochemistry, 2012

Publication

<1 %

37

Hossain, Md. Belal, Md. Mosaddequr Rahman,
Md. Golam Sarwer, Md. Yusuf Ali, Ferdous
Ahamed, Sharmeen Rahman, Bernerd
Fulanda, Mohammad Mustafizur Rahman,
Bharat Raj Subba, and Md. Yeamin Hossain.
"Comparative Study of Carp Pituitary Gland
(PG) Extract and Synthetic Hormone Ovaprim
Used in the Induced Breeding of Stinging
Catfish, *Heteropneustes fossilis* (Siluriformes:
Heteropneustidae)", Our Nature, 2013.

Publication

<1 %

38

Md Saiful Islam, Md Kawser Ahmed, Mohammad Raknuzzaman, Md Habibullah-Al-Mamun, Shigeki Masunaga. "Metal Speciation in Sediment and Their Bioaccumulation in Fish Species of Three Urban Rivers in Bangladesh", Archives of Environmental Contamination and Toxicology, 2014

Publication

<1 %

39

Siti Maulida, Kartini Eriani, Nur Fadli, Firman M. Nur, Luvi S. Handayani, Zainal A. Muchlisin. " The effect of extender on the quality of climbing perch () sperm after refrigerated storage ", Aquaculture Research, 2022

Publication

<1 %

40

Y Kishimoto. "Sulindac and a cyclooxygenase-2 inhibitor, etodolac, increase APC mRNA in the colon of rats treated with azoxymethane", Gut, 2000

Publication

<1 %

41

dspace.stir.ac.uk

Internet Source

<1 %

42

eprints.icar.org.in

Internet Source

<1 %

43

oro.open.ac.uk

Internet Source

<1 %

44

www.intechopen.com

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On