



1. Submitted to the journal “Pakistan Journal of Phytopathology” (28-06-2023)
2. Review Version (26-11-2023)
3. Initiated (29-11-2023)
4. First revision: Accepted with minor revision (03-12-2023)
5. First revision submitted (06-12-2023)
6. Last Modified (07-12-2023)
7. Second revision: Minor revisions (05-12-2023)
8. Second revision submitted (07-12-2023)
9. Accept Submission (07-12-2023)
10. Copyediting (27-12-2023)



Pakistan Journal of Phytopathology
An official publication of Pakistan Phytopathological Society



HEC Recognized
ISSN: 1019-763X (Print), 2305-0284 (Online)

HOME
ABOUT
USER HOME
SEARCH
CURRENT
ARCHIVES
ANNOUNCEMENTS
THESIS ABSTRACTS

USER

You are logged in as...
saipul_abbas

- [My Profile](#)
- [Log Out](#)

Home > User > Author > Submissions > #1037 > Review

#1037 Review

SUMMARY REVIEW EDITING SUBMISSION

Authors	Mariana Mariana, Saipul Abbas, Salamiah Salamiah, Yusriadi Marsuni, Samharinto Soedijo, Muslimin Sepe, Muhammad I. Pramudi, Ismed S. Budi, Dewi Fitriyanti, Ely Liestiany, Lyswiana Aphrodyanti, Noor Aidawati, Helda O. Rosa
Title	IN SILICO ANALYSIS OF ASPERGILLUS FLAVUS FUNGAL PROTEINS: STRUCTURAL AND FUNCTIONAL INSIGHTS USING ITS PRIMERS
Section	Research Articles
Editor	Amajd Gondal

PEER REVIEW

Round 1

Review Version	1037-4555-1-RV.DOCX	2023-11-26
Initiated	2023-11-29	
Last modified	2023-12-07	
Uploaded file	Reviewer B 1037-4569-1-RV.PDF	2023-12-03
	Reviewer A 1037-4581-1-RV.DOCX	2023-12-05

EDITOR DECISION

Decision	Accept Submission	2023-12-07
Notify Editor	Editor/Author Email Record	2023-12-07
Editor Version	None	
Author Version	1037-4582-1-ED.DOCX	2023-12-06 DELETE
	1037-4582-2-ED.DOCX	2023-12-07 DELETE
	1037-4582-3-ED.DOCX	2023-12-09 DELETE
	1037-4582-4-ED.DOCX	2023-12-27 DELETE

Upload Author Version

LANGUAGE

Select Language

English

KEYWORDS

Alternaria solani Biological control Botanicals Cultivars Eggplant FHB species Fungi Fungicides Fusarium oxysporum Incidence Isolation Petri-dish assay Plant extracts Rice Trichoderma spp Wheat disease incidence identification pathogenic variation races virulence

JOURNAL CONTENT

Search

Search Scope

All

Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)

AUTHOR


Submissions

- [Active \(2\)](#)
- [Archive \(2\)](#)

Accept Submission (07-12-2023)

← 📄 🕒 🗑️ 📧 🕒 🔄 📁 🗑️ ⋮ 290 of 962 < > 📧

[PJP] Editor Decision **External** **Inbox** x


 **Dr. Muhammad Atiq** <matiq@pakps.com>
to me ▾ Thu, Dec 7, 2023, 1:25 PM ☆ ↶ ⋮

Saipul Abbas:

We have reached a decision regarding your submission to Pakistan Journal of Phytopathology, "In Silico Analysis of Aspergillus flavus Fungal Proteins: Structural and Functional Insights Using ITS Primers".

Our decision is to:

Dr. Muhammad Atiq
Associate Professor
Pakistan Journal of Phytopathology/
Phone 0333-4396981
matiq@pakps.com
<http://pjp.pakps.com>

 **Dr. Muhammad Atiq** <matiq@pakps.com>
to me ▾ Thu, Dec 7, 2023, 1:37 PM ☆ ↶ ⋮

Saipul Abbas:

We have reached a decision regarding your submission to Pakistan Journal of Phytopathology, "In Silico Analysis of Aspergillus flavus Fungal Proteins: Structural and Functional Insights Using ITS Primers".


Our decision is to: Accept Submission

Dr. Muhammad Atiq
Associate Professor
Pakistan Journal of Phytopathology/
Phone 0333-4396981
matiq@pakps.com
<http://pjp.pakps.com>

Copyediting (27-12-2023)

← 📄 🕒 🗑️ 📧 🕒 🔄 📁 🗑️ ⋮ 257 of 962 < > 📧

Galley **External** **Inbox** x


 **Muhammad Atiq** <matiq@pakps.com>
to me ▾ Wed, Dec 27, 2023, 1:27 PM ☆ ↶ ⋮


AOA

Respected author,

Piz see the attachment regarding galley of yours accepted paper. Put contribution of authors at the end of paper in a box. and send back me paper upto tomorrow because journal is ready to publish on 30th December.

One attachment • Scanned by Gmail ⓘ



 **SAIPUL ABBAS** <saipul.abbas@ulm.ac.id>
to Muhammad ▾ Wed, Dec 27, 2023, 2:32 PM ☆ ↶ ⋮

Dear Editor, Pakistan Journal of Phytopathology

I hope this email finds you well. I have completed the addition of the Author Contribution section at the end of the manuscript, and Proof of Journal Payment.

Thank you for your time and for the chance to submit my work.

Best regards,

Saipul Abbas
Lambung Mangkurat University
081533333333

Layout (29-12-2023)

LAYOUT

Layout Editor	None				
Layout Version		REQUEST	UNDERWAY	COMPLETE	VIEWS
None		—	—	—	
Galley Format		FILE			
1. PDF VIEW PROOF		1037-4643-1-PB.PDF	2023-12-29		101
Supplementary Files		FILE			
			None		

Published (30-12-2023)

AUTHOR FEES

Article Publication Fee Paid December 29, 2023 – 07:28 AM

STATUS

Status Published Vol 35, No 2 (2023): Pak. J. Phytopath.
Initiated 2023-12-30
Last modified 2024-01-02

<https://www.pjp.pakps.com/index.php/PJP/article/view/1037>

Pakistan Journal of Phytopathology

An official publication of Pakistan Phytopathological Society



HEC Recognized

ISSN: 1019-763X (Print), 2305-0284 (Online)

ER HOME	SEARCH	CURRENT	ARCHIVES	ANNOUNCEMENTS	THESES ABSTRACTS
-------------------------	------------------------	-------------------------	--------------------------	-------------------------------	----------------------------------

[Home](#) > [Vol 35, No 2 \(2023\)](#) > [Mariana](#)

IN SILICO ANALYSIS OF ASPERGILLUS FLAVUS FUNGAL PROTEINS: STRUCTURAL AND FUNCTIONAL INSIGHTS USING ITS PRIMERS

Mariana Mariana, Saipul Abbas, Salamiah Salamiah, Yusriadi Marsuni, Samharinto Soedjjo, Muslimin Sepe, Muhammad I. Pramudi, Ismed S. Budi, Dewi Fitriyanti, Elly Liestiany, Lyswiana Aphrodyanti, Noor Aidawati, Helda O. Rosa

Abstract

Aspergillus spp. is a type of fungi that can contaminate and damage various types of food and agricultural products. To better understand the structure and function of proteins, *in silico* research was conducted using Internal Transcribed Spacer (ITS) primers and SWISS Model protein prediction tools. The objective of this study was to identify fungal species using the BLAST method and to analyze the structure and function of proteins from *Aspergillus* spp. The methods used included BLAST for species identification, Web Expsy to translate DNA sequences into proteins, SWISS Models to model protein structures, SAVES to validate protein structures, and STRING to analyze the function of proteins. The results of the BLAST analysis showed that the identified fungal species were *Aspergillus flavus*, *A. tamarii*, and *A. nomius*. Furthermore, the results of translating DNA sequences into proteins using Web Expsy showed that there were three open reading frames with the highest residual values of 119 and 83, while the lowest residual value was 4. Only two of these frames met the protein criteria. Moreover, the results of protein structure modeling using the SWISS Model method produced a fairly accurate *Aspergillus* spp. protein structure model with a validation value of protein structure using ERRAT (SAVES V6.0) of 100%. Additionally, the results of protein function analysis using STRING showed that the *Aspergillus* spp. protein has a function in producing enzymes that play a role in the metabolic process of cells.

Keywords

Aspergillus flavus; in silico; ITS primer; protein structure; protein function; SWISS Model.

LANGUAGE

Select Language
English
Submit

KEYWORDS

Alternaria solani Biological control Botanicals Cultivars Eggplant FHB species Fungi Fungicides Fusarium oxysporum Incidence Isolation Petri-dish assay Plant extracts Rice Trichoderma spp Wheat disease incidence identification pathogenic variation races virulence

In Silico Analysis of *Aspergillus flavus* Fungal Proteins: Structural and Functional Insights Using ITS Primer

Mariana Mariana, Saipul Abbas*, Salamiah Salamiah, Samharinto Samharinto, Yusri
Muhammad Indar Pramudi, Ismed Setya Budi, Dewi Fitriyanti, Elly Liestiany, L

Department of Plant Protection, Faculty of Agriculture, Lambung Mangkurat University,
Banjarbaru. South Kalimantan 70714 Indonesia

*Corresponding author: Email: saipul.abbas@ulm.ac.id

ABSTRACT

Aspergillus flavus is a type of fungus that can contaminate and damage agricultural products. To better understand the structure and function of *Aspergillus flavus*, research was conducted using Internal Transcribed Spacer (ITS) primer and protein prediction tools. The objective of this study was to identify fungal species using BLAST method and to analyze the structure and function of proteins from *Aspergillus spp.* methods used included BLAST for species identification, Web Expasy for translating DNA sequences into proteins, SWISS Models to model protein structures, SAHARA for protein structures, and STRING to analyze the function of proteins. The results showed that the identified fungal species were *Aspergillus flavus*, *Aspergillus tamarii*, and *Aspergillus nomius*. Furthermore, the results of translating DNA sequences into proteins using Web Expasy showed that there were three open reading frames with the highest residual values of 119 and 83, while the lowest residual value was 4. Only two of these frames met the protein criteria. Moreover, the results of protein structure modeling using the SWISS Model method produced a fairly accurate *Aspergillus spp.* protein structure model with a validation value of protein structure using ERRAT (SAVES V6.0) of 100%. Additionally, the results of protein function analysis using STRING showed that the *Aspergillus spp.* protein has a function in producing enzymes that play a role in the metabolic process of cells.

Keywords: *Aspergillus flavus*, in silico, ITS primer, protein structure, protein structure model

INTRODUCTION

Aspergillus spp. fungus is a type of fungus that often causes contamination and damage to various agricultural and food products (Gourama & Bullerman, 1995) and processed products (Ayofemi Olalekan Adeyeye, 2020). One of the *Aspergillus spp.* that can contaminate feed ingredients is *the species Aspergillus* can cause aflatoxin pollution (Klich, 2007), which is a harmful substance to human health and livestock (Balina et al, 2018). *Aspergillus flavus* is a group of fungi in agricultural environments, especially in food crops such as corn, rice, and wheat (Keller, 2011). Contamination by *Aspergillus spp.* can cause damage to crops and foodstuffs and can cause health problems in humans and farm animals (Yu et al, 2005). Therefore, research on *Aspergillus spp.* is very important in agriculture to prevent and control contamination of *Aspergillus spp.* on agricultural products and foodstuffs.

Previous studies have highlighted the negative impact of *Aspergillus* spp. on agricultural products and food items. Ramirez et al. (2018) noted that this type of fungus can cause damage to grains, legumes, and processed products. Similarly, Balina et al. (2018) indicated that *A. flavus* can be a major cause of aflatoxin contamination in feed materials, posing risks to livestock and humans who consume them. The significance of research on *A. flavus* is further underscored by the findings of Bhatnagar-Mathur et al. (2015), demonstrating the frequent presence of *A. flavus* in agricultural environments, with a high tendency to infect staple food crops. Therefore, contamination risk from *A. flavus* not only affects the quality of agricultural products and food items but also poses a serious health risk to consumers. In silico research has become a useful tool for studying the structure and function of proteins (Dutta et al, 2018), as it allows for quick and efficient hypothesis testing and analysis (Kirubakaran et al, 2013). In this study, we used an in silico technique to predict the structure and function of the *A. flavus* protein by combining the primary use of the Internal Transcribed Spacer (ITS) and the SWISS Model protein prediction tool. This in silico approach is essential to understanding this fungus and is beneficial to the food and agriculture industries.

The formulation of the problem of this study was to analyze the BLAST results of these *Aspergillus* spp sequences, examine the protein structure related to contamination, evaluate the validity and confidence of the protein structure of *Aspergillus* spp. which was found, as well as examining the function of the protein *Aspergillus* spp. associated with contamination of corn feed and other foodstuffs, as well as to understand the structure and function of the *Aspergillus* spp protein associated with the contamination aforementioned. Therefore, this research focused on the analysis of *Aspergillus* spp. sequence, protein structure analysis, and protein function analysis, with the hope of providing new insights and solutions to the problem of contamination in food.

This study aims to analyze the species *Aspergillus* spp. which often contaminates corn feed and other foodstuffs. This study is expected to provide information about the species *Aspergillus* spp. which contaminates corn feed and other foodstuffs, as well as the structure and function of the protein *Aspergillus* spp. The results of this study will contribute to the development of methods for detecting and preventing *Aspergillus* spp. in corn feed and other foodstuffs.

nsakr
2023-12-03 09:14:48
ion of

please select another word in place of it is,
used in the previous sentence.

MATERIALS AND METHODS

Identification of fungus species

This step involved the use of an Internal Transcribed Spacer (ITS) Primer to verify fungus species. This stage of research was carried out by taking DNA samples from fungi and conducting DNA amplification using primary ITS. DNA amplification was performed using the Polymerase Chain Reaction (PCR) technique. This step aimed to verify the species of fungus analyzed. In the PCR technique, DNA samples were placed together with ITS primers (ITS1 5'-TCCGTAGGTGAACCTGCGG-3', and ITS4 5'-TCCTCCGCTTATTGATATGC-3') and polymerase enzymes, and a heating and cooling process was carried out to start and end the reaction. The

result of the amplification process was more DNA sequences associated with ITS. After successful DNA amplification, the obtained DNA sequence was analyzed to verify the fungus species. Sequencing analysis was carried out using bioinformatics software such as BLAST (*Basic Local Alignment Search Tool*) (Lobo, 2008).

Translate DNA Sequencing into Proteins

After the fungus DNA sequence was successfully analyzed, the DNA sequence was then translated into proteins. This step was done using a tool such as the ExPASy Translate Tool (Bashir, 2022). This tool converted the DNA sequence into a protein sequence based on the genetic code (Ayaz et al, 2020). The results of the translation showed the amino acid sequence contained in the fungus protein.

Analysis of protein structure

This stage used the SWISS Model method to predict the protein structure of the fungus *Aspergillus* spp. At this stage, the input of protein sequencing data from ExPASy tools was carried out, and then protein modeling was made using the SWISS Model (Waterhouse et al, 2018). Analysis using the SWISS Model provided information regarding protein domains, protein motifs, and other information related to the structure and function of proteins (Bordoli & Schwede, 2012). The results of the analysis of the structure and function of the protein were re-examined to ensure the validity of the results. Validation was done by comparing the results with a trusted protein database such as UniProt.

Protein Function Analysis

The research method of protein function analysis with STRING involved the use of a database of protein interactions and the analysis of protein networks (Franceschini et al, 2012). In this study, *Aspergillus flavus* protein data were used in the STRING database. Protein data were inputted into a STRING database containing information about protein interactions formed based on data from literature databases, scientific publications, and experimental studies (Szklarczyk et al, 2010). Analysis of protein interactions formed using algorithms available on the STRING database. This algorithm generated information about the interaction of formed proteins, the forces of interaction, and their relationship with biological functions. Interpretation of the results of protein interaction analysis was done to obtain information about the biological function of the protein being analyzed and how the protein interaction was related to the biological function found. This method could be used to analyze the function of proteins and understand how proteins interacted with each other in a network of proteins (Szklarczyk et al, 2023). Using this method, we could gain a better understanding of how proteins were involved in biological processes and their possible effects on health and disease.

RESULTS AND DISCUSSION

Identification of fungus species by BLAST Method

The results of the fungus species identification using BLAST revealed a similarity of 99.61% with *Aspergillus tamarii*, 98.94% with *Aspergillus flavus*, and 96.33% with *Aspergillus nomius*. These findings suggested that the sample likely belonged to the *Aspergillus* group, with a higher similarity to *Aspergillus tamarii* compared to the other two species. *Aspergillus tamarii* is recognized for its capability to produce various bioactive compounds (Bose et al, 2019), including phenol, flavonoid and IAA and its antifungal activity, applications in the Agriculture and food industries. However, it is in *Aspergillus tamarii* can also cause infections in humans and animals. Similarly, *Aspergillus flavus* is known for its production of toxic compounds, which can be harmful to human and animal health (Fouad et al, 2019). Therefore, proper handling measures are necessary to prevent the spread of this species. Moreover, *Aspergillus nomius*, another identified species in the sample, can produce aflatoxin and the potential to become pathogenic in humans and animals (Zain, 2011). Consequently, accurate identification of the fungus species is essential to determine appropriate control measures. In this case, the identification results indicating the highest similarity to *Aspergillus tamarii* and *Aspergillus flavus* can serve as references for further research on the bioactive compounds produced by these species. Nevertheless, further examinations are required to confirm the presence of specific fungus species in the sample and establish appropriate control measures to prevent its spread.

Description	Scientific Name	Max Score	Total Score	Query Cover	E value	Per. Ident	Acc. Len	Accession
Aspergillus tamarii strain M24 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8S...	<i>Aspergillus tamarii</i>	1419	2347	97%	0.0	99.61%	889	ON819577.1
Aspergillus flavus strain Bp5 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1 and 5.8S riboso...	<i>Aspergillus flavus</i>	1343	2094	94%	0.0	98.94%	939	KF221065.1
Aspergillus tamarii strain BPFM17 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1,...	<i>Aspergillus tamarii</i>	1321	2151	99%	0.0	96.86%	897	MF359724.1
Aspergillus tamarii strain MF24 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8...	<i>Aspergillus tamarii</i>	1310	1952	98%	0.0	96.83%	791	MF359732.1
Aspergillus sp. AQQWD 17 internal transcribed spacer 1, partial sequence; 5.8S ribosomal RNA gene and internal tran...	<i>Aspergillus sp. A...</i>	1297	2462	100%	0.0	96.37%	1303	KP721583.1
Aspergillus sp. isolate ACSIKS_2100168 small subunit ribosomal RNA gene, partial sequence; internal transcribed spa...	<i>Aspergillus sp.</i>	1291	2430	100%	0.0	96.25%	1187	MN597042.1
Aspergillus sp. isolate SS_12 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8S...	<i>Aspergillus sp.</i>	1286	2440	100%	0.0	96.12%	1303	MT497446.1
Aspergillus flavus isolate 20 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1 and 5.8...	<i>Aspergillus flavus</i>	1273	1946	91%	0.0	98.18%	715	MK120614.1
Aspergillus nomius strain Tur4 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1 and 5.8S ribos...	<i>Aspergillus nomiae</i>	1240	1891	95%	0.0	96.33%	842	KF221090.1
Aspergillus flavus strain Beca_43 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5...	<i>Aspergillus flavus</i>	1234	2105	99%	0.0	96.80%	849	KY234269.1

Figure 1: The results of the fungus species identification using BLAST Method (GenBank)

Translate DNA Sequencing into Proteins

The results obtained from the translation of *Aspergillus* spp DNA sequences into proteins using the ExPASy web revealed the presence of three open reading frames (ORFs), indicated in red, with residual values of 119 and 83 as the highest, and 4 as the lowest. Proteins are typically categorized if they consist of more than 50 residues (Damodaran & Parkin, 2017). In this case, only two out of the three ORFs met the criteria to be considered as proteins. The identification of an ORF qualifying as a protein may indicate the presence of a gene within the *Aspergillus* DNA sequence. Translating DNA sequences into proteins holds significant importance in molecular biology research as it allows for a better understanding of protein functions within a specific organism (Gasteiger et al, 2003). Consequently, these findings serve as a foundation for future investigations concerning the *Aspergillus* gene and its potential

applications in various fields, such as agriculture and the food industry. Nevertheless, it is crucial to note that these results should undergo further verification using other techniques, such as protein structure analysis, to ensure the functional integrity of the resulting proteins. Additionally, the authenticity and suitability of the *Aspergillus* DNA sequence utilized in this study should be validated about the specific *Aspergillus* species under investigation.



Figure 2: The results of *Aspergillus spp* DNA sequences into proteins using the ExpASY web

Table 1: The results of Open Reading Frames (ORFs) with residual values from 4 to 119

Open Reading Frame	Residue	Types of Molecules	Code Peptides/Proteins
MNGGPAEATKVQ	12	Peptide	A1
MAGPPKQLRYSKHGWEVGLARNPTLGND PSVGEPAEGSLPSVGFASPTSHPCLLYLSC FGGPAIHGRRGLSAPGPRPETPRTLSDLV KSELIVSQSVKTFNNGSLGSGIDEERSEMR	119	Protein	A2
MPVRASLLPIKHGLCVGSSSPLRGGGRAPKA AAAPRILERMGLCHPLCRPGRRLPNANQS FSRLTSDQVGI PAELKHINKAE	83	Protein	A3
MILP	4	Peptide	A4
MAY	4	Peptide	A5
MAAGGSQPRARARRRHHHELCLI	22	Peptide	A6
MDLLVPASMKNAAKCDN	17	Peptide	A7

Protein Structure Modeling with the SWISS Model method

The results of modeling the protein structure of *Aspergillus spp* using the SWISS Model method show the presence of a 3D structure of proteins formed (Bordoli & Schwede, 2012). Protein structure modeling is an important method in molecular biology research because it can provide information about the shape and function of proteins in a particular organism (Breda et al, 2007). The results of modeling the protein structure of *Aspergillus spp* can be used as a basis for conducting further analysis of protein function, including interactions with other molecules, enzymatic activity, and possible drug development. In addition, these results can also help in understanding the relationship between the structure of proteins and their biological activity. However, keep in mind that the results of modeling protein structures using computational methods such as the SWISS Model still have limitations, especially in terms of accuracy. Therefore, these results still need to be verified using other techniques, such as X-ray crystallography or core magnetic resonance spectroscopy (NMR), to ensure the accuracy of the resulting protein structure. In addition, modeling the protein structure of *Aspergillus spp* can also provide information about the evolutionary relationship of proteins between different *Aspergillus* species. Modeling the structure of the *Aspergillus spp* protein

can help understand the differences and similarities between different *Aspergillus* proteins so that it can be used as a basis for specific drug and therapeutic development.




Figure 3: Protein Structure Prediction from Internal Transcribed Spacer (ITS) primer of *Aspergillus flavus* fungus using Protein Prediction from SWISS Model. (Template: A2: 3lb6.1.A A3: 5t7v.1.K)



The results of validation of the *Aspergillus spp* protein structure using SAVES V6.0 show that the resulting protein structure has a fairly good value in terms of accuracy and quality. One of the parameters used to evaluate the validation of protein structure is the Ramachandran plot, which shows the proportion of amino acids placed in the most favored regions within the protein conformation space (Laskowski et al, 2013). In the Ramachandran plot results obtained, the highest Residual value in most favored regions was 80% for the protein code A2, which indicates that most of the amino acids in the protein structure of *Aspergillus spp* were placed in the desired area (Lovell et al, 2003). In addition, the validation results using the ERRAT method also show the highest value of 100%, which indicates that the resulting protein structure has high accuracy (Rozario et al, 2021). The results of the validation of the protein structure of *Aspergillus spp* using SAVES V6.0 are important to ensure that the resulting protein structure is trustworthy and used as a basis for further analysis of protein function and drug development. Validation of protein structure can also help identify and correct parts in the protein structure that may be inaccurate, thereby improving the quality and accuracy of research results.

The QMEAN4 Score indicates the overall quality of the protein structure, which is calculated based on geometric values, non-covalent interactions, and consistency of the structure (Benkert et al, 2008). The higher the QMEAN 4 Score, the better the quality of the resulting protein structure (Benkert et al, 2011). In this case, the QMEAN4 Score A2 value of -1.72 versus A3 of -3.27 can be interpreted as a fairly low value, which indicates that the resulting *Aspergillus spp* protein structure does not have optimal quality. Meanwhile, the Z Score is a parameter used to compare the quality of the resulting protein structure with other similar protein structures, which are calculated based on geometric values and structural consistency. The higher the Z Score value, the better the quality of the resulting protein structure compared to other similar protein structures (Bhattacharya et al, 2007). In this case, a Z Score of -0.16 indicates that the resulting *Aspergillus spp* protein structure is of relatively good quality compared to other similar protein structures. However, keep in mind that as with the QMEAN 4 Score value, the Z Score value also has limitations and possible errors, so this result still needs to be verified using other protein structure validation techniques.


Table 2: Protein Structure Validation using SAVES V6.0

Protein Code	Ramachandran plot				ERRAT quality factor (%)	QMEAN 4 Score	Z Score
	Residu in most favored regions (%)	Residu in additional allowed regions (%)	Residu in generously allowed regions (%)	Residu in disallowed regions (%)			
A2	80.0	20.0	0.0	0.0	100	-1.72	-0.16
A3	75.0	20.0	0.0	5.0	6.6666	-3.27	-3.32


Model A2



PROCHECK



Ramachandran Plot



Model A3




Figure 4: The results of validation of the *Aspergillus spp* protein structure using SAVES V6.0

Protein Function & Interaction of *Aspergillus Flavus*

Aspergillus flavus is a type of fungus known to cause damage to agricultural products such as grains and other foodstuffs (Gourama & Bullerman, 1995). This fungus can also produce some toxic substances such as aflatoxins, which are heterocyclic substances and have detrimental effects on human and animal health (Dhanamjayulu et al, 2019). Heterocyclics in aflatoxins have several main functions in the fungus *Aspergillus flavus*, including: (1) As a toxic compound: Aflatoxins have high toxic activity and can cause food poisoning in humans and animals, which can cause various health problems such as liver damage, cancer, and reproductive problems (Ogodo & Ugbogu, 2016). (2) As an antioxidant compound: Aflatoxin has antioxidant properties and can help the fungus *Aspergillus flavus* survive in a poor environment (Abrar et al, 2013). (3) As an antifungal compound: Aflatoxin can help the fungus *Aspergillus flavus* resist attacks from other fungi that seek to interfere with the growth of the

fungus *Aspergillus flavus* (Bhatnagar-Mathur et al, 2015). Overall, heterocyclic in aflatoxins has an important role in the biology of the fungus *Aspergillus flavus*, but it also has detrimental effects on human and animal health. Therefore, it is important to control the production of aflatoxins and ensure that the food products consumed do not contain this substance in harmful quantities.

Organic cyclic compound binding is the process of binding organic chemical compounds that have a benzene ring (Cram & Cram, 1971). In this case, such binding may take place in the fungus *Aspergillus flavus*. Organic chemical compounds that bind to the fungus *Aspergillus flavus* can affect the biological activity of the fungus (Nguyen et al, 2019). Some organic chemical compounds that can bind to the fungus *Aspergillus flavus* include growth regulators, metabolic regulators, and anti-fungal substances (Hou et al, 2022). The binding of organic chemical compounds to the fungus *Aspergillus flavus* can affect the growth and reproduction of fungi, modulate enzyme activity, affect the production of toxic substances such as aflatoxins, and affect the interaction of fungi with their environment. Overall, the binding function of organic chemical compounds to the fungus *Aspergillus flavus* is essential for understanding the biology of the fungus and how the fungus interacts with its environment. Further research on the binding of organic chemical compounds to the fungus *Aspergillus flavus* can help in controlling the production of aflatoxins and improving the quality of food products.

Nucleic acid binding protein has a very important function in the fungal cells of *Aspergillus flavus*. Nucleic acids are important molecules that carry genetic information and play a key role in various biological processes, such as DNA replication, RNA transcription, and protein synthesis (Blackburn, 2006). Nucleic acid binding proteins can bind to nucleic acid molecules, facilitate interaction with enzymes and transcription factors, and influence the biological activity of nucleic acid molecules (Von Hippel et al, 1984). In the fungus *Aspergillus flavus*, Nucleic acid binding proteins may play a role in regulating gene activity, affecting growth and reproduction, and modulating the fungus' response to the environment. Studies of Nucleic acid binding proteins in the fungus *Aspergillus flavus* can provide information about the mechanisms of gene regulation and how fungi react to their environment. Overall, the function of the Nucleic acid binding protein in the fungus *Aspergillus flavus* is essential for understanding the biology of the fungus and how the fungus interacts with its environment. More research on Nucleic acid binding proteins in the fungus *Aspergillus flavus* can help understand the growth and reproduction mechanisms of fungi, as well as how fungi react to their environment.

RNA binding is a process in which proteins bind to RNA molecules. RNA is an important molecule in cells that plays a key role in biological processes such as transcription, protein synthesis, and gene regulation (Elliott & Ladomery, 2017). RNA protein binding can bind to RNA molecules and facilitate interaction with enzymes and transcription factors (Corley et al, 2020). The function of RNA binding proteins can affect the biological activities of RNA molecules, such as modulating transcription and protein synthesis, affecting RNA stability, and facilitating gene control (Gilsovic et al, 2008). In the fungus, *Aspergillus flavus*, protein RNA binding may play an important role in regulating gene activity, modulating responses to the environment, and facilitating growth and reproduction. Studies on protein RNA binding in the fungus *Aspergillus flavus* may help understand the mechanisms of gene regulation and

how fungi react to their environment. Overall, the function of RNA binding proteins in the fungus *Aspergillus flavus* is critical to understanding the biology of the fungus and how the fungus interacts with its environment. More research on protein RNA binding in the fungus *Aspergillus flavus* may help in understanding the mechanisms of fungal growth and reproduction, as well as how the fungus reacts to its environment.

mRNA binding is a process in which proteins bind to mRNA molecules (Dreyfuss et al, 2002). mRNA is an RNA molecule that contains genetic information from DNA and is used as a template for protein synthesis (Alberts et al, 2002). mRNA protein binding can bind to mRNA molecules and facilitate interaction with enzymes and transcription factors (Corley et al, 2020). In the fungus *Aspergillus flavus*, mRNA protein binding may play an important role in regulating gene activity, modulating responses to the environment, and facilitating growth and reproduction. Studies on mRNA protein binding in the fungus *Aspergillus flavus* may help understand the mechanisms of gene regulation and how fungi react to their environment. Overall, the function of mRNA protein binding in the fungus *Aspergillus flavus* is critical to understanding the biology of the fungus and how the fungus interacts with its environment. More research on mRNA protein binding in the fungus *Aspergillus flavus* may help understand the mechanisms of fungal growth and reproduction, as well as how the fungus reacts to its environment.

Ribosomes are cellular structures that play a key role in the process of protein synthesis (Frank & Spahn, 2006). Ribosomes consist of two subunits, a large subunit and a small subunit, which serve to bind amino acids and form polypeptide chains (Lake, 1981). Proteins that are structural components of ribosomes can play an important role in ensuring the integrity and activity of ribosomes. The function of proteins as structural components of ribosomes can affect the ability of ribosomes to bind to mRNA and facilitate proper protein synthesis (de la Cruz et al, 2015). In the fungus *Aspergillus flavus*, proteins that are structural components of ribosomes may play an important role in ensuring the growth and reproduction of fungi. The study of proteins that are structural components of ribosomes in the fungus *Aspergillus flavus* can help understand the mechanism of protein synthesis and how the fungus reacts to its environment. Overall, the function of proteins as structural components of ribosomes in the fungus *Aspergillus flavus* is essential for understanding the biology of the fungus and how the fungus interacts with its environment. More research on the proteins that are structural components of ribosomes in the fungus *Aspergillus flavus* can help in understanding the mechanisms of fungal growth and reproduction, as well as how the fungus reacts to its environment.




Figure 5: Network of protein-protein interactions visualized with STRING (NCBI taxon-Id: 332952)

B8MXN8	<i>Serine/threonine protein kinase, putative</i>
B8N3J1	<i>Small nucleolar ribonucleoprotein complex subunit Utp14, putative</i>
B8NT05	<i>G-protein complex beta subunit CpcB</i>
B8N430	<i>Nuclear mRNA splicing factor, putative</i>
B8N0Y2	<i>G-patch domain protein (TFIP11), putative</i>
B8N058	<i>PWI domain mRNA processing protein, putative</i>
B8N1X2	<i>Pre-RNA splicing factor Srp2, putative</i>
B8ND38	<i>CaaX prenyl proteinase Rce1</i>
B8MXA7	<i>Small nuclear ribonucleoprotein E; Associated with the spliceosome snRNP U1, U2, U4/U6 and U5.</i>
B8MXX7	<i>Casein kinase, putative; Belongs to the protein kinase superfamily.</i>

Table 3: Molecular Function, Biological Process, Cellular Component, and Subcellular Localization Classification

Molecular Function	Biological Process	Cellular Component	Subcellular Localization
<ul style="list-style-type: none"> Heterocyclic compound binding Organic cyclic compound binding 	<ul style="list-style-type: none"> mRNA splicing, via spliceosome RNA processing Gene expression 	<ul style="list-style-type: none"> Spliceosomal complex Ribonucleoprotein complex Nucleus 	<ul style="list-style-type: none"> U2 snRNP U2-type prespliceosome U2-type spliceosomal complex

• Nucleic acid binding	• Macromolecule metabolic process	• Intracellular membrane-bounded organelle	• Spliceosomal complex
• RNA binding	• Cellular nitrogen compound		• Ribonucleoprotein complex
• mRNA binding	• metabolic process		• Nucleus
• Structural constituent of ribosome	• Nitrogen compound metabolic process		• Protein-containing complex
• rRNA binding	• Primary metabolic process		• Intracellular membrane-bounded organelle
• snoRNA binding	• Cellular metabolic process		• Intracellular
			• Cellular anatomical entity

Conclusion

In silico research was conducted to understand the structure and function of *proteins of Aspergillus spp.* The results of the BLAST analysis showed the presence of three species of fungi, namely *Aspergillus flavus*, *Aspergillus tamarii*, and *Aspergillus nomius*. Translating DNA sequences into proteins using Web Expasy showed that only two proteins met the criteria. Protein structure modeling using SWISS Model resulted in a fairly accurate *Aspergillus spp* protein structure model with protein structure validation values using ERRAT (SAVES V6.0) of 100%. Analysis of protein function using STRING shows that *Aspergillus spp* protein has a function in producing enzymes that play a role in the metabolic process of cells. Thus, this study provides important information about the structure and function of the *protein Aspergillus spp*, which can be used to improve understanding of contamination and damage to food and agricultural products by this type of fungus.

Acknowledgement

A thank you to PT. Arutmin Indonesia Tambang Satui and Agreement/Contract Number: 39/UN8.1.23/PP/2022 through the Matching Fund program "Kedaireka" for financial support for this research activity, and all parties who have helped in the success of this research. All the help and support provided means a lot to us. Thanks.

Reference

- Ayaz, A., Zaman, W., Saqib, S., Ullah, F., & Mahmood, T. (2020). Phylogeny and diversity of Lamiaceae based on rps14 gene in Pakistan. *Genetika*, 52(2), 435-452. <https://doi.org/10.2298/GENSR2002435A>
- Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., & Walter, P. (2002). From DNA to RNA. In *Molecular Biology of the Cell. 4th edition*. Garland Science.

- Abrar, M., Anjum, F. M., Butt, M. S., Pasha, I., Randhawa, M. A., Saeed, F., & Waqas, K. (2013). Aflatoxins: biosynthesis, occurrence, toxicity, and remedies. *Critical reviews in food science and nutrition*, 53(8), 862-874. <https://doi.org/10.1080/10408398.2011.563154>
- Ayofemi Olalekan Adeyeye, S. (2020). Aflatoxigenic fungi and mycotoxins in food: a review. *Critical reviews in food science and nutrition*, 60(5), 709-721. <https://doi.org/10.1080/10408398.2018.1548429>
- Amaiike, S., & Keller, N. P. (2011). *Aspergillus flavus*. *Annual review of phytopathology*, 49, 107-133. <https://doi.org/10.1146/annurev-phyto-072910-095221>
- Bhatnagar-Mathur, P., Sunkara, S., Bhatnagar-Panwar, M., Waliyar, F., & Sharma, K. K. (2015). Biotechnological advances for combating *Aspergillus flavus* and aflatoxin contamination in crops. *Plant science*, 234, 119-132. <https://doi.org/10.1016/j.plantsci.2015.02.009>
- Bhattacharya, A., Tejero, R., & Montelione, G. T. (2007). Evaluating protein structures determined by structural genomics consortia. *Proteins: Structure, Function, and Bioinformatics*, 66(4), 778-795. <https://doi.org/10.1002/prot.21165>
- Blackburn, G. M. (Ed.). (2006). *Nucleic acids in chemistry and biology*. Royal Society of Chemistry.
- Breda, A., Valadares, N. F., de Souza, O. N., & Garratt, R. C. (2007). Protein structure, modelling and applications. In *Bioinformatics in tropical disease research: a practical and case-study approach [Internet]*. National Center for Biotechnology Information (US). <https://www.ncbi.nlm.nih.gov/books/NBK6824/>
- Benkert, P., Tosatto, S. C., & Schomburg, D. (2008). QMEAN: A comprehensive scoringfunction for model quality assessment. *Proteins: Structure, Function, and Bioinformatics*, 71(1), 261-277. <https://doi.org/10.1002/prot.21715>
- Benkert, P., Biasini, M., & Schwede, T. (2011). Toward the estimation of the absolute quality of individual protein structure models. *Bioinformatics*, 27(3), 343-350. <https://doi.org/10.1093/bioinformatics/btq662>
- Bordoli, L., & Schwede, T. (2012). Automated protein structure modeling with SWISS-MODEL Workspace and the Protein Model Portal. *Homology Modeling: Methods and Protocols*, 107-136. https://doi.org/10.1007/978-1-61779-588-6_5
- Bose, P., Gowrie, S. U., & Chaturdevi, G. (2019). Optimization of Culture Conditions for Growth and Production of Bioactive Metabolites by Endophytic Fungus-*Aspergillus tamarii*. *International Journal of Pharmacy and Biological Sciences*, 9(2), 469-478. <https://doi.org/10.21276/ijpbs.2019.9.2.60>
- Balina, A., Kebede, A., & Tamiru, Y. (2018). Review on aflatoxin and its impacts on livestock. *Journal of Dairy and Veterinary Sciences*, 6(2), e555685. <https://doi.org/10.19080/JDVS.2018.06.555685>
- Bashir, S., Janiad, S., Zeeshan, A., Farah, N., Mehboob, S., Butt, H. I., & Qari, S. H. (2022). In Silico Analysis and Function Prediction of Sr22 Gene Product as Stem Rust Resistant Protein. *Annals of the Romanian Society for Cell Biology*, 26(01), 258-262. <http://annalsofrscb.ro>
- Cram, D. J., & Cram, J. M. (1971). Cyclophane chemistry: bent and battered benzene rings. *Accounts of Chemical Research*, 4(6), 204-213. <https://doi.org/10.1021/ar50042a003>

- Corley, M., Burns, M. C., & Yeo, G. W. (2020). How RNA-binding proteins interact with RNA: molecules and mechanisms. *Molecular cell*, 78(1), 9-29. <https://doi.org/10.1016/j.molcel.2020.03.011>
- de la Cruz, J., Karbstein, K., & Woolford Jr, J. L. (2015). Functions of ribosomal proteins in assembly of eukaryotic ribosomes in vivo. *Annual review of biochemistry*, 84, 93-129. <https://doi.org/10.1146/annurev-biochem-060614-033917>
- Dhanamjayulu, P., Boga, R. B., & Mehta, A. (2019). Inhibition of aflatoxin B1 biosynthesis and down regulation of aflR and aflB genes in presence of benzimidazole derivatives without impairing the growth of *Aspergillus flavus*. *Toxicon*, 170, 60-67. <https://doi.org/10.1016/j.toxicon.2019.09.018>
- Damodaran, S., & Parkin, K. L. (2017). Amino acids, peptides, and proteins. In *Fennema's food chemistry* (pp. 235-356). CRC Press.
- Dreyfuss, G., Kim, V. N., & Kataoka, N. (2002). Messenger-RNA-binding proteins and the messages they carry. *Nature reviews Molecular cell biology*, 3(3), 195-205. <https://doi.org/10.1038/nrm760>
- Dutta, B., Banerjee, A., Chakraborty, P., & Bandopadhyay, R. (2018). In silico studies on bacterial xylanase enzyme: Structural and functional insight. *Journal of Genetic Engineering and Biotechnology*, 16(2), 749-756. <https://doi.org/10.1016/j.jgeb.2018.05.003>
- Elliott, D., & Ladomery, M. (2017). *Molecular biology of RNA*. Oxford University Press.
- Frank, J., & Spahn, C. M. (2006). The ribosome and the mechanism of protein synthesis. *Reports on Progress in Physics*, 69(5), 1383. <https://doi.org/10.1088/0034-4885/69/5/R03>
- Fouad, A. M., Ruan, D., El-Senousey, H. K., Chen, W., Jiang, S., & Zheng, C. (2019). Harmful effects and control strategies of aflatoxin b1 produced by *Aspergillus flavus* and *Aspergillus parasiticus* strains on poultry. *Toxins*, 11(3), 176. <https://doi.org/10.3390/toxins11030176>
- Franceschini, A., Szklarczyk, D., Frankild, S., Kuhn, M., Simonovic, M., Roth, A., & Jensen, L. J. (2012). STRING v9. 1: protein-protein interaction networks, with increased coverage and integration. *Nucleic acids research*, 41(D1), D808-D815. <https://doi.org/10.1093/nar/gks1094>
- Glisovic, T., Bachorik, J. L., Yong, J., & Dreyfuss, G. (2008). RNA-binding proteins and post-transcriptional gene regulation. *FEBS letters*, 582(14), 1977-1986. <https://doi.org/10.1016/j.febslet.2008.03.004>
- Gourama, H., & Bullerman, L. B. (1995). *Aspergillus flavus* and *Aspergillus parasiticus*: Aflatoxigenic fungi of concern in foods and feeds: A review. *Journal of Food protection*, 58(12), 1395-1404. <https://doi.org/10.4315/0362-028X-58.12.1395>
- Gasteiger, E., Gattiker, A., Hoogland, C., Ivanyi, I., Appel, R. D., & Bairoch, A. (2003). ExPASy: the proteomics server for in-depth protein knowledge and analysis. *Nucleic acids research*, 31(13), 3784-3788. <https://doi.org/10.1093/nar/gkg563>
- Hou, T., Sana, S. S., Li, H., Xing, Y., Nanda, A., Netala, V. R., & Zhang, Z. (2022). Essential oils and its antibacterial, antifungal and anti-oxidant activity applications: A review. *Food Bioscience*, 47, 101716. <https://doi.org/10.1016/j.fbio.2022.101716>
- Kirubakaran, P., Karthikeyan, M., Singh, K. D., Nagamani, S., & Premkumar, K. (2013). In silico structural and functional analysis of the human TOPK protein by structure modeling and molecular dynamics studies. *Journal of molecular modeling*, 19, 407-419. <https://doi.org/10.1007/s00894-012-1566-1>

- Klich, M. A. (2007). *Aspergillus flavus*: the major producer of aflatoxin. *Molecular plant pathology*, 8(6), 713-722. <https://doi.org/10.1111/j.1364-3703.2007.00436.x>
- Laskowski, R. A., Furnham, N., & Thornton, J. M. (2013). The Ramachandran plot and protein structure validation. In *Biomolecular forms and functions: a celebration of 50 years of the ramachandran map* (pp. 62-75). https://doi.org/10.1142/9789814449144_0005
- Lake, J. A. (1981). The ribosome. *Scientific American*, 245(2), 84-97. <https://doi.org/10.1038/scientificamerican0881-84>
- Lovell, S. C., Davis, I. W., Arendall III, W. B., De Bakker, P. I., Word, J. M., Prisant, M. G., ... & Richardson, D. C. (2003). Structure validation by C α geometry: ϕ , ψ and C β deviation. *Proteins: Structure, Function, and Bioinformatics*, 50(3), 437-450. <https://doi.org/10.1002/prot.10286>
- Lobo, I. (2008). Basic local alignment search tool (BLAST). *Nature Education*, 1(1).
- Nguyen, H. N., Chaves-Lopez, C., Oliveira, R. C., Paparella, A., & Rodrigues, D. F. (2019). Cellular and metabolic approaches to investigate the effects of graphene and graphene oxide in the fungi *Aspergillus flavus* and *Aspergillus niger*. *Carbon*, 143, 419-429. <https://doi.org/10.1016/j.carbon.2018.10.099>
- Ogodo, A. C., & Ugbogu, O. C. (2016). Public health significance of aflatoxin in food industry—A review. *European Journal of Clinical and Biomedical Sciences*, 2(5), 51-58. <https://doi.org/10.11648/j.ejcb.20160205.14>
- Rozario, L. T., Sharker, T., & Nila, T. A. (2021). In silico analysis of deleterious SNPs of human MTUS1 gene and their impacts on subsequent protein structure and function. *PLoS one*, 16(6), e0252932. <https://doi.org/10.1371/journal.pone.0252932>
- Ramirez, M. L., Cendoya, E., Nichea, M. J., Zchetti, V. G. L., & Chulze, S. N. (2018). Impact of toxigenic fungi and mycotoxins in chickpea: A review. *Current Opinion in Food Science*, 23, 32-37. <https://doi.org/10.1016/j.cofs.2018.05.003>
- Szklarczyk, D., Franceschini, A., Kuhn, M., Simonovic, M., Roth, A., Minguez, P., ... & Mering, C. V. (2010). The STRING database in 2011: functional interaction networks of proteins, globally integrated and scored. *Nucleic acids research*, 39(suppl_1), D561-D568. <https://doi.org/10.1093/nar/gkq973>
- Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., ... & von Mering, C. (2023). The STRING database in 2023: protein–protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic acids research*, 51(D1), D638-D646. <https://doi.org/10.1093/nar/gkac1000>
- Tam, E. W., Chen, J. H., Lau, E. C., Ngan, A. H., Fung, K. S., Lee, K. C., ... & Woo, P. C. (2014). Misidentification of *Aspergillus nomius* and *Aspergillus tamaris* as *Aspergillus flavus*: characterization by internal transcribed spacer, β -tubulin, and calmodulin gene sequencing, metabolic fingerprinting, and matrix-assisted laser desorption ionization–time of flight mass spectrometry. *Journal of Clinical Microbiology*, 52(4), 1153-1160. <https://doi.org/10.1128/JCM.03258-13>
- Von Hippel, P. H., Bear, D. G., Morgan, W. D., & McSwiggen, J. A. (1984). Protein-nucleic acid interactions in transcription: a molecular analysis. *Annual Review of Biochemistry*, 53(1), 389-446. <https://doi.org/10.1146/annurev.bi.53.070184.002133>
- Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., ... & Schwede, T. (2018). SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic acids research*, 46(W1), W296-W303. <https://doi.org/10.1093/nar/gky427>

- Yu, J., Cleveland, T. E., Nierman, W. C., & Bennett, J. W. (2005). *Aspergillus flavus* genomics: gateway to human and animal health, food safety, and crop resistance to diseases. *Revista iberoamericana de micología*, 22(4), 194-202. [https://doi.org/10.1016/S1130-1406\(05\)70043-7](https://doi.org/10.1016/S1130-1406(05)70043-7)
- Zain, M. E. (2011). Impact of mycotoxins on humans and animals. *Journal of Saudi chemical society*, 15(2), 129-144. <https://doi.org/10.1016/j.jscs.2010.06.006>

Second revision: Minor revisions (05-12-2023)

**In Silico Analysis of *Aspergillus flavus* Fungal Proteins:
Structural and Functional Insights Using ITS Primers**

**Mariana Mariana, Saipul Abbas*, Salamiah Salamiah, Samharinto Samharinto, Yusriadi Marsuni,
Muslimin Sepe,
Muhammad Indar Pramudi, Ismed Setya Budi, Dewi Fitriyanti, Elly Liestiany, Lyswiana
Aphrodyanti**

Department of Plant Protection, Faculty of Agriculture, Lambung Mangkurat University. Jl.Jend. A.
Yani Km.36 Banjarbaru. South Kalimantan 70714 Indonesia

*Corresponding author: Email: saipul.abbas@ulm.ac.id

ABSTRACT

Aspergillus flavus is a type of fungus that can contaminate and damage various types of food and agricultural products. To better understand the structure and function of proteins, in silico research was conducted using Internal Transcribed Spacer (ITS) primers and SWISS Model protein prediction tools. The objective of this study was to identify fungal species using the BLAST method and to analyze the structure and function of proteins from *Aspergillus spp.* The methods used included BLAST for species identification, Web Expaty to translate DNA sequences into proteins, SWISS Models to model protein structures, SAVES to validate protein structures, and STRING to analyze the function of proteins. The results of the BLAST analysis showed that the identified fungal species were *Aspergillus flavus*, *Aspergillus tamarii*, and *Aspergillus nomius*. Furthermore, the results of translating DNA sequences into proteins using Web Expaty showed that there were three open reading frames with the highest residual values of 119 and 83, while the lowest residual value was 4. Only two of these frames met the protein criteria. Moreover, the results of protein structure modeling using the SWISS Model method produced a fairly accurate *Aspergillus spp* protein structure model with a validation value of protein structure using ERRAT (SAVES V6.0) of 100%. Additionally, the results of protein function analysis using STRING showed that the *Aspergillus spp* protein has a function in producing enzymes that play a role in the metabolic process of cells.

Keywords: *Aspergillus flavus*, in silico, ITS primer, protein structure, protein function, SWISS Model

INTRODUCTION

Aspergillus spp fungus is a type of fungus that often causes contamination and damage to various agricultural and food products (Gourama & Bullerman, 1995), such as grains, nuts, and processed products

(Ayofemi Olalekan Adeyeye, 2020). One of the fungi *Aspergillus* spp that can contaminate feed ingredients is *the species Aspergillus flavus* which can cause aflatoxin pollution (Klich, 2007), which is a harmful substance that can affect human health and livestock (Balina et al, 2018). *Aspergillus flavus* is a group of fungi often found in agricultural environments, especially in food crops such as corn, rice, and legumes (Amaike & Keller, 2011). Contamination by *Aspergillus spp.* can cause damage to agricultural products and foodstuffs and can cause health problems in humans and farm animals that consume them (Yu et al, 2005). Therefore, research on *Aspergillus spp.* is very important in agriculture to prevent and control contamination of *Aspergillus spp.* on agricultural products and foodstuffs.

Previous studies have highlighted the negative impact of *Aspergillus* spp. on agricultural products and food items. Ramirez et al. (2018) noted that this type of fungus can cause damage to grains, legumes, and processed products. Similarly, Balina et al. (2018) indicated that *A. flavus* can be a major cause of aflatoxin contamination in feed materials, posing risks to livestock and humans who consume them. The significance of research on *A. flavus* is further underscored by the findings of Bhatnagar-Mathuret al. (2015), demonstrating the frequent presence of *A. flavus* in agricultural environments, with a high tendency to infect staple food crops. Therefore, contamination risk from *A. flavus* not only affects the quality of agricultural products and food items but also poses a serious health risk to consumers. In silico research has become a useful tool for studying the structure and function of proteins (Dutta et al, 2018), as it allows for quick and efficient hypothesis testing and analysis (Kirubakaran et al, 2013). In this study, we used an in-silico technique to predict the structure and function of the *A. flavus* protein by combining the primary use of the Internal Transcribed Spacer (ITS) and the SWISS Model protein prediction tool. This in silico approach is essential to understanding this fungus and is beneficial to the food and agriculture industries.

The formulation of the problem of this study was to analyze the BLAST results of these *Aspergillus spp* sequences, examine the protein structure related to contamination, evaluate the validity and confidence of the protein structure of *Aspergillus spp.* which was found, as well as examining the function of the protein *Aspergillus spp.* associated with contamination of corn feed and other foodstuffs, as well as to understand the structure and function of the *Aspergillus spp* protein associated with the contamination aforementioned. Therefore, this research focused on the analysis of *Aspergillus spp.* sequence, protein structure analysis, and protein function analysis, with the hope of providing new insights and solutions to the problem of contamination in food.

This study aims to analyze the species *Aspergillus spp.* which often contaminates corn feed and other foodstuffs. This study is expected to provide information about the species *Aspergillus spp.* which contaminates corn feed and other foodstuffs, as well as the structure and function of the protein *Aspergillus spp.* The results of this study are expected to contribute to the development of methods for detecting and preventing contamination of *Aspergillus spp.* in corn feed and other foodstuffs.

MATERIALS AND METHODS

Identification of fungus species

This step involved the use of an Internal Transcribed Spacer (ITS) Primer to **verify** fungus species. This stage of research was carried out by taking DNA samples from fungi and conducting DNA amplification using primary ITS. DNA amplification was performed using the Polymerase Chain Reaction (PCR) technique. This step aimed to verify the species of fungus analyzed. In the PCR technique, DNA samples were placed together with ITS primers (ITS1 5'-TCCGTAGGTGAACCTGCGG-3', and ITS4 5'-TCCTCCGCTTATTGATATGC-3') and polymerase enzymes, and a heating and cooling process was carried out to start and end the reaction. The result of the amplification process was more DNA sequences associated with ITS. After successful DNA amplification, the obtained DNA sequence was analyzed to verify the fungus species. Sequencing analysis was carried out using bioinformatics software such as BLAST (*Basic Local Alignment Search Tool*) (Lobo, 2008).

Translate DNA Sequencing into Proteins

After the fungus DNA sequence was successfully analyzed, the DNA sequence was then translated into proteins. This step was done using a tool such as the ExPasy Translate Tool (Bashir, 2022). This tool converted the DNA sequence into a protein sequence based on the genetic code (Ayaz et al, 2020). The results of the translation showed the amino acid sequence contained in the fungus protein.

Analysis of protein structure

This stage used the SWISS Model method to predict the protein structure of the fungus *Aspergillus* spp. At this stage, the input of protein sequencing data from ExPasy tools was carried out, and then protein modeling was made using the SWISS Model (Waterhouse et al, 2018). Analysis using the SWISS Model provided information regarding protein domains, protein motifs, and other information related to the structure and function of proteins (Bordoli & Schwede, 2012). The results of the analysis of the structure and function of the protein were re-examined to ensure the validity of the results. Validation was done by comparing the results with a trusted protein database such as UniProt.

Protein Function Analysis

The research method of protein function analysis with STRING involved the use of a database of protein interactions and the analysis of protein networks (Franceschini et al, 2012). In this study, *Aspergillus flavus* protein data were used in the STRING database. Protein data were inputted into a STRING database containing information about protein interactions formed based on data from literature databases, scientific publications, and experimental studies (Szkarczyk et al, 2010). Analysis of protein

interactions formed using algorithms available on the STRING database. This algorithm generated information about the interaction of formed proteins, the forces of interaction, and their relationship with biological functions. Interpretation of the results of protein interaction analysis was done to obtain information about the biological function of the protein being analysed and how the protein interaction was related to the biological function found. This method could be used to analyse the function of proteins and understand how proteins interacted with each other in a network of proteins (Szklarczyk et al, 2023). Using this method, we could gain a better understanding of how proteins were involved in biological processes and their possible effects on health and disease.

RESULTS AND DISCUSSION

Identification of fungus species by BLAST Method

The results of the fungal species identification using BLAST revealed a similarity of 99.61% with *Aspergillus tamarii*, 98.94% with *Aspergillus flavus*, and 96.33% with *Aspergillus nomius*. These findings suggested that the sample likely belonged to the *Aspergillus* group, with a higher similarity to *Aspergillus tamarii* compared to the other two species. *Aspergillus tamarii* is recognized for its capability to produce various bioactive compounds (Bose et al, 2019), including phenol, flavonoid and IAA and its antifungal activity, which have potential applications in the Agriculture, and food industries. However, it is important to note that *Aspergillus tamarii* can also cause infections in humans and animals (Tam et al, 2014). Similarly, *Aspergillus flavus* is known for its production of toxic compounds, such as aflatoxin, which can be harmful to human and animal health (Fouad et al, 2019). Therefore, proper handling measures are necessary to prevent the spread of this species. Moreover, *Aspergillus nomius*, another identified species in the sample, can produce aflatoxin and the potential to become pathogenic in humans and animals (Zain, 2011). Consequently, accurate identification of the fungus species is essential to determine appropriate control measures. In this case, the identification results indicating the highest similarity to *Aspergillus tamarii* and *Aspergillus flavus* can serve as references for further research on the bioactive compounds produced by these species. Nevertheless, further examinations are required to confirm the presence of specific fungus species in the sample and establish appropriate control measures to prevent its spread.

select all 10 sequences selected		GenBank	Graphics	Distance tree of results	MSA Viewer			
Description	Scientific Name	Max Score	Total Score	Query Cover	E value	Per. Ident	Acc. Len	Accession
Aspergillus tamarii strain M24 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8S...	Aspergillus tamarii	1419	2347	97%	0.0	99.61%	889	ON819577.1
Aspergillus flavus strain Bp5 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1 and 5.8S riboso...	Aspergillus flavus	1343	2094	94%	0.0	98.94%	939	KF221065.1
Aspergillus tamarii strain BPFM17 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1,...	Aspergillus tamarii	1321	2151	99%	0.0	96.86%	897	MF359724.1
Aspergillus tamarii strain MF24 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8...	Aspergillus tamarii	1310	1952	98%	0.0	96.83%	791	MF359732.1
Aspergillus sp. AQGWD 17 internal transcribed spacer 1, partial sequence; 5.8S ribosomal RNA gene and internal tran...	Aspergillus sp. A...	1297	2462	100%	0.0	96.37%	1303	KP721583.1
Aspergillus sp. isolate ACSIKS_2100168 small subunit ribosomal RNA gene, partial sequence; internal transcribed spa...	Aspergillus sp.	1291	2430	100%	0.0	96.25%	1187	MN597042.1
Aspergillus sp. isolate SS_12 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8...	Aspergillus sp.	1286	2440	100%	0.0	96.12%	1303	MT497446.1
Aspergillus flavus isolate 20 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1 and 5.8...	Aspergillus flavus	1273	1946	91%	0.0	98.18%	715	MK120614.1
Aspergillus nomius strain Tur4 18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1 and 5.8S ribos...	Aspergillus nomiae	1240	1891	95%	0.0	96.33%	842	KF221090.1
Aspergillus flavus strain Beca_43 small subunit ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5...	Aspergillus flavus	1234	2105	99%	0.0	96.80%	849	KY234269.1

Figure 1: The results of the fungus species identification using BLAST Method (GenBank)

Translate DNA Sequencing into Proteins

The results obtained from the translation of *Aspergillus* spp DNA sequences into proteins using the ExpASy web revealed the presence of three open reading frames (ORFs), indicated in red, with residual values of 119 and 83 as the highest, and 4 as the lowest. Proteins are typically categorized if they consist of more than 50 residues (Damodaran & Parkin, 2017). In this case, only two out of the three ORFs met the criteria to be considered as proteins. The identification of an ORF qualifying as a protein may indicate the presence of a gene within the *Aspergillus* DNA sequence. Translating DNA sequences into proteins holds significant importance in molecular biology research as it allows for a better understanding of protein functions within a specific organism (Gasteiger et al, 2003). Consequently, these findings serve as a foundation for future investigations concerning the *Aspergillus* gene and its potential applications in various fields, such as agriculture and the food industry. Nevertheless, it is crucial to note that these results should undergo further verification using other techniques, such as protein structure analysis, to ensure the functional integrity of the resulting proteins. Additionally, the authenticity and suitability of the *Aspergillus* DNA sequence utilized in this study should be validated about the specific *Aspergillus* species under investigation.

<p>5'3' Frame 1</p> <p>S LN-LRYNQ LRLH-I RQSSWCLRRARARG-E PPAAMNGGPAEATKVQ-T RVGGWAR-E PYTR--S FRR-T CGRI ITECRVPSEPNLPFVFTVP-L LRRARHSWPPGALS PGPAP AGDTNSV-S SEV-VDCIAIS-N FQQWISWFRHR-RTQRNAITSVNCRIP-I IESLNAHCAPWYSGGHACPSVIAAHQARLVCWVVVPSGGTGPKGSGGTASDPRAYGALSPA L-ARPALAERKSIFFQVDLGSGRDTR-T-A YQ-GGG</p>
<p>5'3' Frame 2</p> <p>V L TDCDTINSDFTRS D R V R G V S G G R G P G A E S P R R P -M A G P P Q L R Y S K H G W E V G L A R N P T L G N D P S V G E P A E G S L P S V G F L A S P T S H P C L L Y L S C F G G P A I H G R R G L S A P G P R P P E T P R T L S D L V K S E L I V S Q S V K T F N N G S L G S I D E E R S E M R -L V -I A E P R E S S L -T H I A P P G I P G G M P V R A S L L P I K H G L C V G S S S P L R G R A P K A A A P R P I L E R M G L C H P L C R P G R R L P N A N Q S F S R L T S D Q V G I P A E L K H I N K A E E</p>
<p>5'3' Frame 3</p> <p>S -L T A I Q S T Q T S L D Q T E F V V S P A G A G P L R A P G G H E W R A R R S N -G T V N T G G R L G S L G T L H S V M I L P -V N L R K D H Y R V -G S -R A Q P P T R V Y C T L V A S A G P P F M A A G G S Q P R A R A R R R H H E L C L I -S L S -L Y R N Q L K L S T M D L L V P A S M K N A A K C D N -C E L Q N S V N H R V F E R T L R P L V F R G A C L S E R H C C P S S T A C V L G R R R L P S G G D G P Q R R H R V R S S V W G F V T R S V G P A G A C R T Q I N L P F P -P R I R -G Y P L N L S I S I R R R</p>

Figure 2: The results of *Aspergillus* spp DNA sequences into proteins using the ExpASy web

Table 1: The results of Open Reading Frames (ORFs) with residual values from 4 to 119

Open Reading Frame	Residue	Types of Molecules	Code Peptides/Proteins
--------------------	---------	--------------------	------------------------

MNGGPAEATKVQ	12	Peptide	A1
MAGPPKQLRYSKHGWEVGLARNPTLGND PSVGEPAEGLPSVGFLASPTSHPCLLYLSC FGGPAIHGRRGLSAPGPRPPETPRTLSDLV KSELIVSQSVKTFNNGSLGSGIDEERSEMR	119	Protein	A2
MPVRASLLPIKHGLCVGSSSPLRGGRAPKA AAAPRPILERMGLCHPLCRPGRRLPNANQS FSRLTSDQVGIPAELKHINKAEE	83	Protein	A3
MILP	4	Peptide	A4
MAY	4	Peptide	A5
MAAGGSQPRARARRRHHELCLI	22	Peptide	A6
MDLLVPASMKNAAKCDN	17	Peptide	A7

Protein Structure Modeling with the SWISS Model method

The results of modeling the protein structure of *Aspergillus spp* using the SWISS Model method show the presence of a 3D structure of proteins formed (Bordoli & Schwede, 2012). Protein structure modeling is an important method in molecular biology research because it can provide information about the shape and function of proteins in a particular organism (Breda et al, 2007). The results of modeling the protein structure of *Aspergillus spp* can be used as a basis for conducting further analysis of protein function, including interactions with other molecules, enzymatic activity, and possible drug development. In addition, these results can also help in understanding the relationship between the structure of proteins and their biological activity. However, keep in mind that the results of modeling protein structures using computational methods such as the SWISS Model still have limitations, especially in terms of accuracy. Therefore, these results still need to be verified using other techniques, such as X-ray crystallography or core magnetic resonance spectroscopy (NMR), to ensure the accuracy of the resulting protein structure. In addition, modeling the protein structure of *Aspergillus spp* can also provide information about the evolutionary relationship of proteins between different *Aspergillus* species. Modeling the structure of the *Aspergillus spp* protein can help understand the differences and similarities between different *Aspergillus* proteins so that it can be used as a basis for specific drug and therapeutic development.




Figure 3: Protein Structure Prediction from Internal Transcribed Spacer (ITS) primer of *Aspergillus flavus* fungus using Protein Prediction from SWISS Model. (Template: A2: 3lb6.1.A A3: 5t7v.1.K)

The results of validation of the *Aspergillus spp* protein structure using SAVES V6.0 show that the resulting protein structure has a fairly good value in terms of accuracy and quality. One of the parameters used to evaluate the validation of protein structure is the Ramachandran plot, which shows the proportion of amino acids placed in the most favored regions within the protein conformation space (Laskowski et al, 2013). In the Ramachandran plot results obtained, the highest Residual value in most favored regions was 80% for the protein code A2, which indicates that most of the amino acids in the protein structure of *Aspergillus spp* were placed in the desired area (Lovell et al, 2003). In addition, the validation results using the ERRAT method also show the highest value of 100%, which indicates that the resulting protein structure has high accuracy (Rozario et al, 2021). The results of the validation of the protein structure of *Aspergillus spp* using SAVES V6.0 are important to ensure that the resulting protein structure is trustworthy and used as a basis for further analysis of protein function and drug development. Validation of protein structure can also help identify and correct parts in the protein structure that may be inaccurate, thereby improving the quality and accuracy of research results.


The QMEAN4 Score indicates the overall **quality** of the protein structure, which is calculated based on geometric values, non-covalent interactions, and consistency of the structure (Benkert et al, 2008). The higher the QMEAN 4 score, the better the quality of the resulting protein structure (Benkert et al, 2011). In this case, the QMEAN4 Score A2 value of -1.72 versus A3 of -3.27 can be interpreted as a low value, which indicates that the resulting *Aspergillus spp* protein structure does not have optimal quality. Meanwhile, the Z Score is a parameter used to compare the quality of the resulting protein structure with other similar protein structures, which are calculated based on geometric values and structural consistency. The higher the Z Score value, the better the quality of the resulting protein structure compared to other similar protein structures (Bhattacharya et al, 2007). In this case, a Z Score of -0.16 indicates that the resulting *Aspergillus spp* protein structure is of relatively good quality compared to other similar protein structures. However, keep in mind that as with the QMEAN 4 Score value, the Z Score value also has limitations and possible errors, so this result still needs to be verified using other protein structure validation techniques.

Table 2: Protein Structure Validation using SAVES V6.0

Protein Code	Ramachandran plot				ERRA T quality factor (%)	QMEAN 4 score	Z Score
	Residue in most favored regions (%)	Residue in additional allowed regions (%)	Residue in generously allowed regions (%)	Residue in disallowed regions (%)			
A2	80.0	20.0	0.0	0.0	100	-1.72	-0.16


A3 75.0 20.0 0.0 5.0 6.6666 -3.27 -3.32

Model A2



PROCHECK

Ramachandran Plot



Model A3




Figure 4: The results of validation of the *Aspergillus spp* protein structure using SAVES V6.0

Protein Function & Interaction of *Aspergillus Flavus*

Aspergillus flavus is a type of fungus known to cause damage to agricultural products such as grains and other foodstuffs (Gourama & Bullerman, 1995). This fungus can also produce some toxic substances such as aflatoxins, which are heterocyclic substances and have detrimental effects on human and animal health (Dhanamjayulu et al, 2019). Heterocyclics in aflatoxins have several main functions in the fungus *Aspergillus flavus*, including: (1) As a toxic compound: Aflatoxins have high toxic activity and can cause food poisoning in humans and animals, which can cause various health problems such as liver damage, cancer, and reproductive problems (Ogodo & Ugbogu, 2016). (2) As an antioxidant compound: Aflatoxin has antioxidant properties and can help the fungus *Aspergillus flavus* survive in a poor

environment (Abrar et al, 2013). (3) As an antifungal compound: Aflatoxin can help the fungus *Aspergillus flavus* resist attacks from other fungi that seek to interfere with the growth of the fungus *Aspergillus flavus* (Bhatnagar-Mathur et al, 2015). Overall, heterocyclic in aflatoxins has an important role in the biology of the fungus *Aspergillus flavus*, but it also has detrimental effects on human and animal health. Therefore, it is important to control the production of aflatoxins and ensure that the food products consumed do not contain this substance in harmful quantities.

Organic cyclic compound binding is the process of binding organic chemical compounds that have a benzene ring (Cram & Cram, 1971). In this case, such binding may take place in the fungus *Aspergillus flavus*. Organic chemical compounds that bind to the fungus *Aspergillus flavus* can affect the biological activity of the fungus (Nguyen et al, 2019). Some organic chemical compounds that can bind to the fungus *Aspergillus flavus* include growth regulators, metabolic regulators, and anti-fungal substances (Hou et al, 2022). The binding of organic chemical compounds to the fungus *Aspergillus flavus* can affect the growth and reproduction of fungi, modulate enzyme activity, affect the production of toxic substances such as aflatoxins, and affect the interaction of fungi with their environment. Overall, the binding function of organic chemical compounds to the fungus *Aspergillus flavus* is essential for understanding the biology of the fungus and how the fungus interacts with its environment. Further research on the binding of organic chemical compounds to the fungus *Aspergillus flavus* can help in controlling the production of aflatoxins and improving the quality of food products.

Nucleic acid binding protein has a very important function in the fungal cells of *Aspergillus flavus*. Nucleic acids are important molecules that carry genetic information and play a key role in various biological processes, such as DNA replication, RNA transcription, and protein synthesis (Blackburn, 2006). Nucleic acid binding proteins can bind to nucleic acid molecules, facilitate interaction with enzymes and transcription factors, and influence the biological activity of nucleic acid molecules (Von Hippel et al, 1984). In the fungus *Aspergillus flavus*, Nucleic acid binding proteins may play a role in regulating gene activity, affecting growth and reproduction, and modulating the fungus' response to the environment. Studies of Nucleic acid binding proteins in the fungus *Aspergillus flavus* can provide information about the mechanisms of gene regulation and how fungi react to their environment. Overall, the function of the Nucleic acid binding protein in the fungus *Aspergillus flavus* is essential for understanding the biology of the fungus and how the fungus interacts with its environment. More research on Nucleic acid binding proteins in the fungus *Aspergillus flavus* can help understand the growth and reproduction mechanisms of fungi, as well as how fungi react to their environment.

RNA binding is a process in which proteins bind to RNA molecules. RNA is an important molecule in cells that plays a key role in biological processes such as transcription, protein synthesis, and gene regulation (Elliott & Lodomery, 2017). RNA protein binding can bind to RNA molecules and facilitate interaction with enzymes and transcription factors (Corley et al, 2020). The function of RNA binding proteins can affect the biological activities of RNA molecules, such as modulating transcription and protein synthesis, affecting RNA stability, and facilitating gene control (Gilsovic et al, 2008). In the

fungus, *Aspergillus flavus*, protein RNA binding may play an important role in regulating gene activity, modulating responses to the environment, and facilitating growth and reproduction. Studies on protein RNA binding in the fungus *Aspergillus flavus* may help understand the mechanisms of gene regulation and how fungi react to their environment. Overall, the function of RNA binding proteins in the fungus *Aspergillus flavus* is critical to understanding the biology of the fungus and how the fungus interacts with its environment. More research on protein RNA binding in the fungus *Aspergillus flavus* may help in understanding the mechanisms of fungal growth and reproduction, as well as how the fungus reacts to its environment.

mRNA binding is a process in which proteins bind to mRNA molecules (Dreyfuss et al, 2002). mRNA is an RNA molecule that contains genetic information from DNA and is used as a template for protein synthesis (Alberts et al, 2002). mRNA protein binding can bind to mRNA molecules and facilitate interaction with enzymes and transcription factors (Corley et al, 2020). In the fungus *Aspergillus flavus*, mRNA protein binding may play an important role in regulating gene activity, modulating responses to the environment, and facilitating growth and reproduction. Studies on mRNA protein binding in the fungus *Aspergillus flavus* may help understand the mechanisms of gene regulation and how fungi react to their environment. Overall, the function of mRNA protein binding in the fungus *Aspergillus flavus* is critical to understanding the biology of the fungus and how the fungus interacts with its environment. More research on mRNA protein binding in the fungus *Aspergillus flavus* may help understand the mechanisms of fungal growth and reproduction, as well as how the fungus reacts to its environment.

Ribosomes are cellular structures that play a key role in the process of protein synthesis (Frank & Spahn, 2006). Ribosomes consist of two subunits, a large subunit and a small subunit, which serve to bind amino acids and form polypeptide chains (Lake, 1981). Proteins that are structural components of ribosomes can play an important role in ensuring the integrity and activity of ribosomes. The function of proteins as structural components of ribosomes can affect the ability of ribosomes to bind to mRNA and facilitate proper protein synthesis (de la Cruz et al, 2015). In the fungus *Aspergillus flavus*, proteins that are structural components of ribosomes may play an important role in ensuring the growth and reproduction of fungi. The study of proteins that are structural components of ribosomes in the fungus *Aspergillus flavus* can help understand the mechanism of protein synthesis and how the fungus reacts to its environment. Overall, the function of proteins as structural components of ribosomes in the fungus *Aspergillus flavus* is essential for understanding the biology of the fungus and how the fungus interacts with its environment. More research on the proteins that are structural components of ribosomes in the fungus *Aspergillus flavus* can help in understanding the mechanisms of fungal growth and reproduction, as well as how the fungus reacts to its environment.




Figure 5: Network of protein-protein interactions visualized with STRING (NCBI taxon-Id: 332952)

	B8MXN8	Serine/threonine protein kinase, putative
	B8N3J1	Small nucleolar ribonucleoprotein complex subunit Utp14, putative
	B8NT05	G-protein complex beta subunit CpcB
	B8N430	Nuclear mRNA splicing factor, putative
	B8N0Y2	G-patch domain protein (TFIP11), putative
	B8N058	PWI domain mRNA processing protein, putative
	B8N1X2	Pre-RNA splicing factor Srp2, putative
	B8ND38	CaaX prenyl proteinase Rce1
	B8MXA7	Small nuclear ribonucleoprotein E; Associated with the spliceosome snRNP U1, U2, U4/U6 and U5.
	B8MXX7	Casein kinase, putative; Belongs to the protein kinase superfamily.

Table 3: Molecular Function, Biological Process, Cellular Component, and Subcellular Localization Classification

Molecular Function	Biological Process	Cellular Component	Sucellular Localization
<ul style="list-style-type: none"> Heterocyclic compound binding 	<ul style="list-style-type: none"> mRNA splicing, via spliceosome RNA processing 	<ul style="list-style-type: none"> Spliceosomal complex 	<ul style="list-style-type: none"> U2 snRNP U2-type prespliceosome

• Organic cyclic compound binding	• Gene expression	• Ribonucleoprotein complex	• U2-type spliceosomal complex
• Nucleic acid binding	• Macromolecule metabolic process	• Nucleus	• Spliceosomal complex
• RNA binding	• Cellular nitrogen compound metabolic process	• Intracellular membrane-bounded organelle	• Ribonucleoprotein complex
• mRNA binding	• Nitrogen compound metabolic process		• Nucleus
• Structural constituent of ribosome	• Primary metabolic process		• Protein-containing complex
• rRNA binding	• Cellular metabolic process		• Intracellular membrane-bounded organelle
• snoRNA binding			• Intracellular
			• Cellular anatomical entity

Conclusion

In silico research was conducted to understand the structure and function of *proteins of Aspergillus spp.* The results of the BLAST analysis showed the presence of three species of fungi, namely *Aspergillus flavus*, *Aspergillus tamarii*, and *Aspergillus nomius*. Translating DNA sequences into proteins using Web Exspasy showed that only two proteins met the criteria. Protein structure modeling using SWISS Model resulted in a fairly accurate *Aspergillus spp* protein structure model with protein structure validation values using ERRAT (SAVES V6.0) of 100%. Analysis of protein function using STRING shows that *Aspergillus spp* protein has a function in producing enzymes that play a role in the metabolic process of cells. Thus, this study provides important information about the structure and function of the *protein Aspergillus spp*, which can be used to improve understanding of contamination and damage to food and agricultural products by this type of fungus.

Acknowledgement

A thank you to PT. Arutmin Indonesia Tambang Satui and Agreement/Contract Number: 39/UN8.1.23/PP/2022 through the Matching Fund program "Kedaireka" for financial support for this research activity, and all parties who have helped in the success of this research. All the help and support provided means a lot to us. Thanks.

Reference


- Ayaz, A., Zaman, W., Saqib, S., Ullah, F., & Mahmood, T. (2020). Phylogeny and diversity of Lamiaceae based on rps14 gene in Pakistan. *Genetika*, 52(2), 435-452. <https://doi.org/10.2298/GENSR2002435A>
- Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., & Walter, P. (2002). From DNA to RNA. In *Molecular Biology of the Cell. 4th edition*. Garland Science.
- Abrar, M., Anjum, F. M., Butt, M. S., Pasha, I., Randhawa, M. A., Saeed, F., & Waqas, K. (2013). Aflatoxins: biosynthesis, occurrence, toxicity, and remedies. *Critical reviews in food science and nutrition*, 53(8), 862-874. <https://doi.org/10.1080/10408398.2011.563154>
- Ayofemi Olalekan Adeyeye, S. (2020). Aflatoxigenic fungi and mycotoxins in food: a review. *Critical reviews in food science and nutrition*, 60(5), 709-721. <https://doi.org/10.1080/10408398.2018.1548429>
- Amaike, S., & Keller, N. P. (2011). *Aspergillus flavus*. *Annual review of phytopathology*, 49, 107-133. <https://doi.org/10.1146/annurev-phyto-072910-095221>
- Bhatnagar-Mathur, P., Sunkara, S., Bhatnagar-Panwar, M., Waliyar, F., & Sharma, K. K. (2015). Biotechnological advances for combating *Aspergillus flavus* and aflatoxin contamination in crops. *Plant science*, 234, 119-132. <https://doi.org/10.1016/j.plantsci.2015.02.009>
- Bhattacharya, A., Tejero, R., & Montelione, G. T. (2007). Evaluating protein structures determined by structural genomics consortia. *Proteins: Structure, Function, and Bioinformatics*, 66(4), 778-795. <https://doi.org/10.1002/prot.21165>
- Blackburn, G. M. (Ed.). (2006). *Nucleic acids in chemistry and biology*. Royal Society of Chemistry.
- Breda, A., Valadares, N. F., de Souza, O. N., & Garratt, R. C. (2007). Protein structure, modelling and applications. In *Bioinformatics in tropical disease research: a practical and case-study approach [Internet]*. National Center for Biotechnology Information (US). <https://www.ncbi.nlm.nih.gov/books/NBK6824/>
- Benkert, P., Tosatto, S. C., & Schomburg, D. (2008). QMEAN: A comprehensive scoring function for model quality assessment. *Proteins: Structure, Function, and Bioinformatics*, 71(1), 261-277. <https://doi.org/10.1002/prot.21715>
- Benkert, P., Biasini, M., & Schwede, T. (2011). Toward the estimation of the absolute quality of individual protein structure models. *Bioinformatics*, 27(3), 343-350. <https://doi.org/10.1093/bioinformatics/btq662>
- Bordoli, L., & Schwede, T. (2012). Automated protein structure modeling with SWISS-MODEL Workspace and the Protein Model Portal. *Homology Modeling: Methods and Protocols*, 107-136. https://doi.org/10.1007/978-1-61779-588-6_5
- Bose, P., Gowrie, S. U., & Chaturdevi, G. (2019). Optimization of Culture Conditions for Growth and Production of Bioactive Metabolites by Endophytic Fungus-*Aspergillus tamarii*. *International Journal of Pharmacy and Biological Sciences*, 9(2), 469-478. <https://doi.org/10.21276/ijpbs.2019.9.2.60>

- Balina, A., Kebede, A., & Tamiru, Y. (2018). Review on aflatoxin and its impacts on livestock. *Journal of Dairy and Veterinary Sciences*, 6(2), e555685. <https://doi.org/10.19080/JDVS.2018.06.555685>
- Bashir, S., Janiad, S., Zeeshan, A., Farah, N., Mehboob, S., Butt, H. I., & Qari, S. H. (2022). In Silico Analysis and Function Prediction of Sr22 Gene Product as Stem Rust Resistant Protein. *Annals of the Romanian Society for Cell Biology*, 26(01), 258-262. <http://annalsofrscb.ro>
- Cram, D. J., & Cram, J. M. (1971). Cyclophane chemistry: bent and battered benzene rings. *Accounts of Chemical Research*, 4(6), 204-213. <https://doi.org/10.1021/ar50042a003>
- Corley, M., Burns, M. C., & Yeo, G. W. (2020). How RNA-binding proteins interact with RNA: molecules and mechanisms. *Molecular cell*, 78(1), 9-29. <https://doi.org/10.1016/j.molcel.2020.03.011>
- de la Cruz, J., Karbstein, K., & Woolford Jr, J. L. (2015). Functions of ribosomal proteins in assembly of eukaryotic ribosomes in vivo. *Annual review of biochemistry*, 84, 93-129. <https://doi.org/10.1146/annurev-biochem-060614-033917>
- Dhanamjayulu, P., Boga, R. B., & Mehta, A. (2019). Inhibition of aflatoxin B1 biosynthesis and down regulation of aflR and aflB genes in presence of benzimidazole derivatives without impairing the growth of *Aspergillus flavus*. *Toxicon*, 170, 60-67. <https://doi.org/10.1016/j.toxicon.2019.09.018>
- Damodaran, S., & Parkin, K. L. (2017). Amino acids, peptides, and proteins. In *Fennema's food chemistry* (pp. 235-356). CRC Press.
- Dreyfuss, G., Kim, V. N., & Kataoka, N. (2002). Messenger-RNA-binding proteins and the messages they carry. *Nature reviews Molecular cell biology*, 3(3), 195-205. <https://doi.org/10.1038/nrm760>
- Dutta, B., Banerjee, A., Chakraborty, P., & Bandopadhyay, R. (2018). In silico studies on bacterial xylanase enzyme: Structural and functional insight. *Journal of Genetic Engineering and Biotechnology*, 16(2), 749-756. <https://doi.org/10.1016/j.jgeb.2018.05.003>
- Elliott, D., & Lodomery, M. (2017). *Molecular biology of RNA*. Oxford University Press.
- Frank, J., & Spahn, C. M. (2006). The ribosome and the mechanism of protein synthesis. *Reports on Progress in Physics*, 69(5), 1383. <https://doi.org/10.1088/0034-4885/69/5/R03>
- Fouad, A. M., Ruan, D., El-Senousey, H. K., Chen, W., Jiang, S., & Zheng, C. (2019). Harmful effects and control strategies of aflatoxin b1 produced by *Aspergillus flavus* and *Aspergillus parasiticus* strains on poultry. *Toxins*, 11(3), 176. <https://doi.org/10.3390/toxins11030176>
- Franceschini, A., Szklarczyk, D., Frankild, S., Kuhn, M., Simonovic, M., Roth, A., ... & Jensen, L. J. (2012). STRING v9. 1: protein-protein interaction networks, with increased coverage and integration. *Nucleic acids research*, 41(D1), D808-D815. <https://doi.org/10.1093/nar/gks1094>
- Glisovic, T., Bachorik, J. L., Yong, J., & Dreyfuss, G. (2008). RNA-binding proteins and post-transcriptional gene regulation. *FEBS letters*, 582(14), 1977-1986. <https://doi.org/10.1016/j.febslet.2008.03.004>
- Gourama, H., & Bullerman, L. B. (1995). *Aspergillus flavus* and *Aspergillus parasiticus*: Aflatoxigenic fungi of concern in foods and feeds: A review. *Journal of Food protection*, 58(12), 1395-1404. <https://doi.org/10.4315/0362-028X-58.12.1395>

- Gasteiger, E., Gattiker, A., Hoogland, C., Ivanyi, I., Appel, R. D., & Bairoch, A. (2003). ExPASy: the proteomics server for in-depth protein knowledge and analysis. *Nucleic acids research*, 31(13), 3784-3788. <https://doi.org/10.1093/nar/gkg563>
- Hou, T., Sana, S. S., Li, H., Xing, Y., Nanda, A., Netala, V. R., & Zhang, Z. (2022). Essential oils and its antibacterial, antifungal and anti-oxidant activity applications: A review. *Food Bioscience*, 47, 101716. <https://doi.org/10.1016/j.fbio.2022.101716>
- Kirubakaran, P., Karthikeyan, M., Singh, K. D., Nagamani, S., & Premkumar, K. (2013). In silico structural and functional analysis of the human TOPK protein by structure modeling and molecular dynamics studies. *Journal of molecular modeling*, 19, 407-419. <https://doi.org/10.1007/s00894-012-1566-1>
- Klich, M. A. (2007). *Aspergillus flavus*: the major producer of aflatoxin. *Molecular plant pathology*, 8(6), 713-722. <https://doi.org/10.1111/j.1364-3703.2007.00436.x>
- Laskowski, R. A., Furnham, N., & Thornton, J. M. (2013). The Ramachandran plot and protein structure validation. In *Biomolecular forms and functions: a celebration of 50 years of the ramachandran map* (pp. 62-75). https://doi.org/10.1142/9789814449144_0005
- Lake, J. A. (1981). The ribosome. *Scientific American*, 245(2), 84-97. <https://doi.org/10.1038/scientificamerican0881-84>
- Lovell, S. C., Davis, I. W., Arendall III, W. B., De Bakker, P. I., Word, J. M., Prisant, M. G., ... & Richardson, D. C. (2003). Structure validation by C α geometry: ϕ , ψ and C β deviation. *Proteins: Structure, Function, and Bioinformatics*, 50(3), 437-450. <https://doi.org/10.1002/prot.10286>
- Lobo, I. (2008). Basic local alignment search tool (BLAST). *Nature Education*, 1(1).
- Nguyen, H. N., Chaves-Lopez, C., Oliveira, R. C., Paparella, A., & Rodrigues, D. F. (2019). Cellular and metabolic approaches to investigate the effects of graphene and graphene oxide in the fungi *Aspergillus flavus* and *Aspergillus niger*. *Carbon*, 143, 419-429. <https://doi.org/10.1016/j.carbon.2018.10.099>
- Ogodo, A. C., & Ugbogu, O. C. (2016). Public health significance of aflatoxin in food industry—A review. *European Journal of Clinical and Biomedical Sciences*, 2(5), 51-58. <https://doi.org/10.11648/j.ejcb.20160205.14>
- Rozario, L. T., Sharker, T., & Nila, T. A. (2021). In silico analysis of deleterious SNPs of human MTUS1 gene and their impacts on subsequent protein structure and function. *PloS one*, 16(6), e0252932. <https://doi.org/10.1371/journal.pone.0252932>
- Ramirez, M. L., Cendoya, E., Nichea, M. J., Zchetti, V. G. L., & Chulze, S. N. (2018). Impact of toxigenic fungi and mycotoxins in chickpea: A review. *Current Opinion in Food Science*, 23, 32-37. <https://doi.org/10.1016/j.cofs.2018.05.003>
- Szklarczyk, D., Franceschini, A., Kuhn, M., Simonovic, M., Roth, A., Minguetz, P., ... & Mering, C. V. (2010). The STRING database in 2011: functional interaction networks of proteins, globally integrated and scored. *Nucleic acids research*, 39(suppl_1), D561-D568. <https://doi.org/10.1093/nar/gkq973>
- Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., ... & von Mering, C. (2023). The STRING database in 2023: protein–protein association networks and functional

- enrichment analyses for any sequenced genome of interest. *Nucleic acids research*, 51(D1), D638-D646. <https://doi.org/10.1093/nar/gkac1000>
- Tam, E. W., Chen, J. H., Lau, E. C., Ngan, A. H., Fung, K. S., Lee, K. C., ... & Woo, P. C. (2014). Misidentification of *Aspergillus nomius* and *Aspergillus tamaris* as *Aspergillus flavus*: characterization by internal transcribed spacer, β -tubulin, and calmodulin gene sequencing, metabolic fingerprinting, and matrix-assisted laser desorption ionization–time of flight mass spectrometry. *Journal of Clinical Microbiology*, 52(4), 1153-1160. <https://doi.org/10.1128/JCM.03258-13>
- Von Hippel, P. H., Bear, D. G., Morgan, W. D., & McSwiggen, J. A. (1984). Protein-nucleic acid interactions in transcription: a molecular analysis. *Annual Review of Biochemistry*, 53(1), 389-446. <https://doi.org/10.1146/annurev.bi.53.070184.002133>
- Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., ... & Schwede, T. (2018). SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic acids research*, 46(W1), W296-W303. <https://doi.org/10.1093/nar/gky427>
- Yu, J., Cleveland, T. E., Nierman, W. C., & Bennett, J. W. (2005). *Aspergillus flavus* genomics: gateway to human and animal health, food safety, and crop resistance to diseases. *Revista iberoamericana de micología*, 22(4), 194-202. [https://doi.org/10.1016/S1130-1406\(05\)70043-7](https://doi.org/10.1016/S1130-1406(05)70043-7)
- Zain, M. E. (2011). Impact of mycotoxins on humans and animals. *Journal of Saudi chemical society*, 15(2), 129-144. <https://doi.org/10.1016/j.jscs.2010.06.006>

Copyediting Revision (27-12-2023)



Official publication of Pakistan Phytopathological Society
Pakistan Journal of Phytopathology
ISSN: 1019-763X (Print), 2305-0284 (Online)
<http://www.pakps.com>



IN SILICO ANALYSIS OF ASPERGILLUS FLAVUS FUNGAL PROTEINS: STRUCTURAL AND FUNCTIONAL INSIGHTS USING ITS PRIMERS

Mariana Mariana, Saipul Abbas*, Salamiah Salamiah, Samharinto Samharinto, Yusriadi Marsuni, Muslimin Sepe,

Muhammad I. Pramudi, Ismed S. Budi, Dewi Fitriyanti, Elly Liestiany, Lyswiana Aphrodyanti

Department of Plant Protection, Faculty of Agriculture, Lambung Mangkurat University. Jl.Jend. A. Yani Km.36 Banjarbaru. South Kalimantan 70714 Indonesia.

ABSTRACT

Aspergillus spp. is a type of fungi that can contaminate and damage various types of food and agricultural products. To better understand the structure and function of proteins, *in silico* research was conducted using Internal Transcribed Spacer (ITS) primers and SWISS Model protein prediction tools. The objective of this study was to identify fungal species using the BLAST method and to analyze the structure and function of proteins from *Aspergillus* spp. The methods used included BLAST for species identification, Web Expasy to translate DNA sequences into proteins, SWISS Models to model protein structures, SAVES to validate protein structures, and STRING to analyze the function of proteins. The results of the BLAST analysis showed that the identified fungal species were *Aspergillus flavus*, *A. tamarii*, and *A. nomius*. Furthermore, the results of translating DNA sequences into proteins using Web Expasy showed that there were three open reading frames with the highest residual values of 119 and 83, while the lowest residual value was 4. Only two of these frames met the protein criteria. Moreover, the results of protein structure modeling using the SWISS Model method produced a fairly accurate *Aspergillus* spp. protein structure model with a validation value of protein structure using ERRAT (SAVES V6.0) of 100%. Additionally, the results of protein function analysis using STRING showed that the *Aspergillus* spp. protein has a function in producing enzymes that play a role in the metabolic process of cells.

Keywords: *Aspergillus flavus*, *in silico*, ITS primer, protein structure, protein function, "SWISS Model".

INTRODUCTION

Aspergillus spp. fungus is a type of fungus that often causes contamination and damage to various agricultural and food products (Gourama & Bullerman, 1995), such as grains, nuts, and processed products (Ayofemi Olalekan Adeyeye, 2020). One of the fungi *Aspergillus* spp. that can contaminate feed ingredients is the species *A. flavus* which can cause aflatoxin pollution (Klich, 2007), which is a harmful substance that can affect human health and livestock (Balina et al, 2018). *A. flavus* is a group of fungi often found in agricultural products, especially in food crops and legumes (Amaike & Keller, 2011). Contamination by *A. flavus* can

cause damage to agricultural products and foodstuffs and can cause health problems in humans and farm animals that consume them (Yu et al, 2005). Therefore, research on *Aspergillus* spp. is very important in agriculture to prevent and control contamination of *Aspergillus* spp. on agricultural products and foodstuffs.

Previous studies have highlighted the negative impact of *Aspergillus* spp. on agricultural products and food items. Ramirez et al. (2018) noted that this type of fungus can cause damage to grains, legumes, and processed products. Similarly, Balina et al. (2018) indicated that *A. flavus* can be a major cause of aflatoxin contamination in feed materials, posing risks to livestock and humans who consume them. The significance of research on *A. flavus* is further underscored by the findings of Bhatnagar-Mathur et al. (2015), demonstrating the frequent presence of *A. flavus* in agricultural environments, with a high tendency to infect staple food crops. Therefore, contamination risk from *A. flavus* not only affects the quality of agricultural products and food items but also poses a serious health risk to consumers. *In silico* research has become a useful tool for studying the structure and function of proteins (Dutta et al, 2018), as it allows for quick and efficient hypothesis testing and analysis (Kirubakaran et al, 2013). In this study, we used an *in silico* technique to predict the structure and function of the *A. flavus* protein by combining the primary use of the

Submitted: June 28, 2023
Revised: July 17, 2023
Accepted for publication: December 25, 2021

* Corresponding Author:

Email: saipul.abbas@ulm.ac.id

© 2017 Pak. J. Phytopathol. All rights reserved.

Internal Transcribed Spacer (ITS) and the SWISS Model protein prediction tool. This *in silico* approach is essential to understanding this fungus and is beneficial to the food and agriculture industries.

The formulation of the problem of this study was to analyze the BLAST results of these *Aspergillus* spp. sequences, examine the protein structure related to contamination, evaluate the validity and confidence of the protein structure of *Aspergillus* spp. which was found, as well as examining the function of the protein *Aspergillus* spp. associated with contamination of corn feed and other foodstuffs, as well as to understand the structure and function of the *Aspergillus* spp. protein associated with the contamination aforementioned. Therefore, this research focused on the analysis of *Aspergillus* spp. sequence, protein structure analysis, and protein function analysis, with the hope of providing new insights and solutions to the problem of contamination in food.

This study aims to analyze the species *Aspergillus* spp. which often contaminates corn feed and other foodstuffs. This study is expected to provide information about the species *Aspergillus* spp. which contaminates corn feed and other foodstuffs, as well as the structure and function of the protein *Aspergillus* spp. The results of this study are anticipated to contribute to the development of methods for detecting and preventing contamination of *Aspergillus* spp. in corn feed and other foodstuffs.

MATERIALS AND METHODS

Identification of fungus species: This step involved the use of an Internal Transcribed Spacer (ITS) Primer to verify fungus species. This stage of research was carried out by taking DNA samples from fungi and conducting DNA amplification using primary ITS. DNA amplification was performed using the Polymerase Chain Reaction (PCR) technique. This step aimed to verify the species of fungus analyzed. In the PCR technique, DNA samples were placed together with ITS

primers (ITS1 5'-TCCGTAGGTGAACCTGCGG-3', and ITS4 5'-TCCTCCGCTTATTGATATGC-3') and polymerase enzymes, and a heating and cooling process was carried out to start and end the reaction. The result of the amplification process was more DNA sequences associated with ITS. After successful DNA amplification, the obtained DNA sequence was analyzed to verify the fungus species. Sequencing analysis was carried out using bioinformatics software such as BLAST (*Basic Local Alignment Search Tool*) (Lobo, 2008).

Translate DNA Sequencing into Proteins: After the fungus DNA sequence was successfully analyzed, the DNA sequence was then translated into proteins. This step was done using a tool such as the ExPasy Translate Tool (Bashir, 2022). This tool converted the DNA sequence into a protein sequence based on the genetic code (Ayaz et al, 2020). The results of the translation showed the amino acid sequence contained in the fungus protein.

Analysis of protein structure: This stage used the SWISS Model method to predict the protein structure of the fungus *Aspergillus* spp. At this stage, the input of protein sequencing data from ExPasy tools was carried out, and then protein modeling was made using the SWISS Model (Waterhouse et al, 2018). Analysis using the SWISS Model provided information regarding protein domains, protein motifs, and other information related to the structure and function of proteins (Bordoli & Schwede, 2012). The results of the analysis of the structure and function of the protein were re-examined to ensure the validity of the results. Validation was done by comparing the results with a trusted protein database such as UniProt.

Protein Function Analysis: The research method of protein function analysis with STRING involved the use of a database of protein interactions and the analysis of protein networks (Franceschini et al, 2012). In this study, *A. flavus* protein data were used in the STRING database. Protein data were inputted into a STRING database containing

information about protein interactions formed based on data from literature databases, scientific publications, and experimental studies (Szklarczyk et al, 2010). Analysis of protein interactions formed using algorithms available on the STRING database. This algorithm generated information about the interaction of formed proteins, the forces of interaction, and their relationship with biological functions. Interpretation of the results of protein interaction analysis was done to obtain information about the biological function of the protein being analyzed and how the protein interaction was related to the biological function found. This method could be used to analyze the function of proteins and understand how proteins interacted with each other in a network of proteins (Szklarczyk et al, 2023). Using this method, we could gain a better understanding of how proteins were involved in biological processes and their possible effects on health and disease.

RESULTS AND DISCUSSION

Identification of fungus species by BLAST

Method: The results of the fungus species identification using BLAST revealed a similarity of 99.61% with *Aspergillus tamaraii*, 98.94% with *Aspergillus flavus*, and 96.33% with *Aspergillus nomius*. These findings suggested that the sample likely belonged to the *Aspergillus* group, with a

higher similarity to *A. tamaraii* compared to the other two species. *A. tamaraii* is recognized for its capability to produce various bioactive compounds (Bose et al, 2019), including phenol, flavonoid and IAA (Indole-3-Acetic Acid) and its antifungal activity, which have potential applications in the Agriculture and food industries. However, it is important to note that *A. tamaraii* can also cause infections in humans and animals (Tam et al, 2014). Similarly, *A. flavus* is known for its production of toxic compounds, such as aflatoxin, which can be harmful to human and animal health (Fouad et al, 2019). Therefore, proper handling measures are necessary to prevent the spread of this species. Moreover, *A. nomius*, another identified species in the sample, can produce aflatoxin and the potential to become pathogenic in humans and animals (Zain, 2011). Consequently, accurate identification of the fungus species is essential to determine appropriate control measures. In this case, the identification results indicating the highest similarity to *A. tamaraii* and *A. flavus* can serve as references for further research on the bioactive compounds produced by these species. Nevertheless, further examinations are required to confirm the presence of specific fungus species in the sample and establish appropriate control measures to prevent its spread.

select all 10 sequences selected		GenBank	Graphics	Distance tree of results	MSA Viewer			
Description	Scientific Name	Max Score	Total Score	Query Cover	E value	Per. Ident	Acc. Len	Accession
<input checked="" type="checkbox"/> Aspergillus tamaraii strain M24 small subunit ribosomal RNA gene, partial sequence: internal transcribed spacer 1, 5.8S...	Aspergillus tamaraii	1419	2347	97%	0.0	99.61%	889	ON819577.1
<input checked="" type="checkbox"/> Aspergillus flavus strain Bp5 18S ribosomal RNA gene, partial sequence: internal transcribed spacer 1 and 5.8S riboso...	Aspergillus flavus	1343	2094	94%	0.0	98.94%	939	KF221065.1
<input checked="" type="checkbox"/> Aspergillus tamaraii strain BPF17 small subunit ribosomal RNA gene, partial sequence: internal transcribed spacer 1,...	Aspergillus tamaraii	1321	2151	99%	0.0	96.86%	897	MF359724.1
<input checked="" type="checkbox"/> Aspergillus tamaraii strain MF24 small subunit ribosomal RNA gene, partial sequence: internal transcribed spacer 1, 5.8...	Aspergillus tamaraii	1310	1952	98%	0.0	96.83%	791	MF359732.1
<input checked="" type="checkbox"/> Aspergillus sp. AQQGD 17 internal transcribed spacer 1, partial sequence: 5.8S ribosomal RNA gene and internal tran...	Aspergillus sp. A...	1297	2462	100%	0.0	96.37%	1303	KP721583.1
<input checked="" type="checkbox"/> Aspergillus sp. isolate ACSIKS_2100168 small subunit ribosomal RNA gene, partial sequence: internal transcribed spa...	Aspergillus sp.	1291	2430	100%	0.0	96.25%	1187	MN597042.1
<input checked="" type="checkbox"/> Aspergillus sp. isolate SS_12 small subunit ribosomal RNA gene, partial sequence: internal transcribed spacer 1, 5.8S...	Aspergillus sp.	1286	2440	100%	0.0	96.12%	1303	MT497446.1
<input checked="" type="checkbox"/> Aspergillus flavus isolate 20 small subunit ribosomal RNA gene, partial sequence: internal transcribed spacer 1 and 5.8...	Aspergillus flavus	1273	1946	91%	0.0	98.18%	715	MK120614.1
<input checked="" type="checkbox"/> Aspergillus nomius strain Tur4 18S ribosomal RNA gene, partial sequence: internal transcribed spacer 1 and 5.8S ribos...	Aspergillus nomiae	1240	1891	95%	0.0	96.33%	842	KF221090.1
<input checked="" type="checkbox"/> Aspergillus flavus strain Beca_43 small subunit ribosomal RNA gene, partial sequence: internal transcribed spacer 1, 5...	Aspergillus flavus	1234	2105	99%	0.0	96.80%	849	KY234269.1

Figure 1. The results of the fungus species identification using BLAST Method (GenBank)

Translate DNA Sequencing into Proteins: The results obtained from the translation of *Aspergillus* spp. DNA sequences into proteins using the ExPASy web revealed the presence of three open reading frames (ORFs), indicated in red, with residual values of 119 and 83 as the highest, and 4 as the lowest. Proteins are typically categorized if they consist of more than 50 residues (Damodaran & Parkin, 2017). In this case, only two out of the three ORFs met the criteria to be considered as proteins. The identification of an ORF qualifying as a protein may indicate the presence of a gene within the *Aspergillus* DNA sequence. Translating DNA sequences into proteins holds significant importance in molecular

biology research as it allows for a better understanding of protein functions within a specific organism (Gasteiger et al, 2003). Consequently, these findings serve as a foundation for future investigations concerning the *Aspergillus* gene and its potential applications in various fields, such as agriculture and the food industry. Nevertheless, it is crucial to note that these results should undergo further verification using other techniques, such as protein structure analysis, to ensure the functional integrity of the resulting proteins. Additionally, the authenticity and suitability of the *Aspergillus* DNA sequence utilized in this study should be validated about the specific *Aspergillus* species under investigation.



Figure 2. The results of *Aspergillus* spp. DNA sequences into proteins using the ExPASy web Table 1. The results of Open Reading Frames (ORFs) with residual values from 4 to 119

Open Reading Frame	Residue	Types of Molecules	Code Peptides/Proteins
MNGGPAEATKVQ	12	Peptide	A1
MAGPPKQLRYSKHGWEVGLARNPTLGND PSVGEPAEGLSPVGFASPTSHPCLLYLSC FGGPAIHGRRGLSAPGPRPETPRTLSDLV KSELIVSQSVKTFNNGSLGSGIDEERSEMR	119	Protein	A2
MPVRASLLPIKHGLCVGSSSPLRGGRAPKA AAAPRILERMGLCHPLCRPGRRLPNANQS FSRLTSDQVGIPAEKHKINKAEE	83	Protein	A3
MILP	4	Peptide	A4
MAY	4	Peptide	A5

MAAGGSQPRARARRRHHELCLI	22	Peptide	A6
MDLLVPASMKNAAKCDN	17	Peptide	A7

Protein Structure Modeling with the SWISS Model method: The results of modeling the protein structure of *Aspergillus* spp. using the SWISS Model method show the presence of a 3D structure of proteins formed (Bordoli & Schwede, 2012). Protein structure modeling is an important method in molecular biology research because it can provide information about the shape and function of proteins in a particular organism (Breda et al, 2007). The results of modeling the protein structure of *Aspergillus* spp. can be used as a basis for conducting further analysis of protein function, including interactions with other molecules, enzymatic activity, and possible drug development. In addition, these results can also help in understanding the relationship between the structure of proteins and their biological activity. However, keep in mind that the results of

modeling protein structures using computational methods such as the SWISS Model still have limitations, especially in terms of accuracy. Therefore, these results still need to be verified using other techniques, such as X-ray crystallography or core magnetic resonance spectroscopy (NMR), to ensure the accuracy of the resulting protein structure. In addition, modeling the protein structure of *Aspergillus* spp. can also provide information about the evolutionary relationship of proteins between different *Aspergillus* species. Modeling the structure of the *Aspergillus* spp. protein can help understand the differences and similarities between different *Aspergillus* proteins so that it can be used as a basis for specific drug and therapeutic development.



Figure 3. Protein Structure Prediction from Internal Transcribed Spacer (ITS) primer of *Aspergillus flavus* fungus using Protein Prediction from SWISS Model. (Template: A2: [3lb6.1.A](#) A3: 5t7v.1.K)

The results of validation of the *Aspergillus* spp. protein structure using SAVES V6.0 show that the resulting protein structure has a fairly good value in terms of accuracy and quality. One of the parameters used to evaluate the validation of protein structure is the Ramachandran plot, which shows the proportion of amino acids placed in the

most favored regions within the protein conformation space (Laskowski et al, 2013). In the Ramachandran plot results obtained, the highest Residual value in most favored regions was 80% for the protein code A2, which indicates that most of the amino acids in the protein structure of *Aspergillus* spp. were placed in the desired area

(Lovell et al, 2003). In addition, the validation results using the ERRAT method also show the highest value of 100%, which indicates that the resulting protein structure has high accuracy (Rozario et al, 2021). The results of the validation of the protein structure of *Aspergillus* spp. using SAVES V6.0 are important to ensure that the resulting protein structure is trustworthy and used as a basis for further analysis of protein function and drug development. Validation of protein structure can also help identify and correct parts in the protein structure that may be inaccurate, thereby improving the quality and accuracy of research results.



The QMEAN4 Score indicates the overall quality of the protein structure, which is calculated based on geometric values, non-covalent interactions, and consistency of the structure (Benkert et al, 2008). The higher the QMEAN 4 Score, the better the quality of the resulting protein structure (Benkert et al, 2011). In this case, the QMEAN4 Score A2



value of -1.72 versus A3 of -3.27 can be interpreted as a fairly low value, which indicates that the resulting *Aspergillus* spp. protein structure does not have optimal quality. Meanwhile, the Z Score is a parameter used to compare the quality of the resulting protein structure with other similar protein structures, which are calculated based on geometric values and structural consistency. The higher the Z Score value, the better the quality of the resulting protein structure compared to other similar protein structures (Bhattacharya et al, 2007). In this case, a Z Score of -0.16 indicates that the resulting *Aspergillus* spp. protein structure is of relatively good quality compared to other similar protein structures. However, keep in mind that as with the QMEAN 4 Score value, the Z Score value also has limitations and possible errors, so this result still needs to be verified using other protein structure validation techniques.

Table 2. Protein Structure Validation using SAVES V6.0



Protein Code	Ramachandran plot				ERRAT quality factor (%)	QMEAN 4 Score	Z Score
	Residu in most favored regions (%)	Residu in additional allowed regions (%)	Residu in generously allowed regions (%)	Residu in disallowed regions (%)			
A2	80.0	20.0	0.0	0.0	100	-1.72	-0.16
A3	75.0	20.0	0.0	5.0	6.6666	-3.27	-3.32

Model A2





Model A3






Figure 4. The results of validation of the *Aspergillus* spp. protein structure using SAVES V6.0

Protein Function & Interaction of *Aspergillus Flavus*:

A. flavus is a type of fungus known to cause damage to agricultural products such as grains and other foodstuffs (Gourama & Bullerman, 1995). This fungus can also produce some toxic substances such as aflatoxins, which are heterocyclic substances and have detrimental effects on human and animal health (Dhanamjayulu et al, 2019). Heterocyclics in aflatoxins have several main functions in the fungus *A. flavus*, including: (1) As a toxic compound: Aflatoxins have high toxic activity and can cause food poisoning in humans and animals, which can cause various health problems such as liver damage, cancer, and reproductive problems (Ogodo & Ugboogu, 2016). (2) As an antioxidant compound: Aflatoxin has antioxidant properties and can help the fungus *A. flavus* survive in a poor environment (Abrar et al, 2013). (3) As an antifungal compound: Aflatoxin can help the fungus *A. flavus* resist attacks from other fungi that seek to interfere with the growth of the fungus *A. flavus* (Bhatnagar-Mathur et al, 2015). Overall, heterocyclic in aflatoxins has an important role in the biology of the fungus *A. flavus*, but it also has detrimental effects on human and animal health. Therefore, it is important to control the production of aflatoxins and ensure that the food products consumed do not contain this substance in harmful quantities.

Organic cyclic compound binding is the process of binding organic chemical compounds that have a benzene ring (Cram & Cram, 1971). In this case, such binding may take place in the fungus *A. flavus*. Organic chemical compounds that bind to the fungus *A. flavus* can affect the biological activity of the fungus (Nguyen et al, 2019). Some organic chemical compounds that can bind to the fungus *A. flavus* include growth regulators, metabolic regulators, and anti-fungal substances (Hou et al, 2022). The binding of organic chemical compounds to the fungus *A. flavus* can affect the growth and reproduction of fungi, modulate enzyme activity, affect the production of toxic substances such as aflatoxins, and affect the interaction of fungi with their environment. Overall, the binding function of organic chemical compounds to the fungus *A. flavus* is essential for understanding the biology of the fungus and how the fungus interacts with its environment. Further research on the binding of organic chemical compounds to the fungus *A. flavus* can help in controlling the production of aflatoxins and improving the quality of food products. Nucleic acid binding protein has a very important function in the fungal cells of *A. flavus*. Nucleic acids are important molecules that carry genetic information and play a key role in various biological processes, such as DNA replication, RNA

transcription, and protein synthesis (Blackburn, 2006). Nucleic acid binding proteins can bind to nucleic acid molecules, facilitate interaction with enzymes and transcription factors, and influence the biological activity of nucleic acid molecules (Von Hippel et al, 1984). In the fungus *A. flavus*, Nucleic acid binding proteins may play a role in regulating gene activity, affecting growth and reproduction, and modulating the fungus' response to the environment. Studies of Nucleic acid binding proteins in the fungus *A. flavus* can provide information about the mechanisms of gene regulation and how fungi react to their environment. Overall, the function of the Nucleic acid binding protein in the fungus *A. flavus* is essential for understanding the biology of the fungus and how the fungus interacts with its environment. More research on Nucleic acid binding proteins in the fungus *A. flavus* can help understand the growth and reproduction mechanisms of fungi, as well as how fungi react to their environment.

RNA binding is a process in which proteins bind to RNA molecules. RNA is an important molecule in cells that plays a key role in biological processes such as transcription, protein synthesis, and gene regulation (Elliott & Ladomery, 2017). RNA protein binding can bind to RNA molecules and facilitate interaction with enzymes and transcription factors (Corley et al, 2020). The function of RNA binding proteins can affect the biological activities of RNA molecules, such as modulating transcription and protein synthesis, affecting RNA stability, and facilitating gene control (Gilsovic et al, 2008). In the fungus, *A. flavus*, protein RNA binding may play an important role in regulating gene activity, modulating responses to the environment, and facilitating growth and reproduction. Studies on protein RNA binding in the fungus *A. flavus* may help understand the mechanisms of gene regulation and how fungi react to their environment. Overall, the function of RNA binding proteins in the fungus *A. flavus* is critical to understanding the biology of the fungus and how the fungus interacts with its environment. More research on protein RNA binding in the fungus *A. flavus* may help in understanding the mechanisms of fungal

growth and reproduction, as well as how the fungus reacts to its environment.

mRNA binding is a process in which proteins bind to mRNA molecules (Dreyfuss et al, 2002). mRNA is an RNA molecule that contains genetic information from DNA and is used as a template for protein synthesis (Alberts et al, 2002). mRNA protein binding can bind to mRNA molecules and facilitate interaction with enzymes and transcription factors (Corley et al, 2020). In the fungus *A. flavus*, mRNA protein binding may play an important role in regulating gene activity, modulating responses to the environment, and facilitating growth and reproduction. Studies on mRNA protein binding in the fungus *A. flavus* may help understand the mechanisms of gene regulation and how fungi react to their environment. Overall, the function of mRNA protein binding in the fungus *A. flavus* is critical to understanding the biology of the fungus and how the fungus interacts with its environment. More research on mRNA protein binding in the fungus *A. flavus* may help understand the mechanisms of fungal growth and reproduction, as well as how the fungus reacts to its environment. Ribosomes are cellular structures that play a key role in the process of protein synthesis (Frank & Spahn, 2006). Ribosomes consist of two subunits, a large subunit and a small subunit, which serve to bind amino acids and form polypeptide chains (Lake, 1981). Proteins that are structural components of ribosomes can play an important role in ensuring the integrity and activity of ribosomes. The function of proteins as structural components of ribosomes can affect the ability of ribosomes to bind to mRNA and facilitate proper protein synthesis (de la Cruz et al, 2015). In the fungus *A. flavus*, proteins that are structural components of ribosomes may play an important role in ensuring the growth and reproduction of fungi. The study of proteins that are structural components of ribosomes in the fungus *A. flavus* can help understand the mechanism of protein synthesis and how the fungus reacts to its environment. Overall, the function of proteins as structural components of ribosomes in the fungus *A. flavus* is essential for understanding the biology of

the fungus and how the fungus interacts with its environment. More research on the proteins that are structural components of ribosomes in the

fungus *A. flavus* can help in understanding the mechanisms of fungal growth and reproduction, as well as how the fungus reacts to its environment.




Figure 5: Network of protein-protein interactions visualized with STRING (NCBI taxon-Id: 332952)

B8MXX8	<i>Serine/threonine protein kinase, putative</i>
B8N3J1	<i>Small nucleolar ribonucleoprotein complex subunit Utp14, putative</i>
B8NT05	<i>G-protein complex beta subunit CpcB</i>
B8N430	<i>Nuclear mRNA splicing factor, putative</i>
B8N0Y2	<i>G-patch domain protein (TFIP11), putative</i>
B8N058	<i>PWI domain mRNA processing protein, putative</i>
B8N1X2	<i>Pre-RNA splicing factor Srp2, putative</i>
B8ND38	<i>CaaX prenyl proteinase Rce1</i>
B8MXA7	<i>Small nuclear ribonucleoprotein E; Associated with the spliceosome snRNP U1, U2, U4/U6 and U5.</i>
B8MXX7	<i>Casein kinase, putative; Belongs to the protein kinase superfamily.</i>

Table 3: Molecular Function, Biological Process, Cellular Component, and Subcellular Localization Classification

Molecular Function	Biological Process	Cellular Component	Subcellular Localization
<ul style="list-style-type: none"> Heterocyclic compound binding Organic cyclic compound binding Nucleic acid binding RNA binding mRNA binding Structural constituent of ribosome rRNA binding snoRNA binding 	<ul style="list-style-type: none"> mRNA splicing, via spliceosome RNA processing Gene expression Macromolecule metabolic process Cellular nitrogen compound metabolic process Nitrogen compound metabolic process Primary metabolic process Cellular metabolic process 	<ul style="list-style-type: none"> Spliceosomal complex Ribonucleoprotein complex Nucleus Intracellular membrane-bounded organelle 	<ul style="list-style-type: none"> U2 snRNP U2-type prespliceosome U2-type spliceosomal complex Spliceosomal complex Ribonucleoprotein complex Nucleus Protein-containing complex Intracellular membrane-bounded organelle Intracellular Cellular anatomical entity

CONCLUSION

In silico research was conducted to understand the structure and function of *proteins of Aspergillus* spp. The results of the BLAST analysis showed the presence of three species of fungi, namely *Aspergillus flavus*, *A. tamarii*, and *A. nomius*. Translating DNA sequences into proteins using Web ExPasy showed that only two proteins met the criteria. Protein structure modeling using SWISS Model resulted in a fairly accurate *Aspergillus* spp. protein structure model with protein structure validation values using ERRAT (SAVES V6.0) of 100%. Analysis of protein function using STRING shows that *Aspergillus* spp. protein has a function in producing enzymes that play a role in the metabolic process of cells. Thus, this study provides important information about the structure and function of the *protein Aspergillus* spp. which can be used to improve understanding of contamination and damage to food and agricultural products by this type of fungus.

ACKNOWLEDGEMENT

A thank you to PT. Arutmin Indonesia Tambang Satui and Agreement/Contract Number: 39/UN8.1.23/PP/2022 through the Matching Fund program "Kedaireka" for financial support for this research activity, and all parties who have helped in the success of this research. All the help and support provided means a lot to us. Thanks.

REFERENCES

Ayaz, A., Zaman, W., Saqib, S., Ullah, F., & Mahmood, T. (2020). Phylogeny and diversity of Lamiaceae based on rps14 gene in Pakistan. *Genetika*, 52(2), 435-452. <https://doi.org/10.2298/GENSR2002435A>

Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., & Walter, P. (2002). From DNA to RNA. In *Molecular Biology of the Cell. 4th edition*. Garland Science.

Abrar, M., Anjum, F. M., Butt, M. S., Pasha, I., Randhawa, M. A., Saeed, F., & Waqas, K. (2013). Aflatoxins: biosynthesis, occurrence, toxicity, and remedies. *Critical reviews in food science and nutrition*, 53(8), 862-874.

<https://doi.org/10.1080/10408398.2011.563154>

- Ayofemi Olalekan Adeyeye, S. (2020). Aflatoxigenic fungi and mycotoxins in food: a review. *Critical reviews in food science and nutrition*, 60(5), 709-721. <https://doi.org/10.1080/10408398.2018.1548429>
- Amaike, S., & Keller, N. P. (2011). *Aspergillus flavus*. *Annual review of phytopathology*, 49, 107-133. <https://doi.org/10.1146/annurev-phyto-072910-095221>
- Bhatnagar-Mathur, P., Sunkara, S., Bhatnagar-Panwar, M., Waliyar, F., & Sharma, K. K. (2015). Biotechnological advances for combating *Aspergillus flavus* and aflatoxin contamination in crops. *Plant science*, 234, 119-132. <https://doi.org/10.1016/j.plantsci.2015.02.009>
- Bhattacharya, A., Tejero, R., & Montelione, G. T. (2007). Evaluating protein structures determined by structural genomics consortia. *Proteins: Structure, Function, and Bioinformatics*, 66(4), 778-795. <https://doi.org/10.1002/prot.21165>
- Blackburn, G. M. (Ed.). (2006). *Nucleic acids in chemistry and biology*. Royal Society of Chemistry.
- Breda, A., Valadares, N. F., de Souza, O. N., & Garratt, R. C. (2007). Protein structure, modelling and applications. In *Bioinformatics in tropical disease research: a practical and case-study approach [Internet]*. National Center for Biotechnology Information (US). <https://www.ncbi.nlm.nih.gov/books/NBK6824/>
- Benkert, P., Tosatto, S. C., & Schomburg, D. (2008). QMEAN: A comprehensive scoring function for model quality assessment. *Proteins: Structure, Function, and Bioinformatics*, 71(1), 261-277. <https://doi.org/10.1002/prot.21715>
- Benkert, P., Biasini, M., & Schwede, T. (2011).

- Toward the estimation of the absolute quality of individual protein structure models. *Bioinformatics*, 27(3), 343-350. <https://doi.org/10.1093/bioinformatics/btq662>
- Bordoli, L., & Schwede, T. (2012). Automated protein structure modeling with SWISS-MODEL Workspace and the Protein Model Portal. *Homology Modeling: Methods and Protocols*, 107-136. https://doi.org/10.1007/978-1-61779-588-6_5
- Bose, P., Gowrie, S. U., & Chathurdevi, G. (2019). Optimization of Culture Conditions for Growth and Production of Bioactive Metabolites by Endophytic Fungus-*Aspergillus tamarii*. *International Journal of Pharmacy and Biological Sciences*, 9(2), 469-478. <https://doi.org/10.21276/ijpbs.2019.9.2.60>
- Balina, A., Kebede, A., & Tamiru, Y. (2018). Review on aflatoxin and its impacts on livestock. *Journal of Dairy and Veterinary Sciences*, 6(2), e555685. <https://doi.org/10.19080/JDVS.2018.06.555685>
- Bashir, S., Janiad, S., Zeeshan, A., Farah, N., Mehboob, S., Butt, H. I., & Qari, S. H. (2022). In Silico Analysis and Function Prediction of Sr22 Gene Product as Stem Rust Resistant Protein. *Annals of the Romanian Society for Cell Biology*, 26(01), 258-262. <http://annalsofscrb.ro>
- Cram, D. J., & Cram, J. M. (1971). Cyclophane chemistry: bent and battered benzene rings. *Accounts of Chemical Research*, 4(6), 204-213. <https://doi.org/10.1021/ar50042a003>
- Corley, M., Burns, M. C., & Yeo, G. W. (2020). How RNA-binding proteins interact with RNA: molecules and mechanisms. *Molecular cell*, 78(1), 9-29. <https://doi.org/10.1016/j.molcel.2020.03.011>
- de la Cruz, J., Karbstein, K., & Woolford Jr, J. L. (2015). Functions of ribosomal proteins in assembly of eukaryotic ribosomes in vivo. *Annual review of biochemistry*, 84, 93-129. <https://doi.org/10.1146/annurev-biochem-060614-033917>
- Dhanamjayulu, P., Boga, R. B., & Mehta, A. (2019). Inhibition of aflatoxin B1 biosynthesis and down regulation of aflR and aflB genes in presence of benzimidazole derivatives without impairing the growth of *Aspergillus flavus*. *Toxicon*, 170, 60-67. <https://doi.org/10.1016/j.toxicon.2019.09.018>
- Damodaran, S., & Parkin, K. L. (2017). Amino acids, peptides, and proteins. In *Fennema's food chemistry* (pp. 235-356). CRC Press.
- Dreyfuss, G., Kim, V. N., & Kataoka, N. (2002). Messenger-RNA-binding proteins and the messages they carry. *Nature reviews Molecular cell biology*, 3(3), 195-205. <https://doi.org/10.1038/nrm760>
- Dutta, B., Banerjee, A., Chakraborty, P., & Bandopadhyay, R. (2018). In silico studies on bacterial xylanase enzyme: Structural and functional insight. *Journal of Genetic Engineering and Biotechnology*, 16(2), 749-756. <https://doi.org/10.1016/j.jgeb.2018.05.003>
- Elliott, D., & Ladomery, M. (2017). *Molecular biology of RNA*. Oxford University Press.
- Frank, J., & Spahn, C. M. (2006). The ribosome and the mechanism of protein synthesis. *Reports on Progress in Physics*, 69(5), 1383. <https://doi.org/10.1088/0034-4885/69/5/R03>
- Fouad, A. M., Ruan, D., El-Senousey, H. K., Chen, W., Jiang, S., & Zheng, C. (2019). Harmful effects and control strategies of aflatoxin b1 produced by *Aspergillus flavus* and *Aspergillus parasiticus* strains on poultry. *Toxins*, 11(3), 176. <https://doi.org/10.3390/toxins11030176>
- Franceschini, A., Szklarczyk, D., Frankild, S., Kuhn, M., Simonovic, M., Roth, A., ... & Jensen, L. J.

- (2012). STRING v9. 1: protein-protein interaction networks, with increased coverage and integration. *Nucleic acids research*, 41(D1), D808-D815. <https://doi.org/10.1093/nar/gks1094>
- Glisovic, T., Bachorik, J. L., Yong, J., & Dreyfuss, G. (2008). RNA-binding proteins and post-transcriptional gene regulation. *FEBS letters*, 582(14), 1977-1986. <https://doi.org/10.1016/j.febslet.2008.03.004>
- Gourama, H., & Bullerman, L. B. (1995). *Aspergillus flavus* and *Aspergillus parasiticus*: Aflatoxigenic fungi of concern in foods and feeds: A review. *Journal of Food protection*, 58(12), 1395-1404. <https://doi.org/10.4315/0362-028X-58.12.1395>
- Gasteiger, E., Gattiker, A., Hoogland, C., Ivanyi, I., Appel, R. D., & Bairoch, A. (2003). ExPASy: the proteomics server for in-depth protein knowledge and analysis. *Nucleic acids research*, 31(13), 3784-3788. <https://doi.org/10.1093/nar/gkg563>
- Hou, T., Sana, S. S., Li, H., Xing, Y., Nanda, A., Netala, V. R., & Zhang, Z. (2022). Essential oils and its antibacterial, antifungal and anti-oxidant activity applications: A review. *Food Bioscience*, 47, 101716. <https://doi.org/10.1016/j.fbio.2022.101716>
- Kirubakaran, P., Karthikeyan, M., Singh, K. D., Nagamani, S., & Premkumar, K. (2013). In silico structural and functional analysis of the human TOPK protein by structure modeling and molecular dynamics studies. *Journal of molecular modeling*, 19, 407-419. <https://doi.org/10.1007/s00894-012-1566-1>
- Klich, M. A. (2007). *Aspergillus flavus*: the major producer of aflatoxin. *Molecular plant pathology*, 8(6), 713-722. <https://doi.org/10.1111/j.1364-3703.2007.00436.x>
- Laskowski, R. A., Furnham, N., & Thornton, J. M. (2013). The Ramachandran plot and protein structure validation. In *Biomolecular forms and functions: a celebration of 50 years of the ramachandran map* (pp. 62-75). https://doi.org/10.1142/9789814449144_0005
- Lake, J. A. (1981). The ribosome. *Scientific American*, 245(2), 84-97. <https://doi.org/10.1038/scientificamerican0881-84>
- Lovell, S. C., Davis, I. W., Arendall III, W. B., De Bakker, P. I., Word, J. M., Prisant, M. G., ... & Richardson, D. C. (2003). Structure validation by C α geometry: ϕ , ψ and C β deviation. *Proteins: Structure, Function, and Bioinformatics*, 50(3), 437-450. <https://doi.org/10.1002/prot.10286>
- Lobo, I. (2008). Basic local alignment search tool (BLAST). *Nature Education*, 1(1).
- Nguyen, H. N., Chaves-Lopez, C., Oliveira, R. C., Paparella, A., & Rodrigues, D. F. (2019). Cellular and metabolic approaches to investigate the effects of graphene and graphene oxide in the fungi *Aspergillus flavus* and *Aspergillus niger*. *Carbon*, 143, 419-429. <https://doi.org/10.1016/j.carbon.2018.10.099>
- Ogodo, A. C., & Ugbogu, O. C. (2016). Public health significance of aflatoxin in food industry—A review. *European Journal of Clinical and Biomedical Sciences*, 2(5), 51-58. <https://doi.org/10.11648/j.ejcb.20160205.14>
- Rozario, L. T., Sharker, T., & Nila, T. A. (2021). In silico analysis of deleterious SNPs of human MTUS1 gene and their impacts on subsequent protein structure and function. *PloS one*, 16(6), e0252932. <https://doi.org/10.1371/journal.pone.0252932>
- Ramirez, M. L., Cendoya, E., Nichea, M. J., Zchetti, V. G. L., & Chulze, S. N. (2018). Impact of toxigenic fungi and mycotoxins in chickpea: A review. *Current Opinion in Food Science*, 23, 32-37.

- <https://doi.org/10.1016/j.cofs.2018.05.003>
Szklarczyk, D., Franceschini, A., Kuhn, M., Simonovic, M., Roth, A., Minguez, P., & Mering, C. V. (2010). The STRING database in 2011: functional interaction networks of proteins, globally integrated and scored. *Nucleic acids research*, 39(suppl_1), D561-D568. <https://doi.org/10.1093/nar/gkq973>
- Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., ... & von Mering, C. (2023). The STRING database in 2023: protein–protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic acids research*, 51(D1), D638-D646. <https://doi.org/10.1093/nar/gkac1000>
- Tam, E. W., Chen, J. H., Lau, E. C., Ngan, A. H., Fung, K. S., Lee, K. C., & Woo, P. C. (2014). Misidentification of *Aspergillus nomius* and *Aspergillus tamarii* as *Aspergillus flavus*: characterization by internal transcribed spacer, β -tubulin, and calmodulin gene sequencing, metabolic fingerprinting, and matrix-assisted laser desorption ionization–time of flight mass spectrometry. *Journal of Clinical Microbiology*, 52(4), 1153-1160. <https://doi.org/10.1128/JCM.03258-13>
- Von Hippel, P. H., Bear, D. G., Morgan, W. D., & McSwiggen, J. A. (1984). Protein-nucleic acid interactions in transcription: a molecular analysis. *Annual Review of Biochemistry*, 53(1), 389-446. <https://doi.org/10.1146/annurev.bi.53.070184.002133>
- Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., ... & Schwede, T. (2018). SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic acids research*, 46(W1), W296-W303. <https://doi.org/10.1093/nar/gky427>
- Yu, J., Cleveland, T. E., Nierman, W. C., & Bennett, J. W. (2005). *Aspergillus flavus* genomics: gateway to human and animal health, food safety, and crop resistance to diseases. *Revista iberoamericana de micología*, 22(4), 194-202. [https://doi.org/10.1016/S1130-1406\(05\)70043-7](https://doi.org/10.1016/S1130-1406(05)70043-7)
- Zain, M. E. (2011). Impact of mycotoxins on humans and animals. *Journal of Saudi chemical society*, 15(2), 129-144. <https://doi.org/10.1016/j.jscs.2010.06.006>

Contribution of Authors:

Mariana Mariana	:
Saipul Abbas	:
Salamiah Salamiah	:
Samharinto	:
Samharinto	:
Yusriadi Marsuni	:
Muslimin Sepe	:
Muhammad I.	:
Pramudi	:
Ismed S. Budi	:
Dewi Fitriyanti	:
Elly Liestiany	:
Lyswiana	:
Aphrodyanti	: