

Antifungal Activities of the Combination of Ulin Wood Liquid Smoke and Hiyung Cayenne Pepper Root Endophyte Fungi Against *Colletotrichum capsici*

by Mariana -

Submission date: 18-Jun-2024 11:17AM (UTC+0700)

Submission ID: 2404595238

File name: Turnitin_Agrivita_1.pdf (959.7K)

Word count: 5675

Character count: 31381



Antifungal Activities of the Combination of Ulin Wood Liquid Smoke and Hiyung Cayenne Pepper Root Endophyte Fungi Against *Colletotrichum capsici*

Witiyasti Imaningsih^{1,2*)}, Mariana³⁾, Ahmad Budi Junaedi⁴⁾ and Rasyidah²⁾

¹⁾ Biology Study Program, Mathematics and Science Faculty, Lambung Mangkurat University, South Kalimantan, Indonesia

²⁾ Microbiology Laboratory, Mathematics and Science Faculty, Lambung Mangkurat University, South Kalimantan, Indonesia

³⁾ Plant Protection Study Program, Faculty of Agriculture, Lambung Mangkurat University, South Kalimantan, Indonesia

⁴⁾ Chemistry Studi Program, Mathematics and Science Faculty, Lambung Mangkurat University, South Kalimantan, Indonesia

ARTICLE INFO

Keywords:

Endophytic fungi
Eusideroxylon zwageri Tejism. & Binn.
Hiyung cayenne pepper
 Inhibitory effect

Article History:

Received: November 18, 2019

Accepted: December 14, 2020

^{*)} Corresponding author:
 E-mail: witiyastimaningsih@ulm.ac.id

ABSTRACT

Chili farming faces several constraints, one of which is the pathogenic fungus *Colletotrichum capsici*. To overcome it can be used indigenous endophytic fungus and liquid smoke wood Ulin (*Eusideroxylon zwageri* Tejism. & Binn.) which has the potential as antimicrobial can be used. This research aimed to quantify and measure the effectiveness of an antimicrobial liquid smoke, endophytic filtrate, and the combination to suppress *C. capsici* growth. Subsequently, the research was conducted to apply the liquid smoke, endophytic fungi, and the two combinations of treatments on the growth of *C. capsici*. Thus, the results of this research showed that liquid smoke with a concentration of 0.085-1.75% can inhibit 3.56-62.17% in range. Meanwhile, the endophytic fungi filtrate, of 2% concentration can inhibit 91.69% *C. capsici*. Two of the combination liquid smoke in a concentration of 0.68%, 37.6% and the endophytic fungi filtrate in 2% have a demonstrated to inhibit the growth of *C. capsici* with the highest inhibition into 88.08%. Based on the analysis results, liquid smoke, endophytic fungi filtrate, and a combination of both showed significantly different inhibitory effects between treatments. This indicates that all those three treatments have antimicrobial potential.

INTRODUCTION

Chilli plant is one of the horticultural commodities that has economical value in Indonesia. The demand for chili is increasing, encouraging farmers to cultivate chili plants. One of the chilies that are cultivated by farmers in the Tapin area of South Kalimantan namely *Hiyung* cayenne pepper. The year 2012 *Hiyung* chili pepper is listed on the Center for Crop varieties Protection and Agriculture Licensing Ministry of Agriculture Republic of Indonesia No. 09/PLV/2012 dated 12 April 2012 as a local variety with the name of *Hiyung* cayenne pepper (Balai Penelitian dan Pengembangan Pertanian, 2018), and in June 2012 the Ministry of Agriculture

of the Republic of Indonesia established *Hiyung* cayenne pepper as a national variety. This cayenne pepper can support national chili production because of its high productivity with good market prospects, growing both on swampy and dry land. Compared to the three commercial varieties (Sonar, Bara, and Santika) *Hiyung* cayenne pepper has the highest dry weight, and higher productivity with a longer harvest duration, with the highest capsaicin levels about 699 ppm (Pramudyani, Sabran, & Noor, 2019).

Plant disease is one of the factors of failure in harvesting, including in chili plants (Islam, Schreinemachers, & Kumar, 2020). Species of fungi that can infect chili plants include *Colletotrichum scovillei* (Caires et al., 2014), *Aspergillus spp.*,

5
 ISSN: 0126-0537 Accredited First Grade by Ministry of Research, Technology and Higher Education of The Republic of Indonesia, Decree No: 30/E/KPT/2018

Cite this as: Imaningsih, W., Mariana, Junaedi, A. B., & Rasyidah. (2021). Antifungal activities of the combination of ulin wood liquid smoke and *Hiyung* cayenne pepper root endophyte fungi against *Colletotrichum capsici*. *AGRIVITA Journal of Agricultural Science*, 43(1), 69–78. <https://doi.org/10.17503/agrivita.v1i1.2458>

Fusarium spp., *Colletotrichum* spp. (Frimpong et al., 2019) 8 genera, and 17 species were identified on the basis of morphology, culture characteristics, and DNA sequencing of the internal transcribed spacer (ITS), and many more fungi have been reported. Fungi that cause disease in plants often cause structural and physical damage (Marques, Soares, & Apezato-Da-Gloria, 2013). Safe and environmentally friendly control of pathogenic fungi can be done by utilizing microorganisms derived from the plant itself, such as endophyte microbes (Köhl, Postma, Nicot, Ruocco, & Blum, 2011).

Endophytic microbes can derive from bacteria and fungi groups that have the ability to create a colony, part or all of its life cycle on a plant network without harming its host (Köhl, Postma, Nicot, Ruocco, & Blum, 2011; Selim, El-Beih, AbdEl-Rahman, & El-Diwany, 2012). Endophytic fungi are found in the plant tissue system, including flowers, twigs, leaves, and plant roots. These microorganisms grow and take food from the plant, and can infect healthy crops in certain tissues as well as able to produce mycotoxin, enzymes, and antibiotics (Stone, Polishook, & White Jr, 2004). Endophyte fungi also can inhibit pathogenic microbes that cause plant disease through mechanisms i.e. space and nutrition competitions and producing bioactive compounds such as antibacterial and antifungal (Gao, Dai, & Liu, 2010). Fungi also have the ability to produce plant growth hormones such as auxin (Imaningsih, Kadarsah, & Rusmannurrachmad, 2019) and herbicidal activity (de Souza et al., 2017).

Control of pathogenic fungi as a plant destruction organism can also be done by utilizing liquid smoke vapor condensate of pyrolysis of wood containing the main compounds of acids, phenols, and carbonyl (Lee et al., 2011). The constituent components of the liquid smoke compound have the ability to inhibit the growth of fungi and bacteria (Nami Kartal, Terzi, Kose, Hofmeyr, & Imamura, 2011; Okutucu, Duman, Ucar, Yasa, & Yanik, 2011). *Ulin* wood (*Eusideroxylon zwageri* Teijsm. & Binn.) liquid smoke contains acids and phenol compounds with acidity levels up to pH 2.08 (Junaidi, Apriyati, Abdullah, & Santoso, 2019) so that it has the potential to inhibit the growth of microbes.

Hiyung cayenne pepper originated from the village of Hiyung Tapin Regency of South Kalimantan is a natural resource that must be guarded, therefore it is necessary to do an exploration of endophytes potential as an antimicrobial agent for

pathogenic fungi. The potency of endophytes as an antimicrobial to pathogenic mold is known for its mechanisms including bioactive compounds as antibacterial and antifungal. Acetic acid and phenol which are the main constituent compounds of liquid smoke have the ability as antimicrobial. Endophytic molds and liquid smoke have the same ability to act as antimicrobials, but their potency is unknown when combined between the two. This research was intended to study the ability of endophytic fungi and liquid smoke to inhibit the growth of pathogenic fungi and examine the ability of endophytic fungi added by a combination of liquid smoke at different concentrations to the growth of pathogenic fungi.

MATERIALS AND METHODS

Isolation and Purification of Endophytic Fungi

This research was conducted from November 2018 to April 2019 at the Microbiology Laboratory of the Faculty of Mathematics and Natural Sciences, Lambung Mangkurat University, Banjarbaru, South Kalimantan. Samples of chili cayenne pepper plants were taken from the village of Hiyung, Tapin District, South Kalimantan in November 2018. The chili plants that were taken were put in polybags and coded as sampling information. Isolation and purification of endophytic fungus were carried out to obtain pure isolates by direct planting methods. The procedure is performed based on the method used by Septiana, Sukarno, Sukarno, & Simanjuntak (2017) with modification.

Identification and Screening of Endophytic Fungi

The fungi Identification includes macroscopic and microscopic observation. The macroscopic observation of pure isolates was done by modification of the method of Gerardo-Lugo et al. (2020) performed by observing the shape, color, and diameter of the colony for 7 days, the color and presence of the Hypha region determined on the 7th day. Microscopic observation was done by the method of slide culture (Rosana, Matsuzawa, Gono, & Karuniawa, 2014). Isolation identification refers to the Fungi identification book: *Illustrated Genera of Imperfect Fungi* (Barnett & Hunter, 1998) and *The Genera of Hyphomycetes* (Seifert & Gams, 2011).

Screening of endophytic fungi is conducted through the test of pathogenicity and antagonism. The pathogenicity test was carried out against the endophyte isolates that were derived from the roots of healthy chili plants based on the methods done

Witiyasti Imaningsih et al.: Disease Protection of Hiyung Cayenne Pepper

by Soesanto, Hartono, Mugiastuti, & Widarta (2020). Percent germination is obtained by dividing the number of germinated seeds (normal/abnormal/not growing) by the total number of seeds multiplied by 100%. The antagonism test refers to the dual culture method in the PDA medium (Tomah, Abd Alamer, Li, & Zhang, 2020). The inhibitory percent was obtained by dividing the difference from the diameter of the control pathogen colony and the diameter of the treatment pathogen colony by the diameter of the control pathogen multiplied by 100% (Kunova et al., 2016).

Production of Endophytic Fungus Filtrate

The endophytic fungus was prepared using the method of Qiao, Ling, Yu, Huang, & Wang (2017) that has been modified. The endophytic fungus was inoculated in a slanted PDA medium incubated for 7 days, then harvested by adding 9 ml of sterile distilled water to a tube containing endophytic fungi. The culture was homogeneous using a soft brush to obtain a spore suspension, then the suspension was transferred to another sterile test tube. Fungus suspension is centrifuged at a speed of 3000 rpm for 20 minutes to get the filtrate supernatant. The results of the process were separated using a 0.45 µm syringe filter and used as a crude biocontrol agent for further testing.

Inhibition Ability Test of Liquid Smoke Endophytic Fungi on *C. capsici*

Test the ability to inhibit liquid smoke is carried out by the agar dilution method (Balouri, Sadiki, & Ibsouda, 2016). The liquid smoke used in this study came from Ulin wood (*Eusideroxylon zwageri* Teijsm. & Binn.) obtained from the condensation of Ulin charcoal smoke production in Ranggang Village, Takisung District, Tanah Laut Regency, South Kalimantan. This method is carried out by mixing the media with liquid smoke, and the pathogen *C. capsici* is inoculated on the media that has been mixed with liquid smoke. The concentration used in this test was 0.00%; 0.085%; 0.17%; 0.34%; 0.68%; 1.36%; 1.75%. Observations were made from day 1 to day 7 after inoculation. The minimum inhibitory concentration was determined by the dilution method, using the method of Venkateswarulu et al. (2018) that has been modified. The concentration of endophytic fungus used in this test was 0%; 2%; 4%; 6%; 8%; 10%. Positive control treatment using ketoconazole 200 mg a concentration of 2%. Pathogenic mold pieces 6 mm inoculated in the middle of the PDA

medium positive control treatment and concentration treatment of endophytic fungus filtrate, then incubated at 28°C for 7 days, and observations were made every day. The minimum inhibitory concentration is determined from the presence or absence of pathogenic fungus growth at the lowest concentration of endophytic fungus filtrate.

An inhibitory test of the combination of endophytic fungus and liquid smoke was carried out based on the method carried out by Balouri, Sadiki, & Ibsouda (2016) the combination of endophytic fungus filtrate (EFF) and liquid smoke (LS) was 2% for filtrate and 0.68%, 1.36% for liquid smoke. So the first combination is 2% (EFF) + 0.68% (LS) and 2% (EFF) + 1.36% (EFF).

A 6 mm pathogenic fungus is inoculated in the middle of the PDA medium in a Petri dish. Observation of inhibition of pathogenic fungus was carried out from day 1 to day 7 after inoculation. The inhibitory effect was obtained by dividing the difference between the diameter of the control pathogen colony and the diameter of the treatment pathogen colony by the diameter of the control pathogen multiplied by 100% (Kunova et al., 2016).

Data Analysis

Analysis of the data using One-Way ANOVA at alpha 0.05 was followed by Duncan's test, or when the data is not homogeneous, the Kruskal Wallis test is used. All data were analyzed using IBM SPSS Statistic Version 22, 2013.

RESULTS AND DISCUSSION

Diversity of Endophytic Fungi from Roots of The Hiyung Cayenne Pepper

Isolation and purification of endophytic fungi derived from chili cayenne root were obtained by 8 isolates and given a code name using a sample source that is chili root (ACH). Isolates obtained from chili cayenne roots were characterized by macroscopically and microscopically. Pure isolates obtained were *Trichoderma* sp. ACH1.1, *Trichoderma* sp. ACH1.6, *Trichoderma* sp. ACH2.2, *Botrytis* sp. ACH2.3, *Gliocladium* sp. ACH2.4, *Harmoniella* sp. ACH2.5, *Humicola* sp. ACH2.6, *Cunninghamella* sp. ACH2.7.

The factors affecting the presence of endophytic fungi include the environment and the plant tissue used (Maheswari & Rajagopal, 2013). The Habitat of plant origin is one of the environmental factors affecting the structure and type of microbes that colonize the plant tissue such as roots, stems,

leaves, and branches (Araújo et al., 2002). Sieber & Grünig (2006) mentions that affecting the diversity of endophytes derived from plants is environmental factors, vegetation, and interactions with other types of microbes.

Pathogenicity and Antagonism of Endophytic Fungi from the Roots of Hiyung Cayenne Pepper

Screening of endophytic fungi was carried out by pathogenicity and antagonistic tests. The pathogenicity test was carried out on pure isolates resulting from the isolation of endophytic fungi. The percentage rate of pepper seed germination on the 7 and 14th day after inoculation (DAI) in the pathogenicity test is presented in Table 1. The fungus penetrates its host through mechanical and enzymatic mechanisms (Ashry & Mohamed, 2012). Fungi penetrate the epidermis, the cuticles, and cell walls (Underwood & Somerville, 2008). Enzymes that act as degenerating cell walls are pectinase and cellulase, where these enzymes are used for fungus for the process of penetration and colonization of host plants (Ashry & Mohamed, 2012; Kikot, Hours, & Alconada, 2009).

The antagonistic test was carried out on 3 selected isolates from the results of the pathogenicity test using the dual culture method. The results of antagonistic tests based on diameter measurements of pathogenic fungi colonies and inhibitory effect can be seen in Table 2. This antagonistic nature is consistent with the statements of De la Cruz-Quiroz, Roussos, Rodríguez-Herrera, Hernandez-Castillo, & Aguilar (2018) that fungus is grown side by side and has the ability to grow faster, then these fungi are able to occupy space and suppress the growth of their opponent's fungus. This antagonistic nature occurs because of the same needs as each fungus, nutrition, and growing needs.

Endophytic screening results showed isolates of *Cunninghamella* sp. ACH2.7 was selected as an isolate used for testing endophytic fungus filtrate and in combination using liquid smoke. This was obtained from the Kruskal Wallis test where the isolate had the highest normal growing percentage of sprouts, the lowest abnormal sprouts, and the lowest ungrown sprouts as 29 as the highest percent inhibition of pathogens (Table 1 and Table 2).

Table 1. The percentage rate of pepper seed germination on the 7 and 14th day after inoculation (DAI) in the pathogenicity test

| Endophytic fungi | Germination (%)* | | | | | |
|-----------------------------------|----------------------|---------------|---------------|-----------------------|--------------|---------------|
| | 7 th -DAI | | | 14 th -DAI | | |
| | Normal | Abnormal | Not grow | Normal | Abnormal | Not grow |
| Without endophytic fungi addition | 100±0c | 0±0a | 0±0a | 100±0c | 0±0a | 0±0a |
| <i>Trichoderma</i> sp. ACH1.1 | 0±0ab | 0±0a | 100±0ab | 30±51.96ab | 6.67±5.77a | 63.33±55.07ab |
| <i>Trichoderma</i> sp. ACH1.6 | 0±0ab | 0±0a | 100±0ab | 33.33±41.63ab | 10±10a | 56.67±45.09ab |
| <i>Trichoderma</i> sp. ACH2.2 | 0±0a | 23.33±20.82ab | 76.66±20.81ab | 0±0a | 60±30ab | 40±30ab |
| <i>Botrytis</i> sp. ACH2.3 | 0±0a | 16.67±28.86ab | 83.33±28.86ab | 0±0a | 30±26.46ab | 70±26.45ab |
| <i>Gliocladium</i> sp. ACH2.4 | 0±0a | 0±0a | 100±0b | 0±0a | 26.67±30.55a | 73.33±30.55b |
| <i>Harmoniella</i> sp. ACH2.5 | 0±0abc | 3.33±5.77a | 96.66±5.77ab | 90±0abc | 3.33±5.77a | 6.67±5.77ab |
| <i>Humicola</i> sp. ACH2.6 | 10±17.32abc | 0±0a | 90±17.32ab | 73.33±23.1abc | 13.33±11.55a | 13.33±11.54ab |
| <i>Cunninghamella</i> sp. ACH2.7 | 66.67±25.16bc | 0±0a | 33.33±25.16ab | 90±10bc | 0±0a | 10±10ab |

Remarks: * The number followed by the same letter is not significantly different based on Duncan ($\alpha=0.05$)

Witiyasti Imaningsih et al.: Disease Protection of Hiyung Cayenne Pepper

Table 2. Pathogenic fungi colony diameter and inhibitory effect of endophytic fungi against *C. capsici* 7th day after inoculation (antagonisms test result)

| Endophytic fungi | <i>Colletotrichum capsici</i> Diameter (mm) | Inhibitory effect (%)* |
|----------------------------------|---|------------------------|
| <i>Harmoniella</i> sp. ACH2.5 | 43.33±2.82 | 10.71±4.79 b |
| <i>Humicola</i> sp. ACH2.6 | 46.50±2.55 | 5.11±0.18 c |
| <i>Cunninghamella</i> sp. ACH2.7 | 32.23±1.50 | 33.26±2.54 a |

Remarks: * The number followed by the same letter is not significantly different based on the Kruskal Wallis Test ($\alpha=0.05$)





Fig. 1. Inhibitory effect (%) of liquid smoke against *C. capsici* on the 1-7th day after inoculation the bar indicates standard deviation between replicates. The concentration of liquid smoke followed by the same letter is not significantly different based on Duncan test ($\alpha=0.05$).

The Ability of Liquid Smoke and Endophytic Fungi Inhibits *C. capsici*

Based on the test results of liquid smoke ability inhibit the growth of *C. capsici*, obtained at all concentrations of liquid smoke is able to inhibit growth. The concentration of ironwood liquid smoke from 0.085-1.75% can inhibit *C. capsici* by 3.56-62.17%. But only at concentrations of 0.34%, 0.68%, 1.36%, and 1.75% showed significant inhibition compared to control (0.00% liquid smoke concentration). Inhibitory effect (%) of liquid smoke against *C. capsici* on the 1-7th day after inoculation is shown in Fig. 1. The inhibition effect was significantly different between treatments ($F=11.053$, $P=0.000$). Ironwood liquid smoke inhibits *C. capsici* causing

changes in the diameter of the colony. The increasing concentration of liquid smoke the smaller the colony that forms (Fig. 2). The condition occurs because the content of liquid smoke affects the growth of fungi. Phenol and acid compounds in liquid smoke can damage the structure of the fungus. This is in line with research of Suresh et al. (2019) which uses pyroligneous acid from a mixture of several kinds of wood that has the ability to inhibit the growth of fungi. The acid content of liquid smoke causes acid conditions in the cytoplasm, therefore causing damage to membrane surface tension and loss of active transport, resulting in unstable function and structure of cell components (Hassan, Sand, & El-Kadi, 2012).



25 2. Visual comparison of the growth of *C. capsici* in the addition of liquid smoke with a concentration of (A) 0.00% (B) 0.085% (C) 0.17% (D) 0.34% (E) 0.68% (F) 1.36% (G) 1.75% of the 7th day of incubation.

The best concentration of liquid smoke is then selected for the combination test. Endophytic fungal filtrate tests selected from previous tests were also conducted to determine the minimum inhibitory concentration (MIC) to be used to test the combination of endophytic fungal filtrate and liquid smoke. Based on the test results, endophyte fungi filtrate can inhibit the growth of *C. capsici*. 24 results of the inhibition of endophilic fungus filtrate can be seen in Fig. 3. Inhibitory effects differ significantly between treatments ($F = 41.634$, $P = 0.000$). All concentration treatments differ significantly from control (ketoconazole 2%). Even at a concentration of 2%, *Cunninghamella* sp. ACH2.7 filtrate is already able to inhibit *C. capsici* (91.69%), even better when compared to ketoconazole 2% 200 mg as a positive control.

In addition to inhibiting growth, isolate *Cunninghamella* sp. ACH2.7, also causes the morphology of *C. capsici* to change. One of them is a colony that was originally blackish gray to be lighter gray-white on PDA medium. These discoloration and morphology are likely caused by the ability of compounds produced by endophytic fungi to damage the structure of *C. capsici*, this requires further research. De la Cruz-Quiroz, Roussos, Rodríguez-Herrera, Hernandez-Castillo, & Aguilar (2018) in research on the ability of *Trichoderma* to inhibit *P.capsici* states that the pathogen cell part is used as a source of nutrition for the growth of *Trichoderma*, the same is likely to happen in this study.

The Synergistic of Endophytic Mold and Liquid Smoke Filtrate Inhibits *C. capsici*

Based on the previous test results, 2

17 concentrations of ironwood liquid smoke (LS) were used (0.68% and 1.36%) and MIC concentration of filtrate *Cunninghamella* sp. ACH27(2%) (EFF) to be combined so that the synergy can be known. The combination of 0.68% LS and 2% EFF, as well as 1.36% LS and 2% EFF is able to inhibit the growth of *C. capsici* differs significantly with 2% ketoconazole as control ($F=14.676$, $P=0.000$) (Fig. 4). The utilization of a combination of liquid smoke and endophytic fungus filtrate as an antimicrobial can be developed with the use of the right concentration because both of these ingredients have active compounds that can be utilized. Liquid smoke and endophyte fungi have the same benefits that act as antibacterial and antifungal. In line with Aisyah, Sinaga, Nawangsih, Giyanto, & Pari (2018) that liquid smoke has the ability as an antibacterial agent and stimulates plant growth, in research on liquid smoke from some wood to overcome the banana disease. Gunatilaka (2006) also stated that endophyte fungi can produce secondary metabolites and compounds that act as an antifungal and antibacterial agent.

The combination of liquid smoke and the endophytic filtrate is able to inhibit growth and cause the morphology of *C. capsici* to change. Morphological changes in *C. capsici* due to the presence of active compounds that can damage cell membranes. Phenols from liquid smoke and flavonoids from endophytic fungi can improve the permeability of cell membranes, as well as inhibit the activation of essential enzymes, and the functioning of genetic materials (Konaté et al., 2012).

Witiyasti Imaningsih et al.: Disease Protection of Hiyung Cayenne Pepper

The treatment of a combination of liquid smoke and endophytic fungi filtrate in vitro is able to inhibit the growth of *C. capsici*, so from these results, there is the potential that the combination of those two compounds can be antimicrobial agents that can be used to control disease-causing

pathogens in plants. Utilization of a combination of liquid smoke and endophytic fungi filtrate as antimicrobial agents can be developed with the use of appropriate concentrations because both of these materials have active compounds.




Fig. 3. Inhibitory effect (%) of *Cunninghamella* sp. ACH2.7 filtrate against *C. capsici* on the 1-7th day after inoculation the bar indicates standard deviation between replicates. The concentration of crude extract followed by the same letter is not significantly different based on Duncan test ($\alpha=0.05$)




Fig. 4. Inhibitory effect (%) of the combination of Liquid Smoke (LS) and *Cunninghamella* sp. ACH2.7 filtrate/Endophyte Fungi Filtrate (EEF) against *C. capsici* on the 1-7th day after inoculation the bar indicates standard deviation between replicates. The concentration of crude extract followed by the same letter is not significantly different based on Duncan test ($\alpha=0.05$)

CONCLUSION

The combination of liquid smoke and endophytic fungi filtrate has the ability to inhibit *C. capsici*, and has the potential to be antimicrobial. The highest inhibitory power was generated in a combination of 1.36% liquid smoke and 2% endophytic fungi filtrate at 88.08%. The benefits of the combination of liquid smoke and endophytic fungi filtrate need further research, especially its potential as antimicrobial and appropriate concentration so that it can be utilized to control pathogens that can damage crops especially *Hiyung* cayenne pepper.

11

ACKNOWLEDGEMENT

This study was funded by the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia in 2018-2019. The authors would like to thank the Mr. Junaidi Secretary of Hiyung Village and farmer group "Karya Baru" Hiyung Village, Tapin Tengah District, Tapin Regency, South Kalimantan. And also sincere thanks conveyed to the chairman of local production business group "TALASIANA" at Ranggung Village, Takisung District, Tanah Laut Regency, South Kalimantan.

REFERENCES

- Aisyah, I., Sinaga, M. S., Nawangsih, A. A., Giyanto, & Pari, G. (2018). Utilization of liquid smoke to suppress blood diseases on bananas and its effects on the plant growth. *AGRIVITA Journal of Agricultural Science*, 40(3), 453–460. <https://doi.org/10.17503/agrivita.v40i3.1390>
- Araújo, W. L., Marcon, J., Maccheroni Jr, W., van Elsas, J. D., van Vuurde, J. W. L., & Azevedo, J. L. (2002). Diversity of endophytic bacterial populations and their interaction with *Xylella fastidiosa* in citrus plants. *Applied and Environmental Microbiology*, 68(10), 4906–4914. <https://doi.org/10.1128/aem.68.10.4906-4914.2002>
- Ashry, N. A., & Mohamed, H. I. (2012). Impact of secondary metabolites and related enzymes in flax resistance and/or susceptibility to powdery mildew. *African Journal of Biotechnology*, 11(5), 1073–1077. Retrieved from <https://www.ajol.info/index.php/ajb/article/view/100252>
- Balai Penelitian dan Pengembangan Pertanian. (2018). *Cabai Hiyung, SDG Lokal Kalimantan Selatan*. Retrieved from <http://www.litbang.pertanian.go.id/info-teknologi/3152/>
- Balouiri, M., Sadiki, M., & Ibsouda, S. K. (2016). Methods for *in vitro* evaluating antimicrobial activity: A review. *Journal of Pharmaceutical Analysis*, 6(2), 71–79. <https://doi.org/10.1016/j.jpha.2015.11.005>
- Barnett, H. L., & Hunter, B. B. (1998). *Illustrated genera of imperfect fungi* (4th ed.). St. Paul, USA: American Phytopathological Society (APS Press). Retrieved from https://www.academia.edu/35499449/illustrated_genera_of_imperfect_fungi_fourth_edition_Barnett_y_Hunter_pdf
- Caires, N. P., Pinho, D. B., Souza, J., Silva, M. A., Lisboa, D. O., Pereira, O. L., & Furtado, G. Q. (2014). First report of anthracnose on pepper fruit caused by *Colletotrichum scovillei* in Brazil. *Plant Disease*, 98(10), 1437. <https://doi.org/10.1094/PDIS-04-14-0426-PDN>
- De la Cruz-Quiroz, R., Roussos, S., Rodríguez-Herrera, R., Hernández-Castillo, D., & Aguilar, C. N. (2018). Growth inhibition of *Colletotrichum gloeosporioides* and *Phytophthora capsici* by native Mexican *Trichoderma* strains. *Karbala International Journal of Modern Science*, 4(2), 237–243. <https://doi.org/10.1016/j.kijoms.2018.03.002>
- de Souza, A. R. C., Baldoni, D. B., Lima, J., Porto, V., Marcuz, C., Machado, C., ... Mazutti, M. A. (2017). Selection, isolation, and identification of fungi for bioherbicide production. *Brazilian Journal of Microbiology*, 48(1), 101–108. <https://doi.org/10.1016/j.bjm.2016.09.004>
- Frimpong, G. K., Adekunle, A. A., Ogundipe, O. T., Solanki, M. K., Sadhasivam, S., & Sionov, E. (2019). Identification and toxigenic potential of fungi isolated from capsicum peppers. *Microorganisms*, 7(9), 1–10. <https://doi.org/10.3390/microorganisms7090303>
- Gao, F., Dai, C., & Liu, X. (2010). Mechanisms of fungal endophytes in plant protection against pathogens. *African Journal of Microbiology Research*, 4(13), 1346–1351. Retrieved from [https://academicjournals.org/article/article1380280021_Gao et al.pdf](https://academicjournals.org/article/article1380280021_Gao%20et%20al.pdf)
- Gerardo-Lugo, S. S., Tovar-Pedraza, J. M., Maharachchikumbura, S. S. N., Apodaca-Sánchez, M. A., Correia, K. C., Saucedo-Acosta, C. P., ... Beltrán-Peña, H. (2020). Characterization of *Neopestalotiopsis* species associated with mango grey leaf spot disease in Sinaloa, Mexico. *Pathogens*, 9(10), 788. <https://doi.org/10.3390/pathogens9100788>
- Gunatilaka, A. A. L. (2006). Natural products from plant-associated microorganisms: Distribution, structural diversity, bioactivity, and implications

Witiyasti Imaningsih *et al.*: Disease Protection of Hiyung Cayenne Pepper

- of their occurrence. *Journal of Natural Products*, 69(3), 509–526. <https://doi.org/10.1021/np058128n>
- Hassan, R. A., Sand, M. I., & El-Kadi, S. M. (2012). Effect of some organic acids on fungal growth and their toxins production. *Journal of Agricultural Chemistry and Biotechnology*, 3(9), 391–397. <https://doi.org/10.21608/jacb.2012.55011>
- Imaningsih, W., Kadarsah, A., & Rusmannurrachmad, R. D. T. (2019). The capability of consortium phosphate solubilizing bacteria and IAA producing fungi on promoting elephant grass growth. *Jurnal Biodjati*, 4(1), 138–148. <https://doi.org/10.15575/biodjati.v4i1.4284>
- Islam, A. H. M. S., Schreinemachers, P., & Kumar, S. (2020). Farmers' knowledge, perceptions and management of chili pepper anthracnose disease in Bangladesh. *Crop Protection*, 133, 105139. <https://doi.org/10.1016/j.cropro.2020.105139>
- Junaidi, A. B., Apriyani, H., Abdullah, & Santoso, U. T. (2019). Fraksinasi dan karakterisasi asap cair dari kayu ulin (*Eusideroxylon zwageri* Teijsm. & Binn.) sebagai pelarut kitosan. *Jurnal Riset Industri Hasil Hutan*, 11(2), 53–64. <https://doi.org/10.24111/jrihh.v11i2.4861>
- Kikot, G. E., Hours, R. A., & Alconada, T. M. (2009). Contribution of cell wall degrading enzymes to pathogenesis of *Fusarium graminearum*: A review. *Journal of Basic Microbiology*, 49(3), 231–241. <https://doi.org/10.1002/jobm.200800231>
- Köhl, J., Postma, J., Nicot, P., Ruocco, M., & Blum, B. (2011). Stepwise screening of microorganisms for commercial use in biological control of plant-pathogenic fungi and bacteria. *Biological Control*, 57(1), 1–12. <https://doi.org/10.1016/j.biocontrol.2010.12.004>
- Konaté, K., Hilou, A., Mavoungou, J. F., Lepengué, A. N., Souza, A., Barro, N., ... Nacoulma, O. G. (2012). Antimicrobial activity of polyphenol-rich fractions from *Sida alba* L. (Malvaceae) against co-trimoxazol-resistant bacteria strains. *Annals of Clinical Microbiology and Antimicrobials*, 11, 5. <https://doi.org/10.1186/1476-0711-11-5>
- Kunova, A., Bonaldi, M., Saracchi, M., Pizzatti, C., Chen, X., & Cortesi, P. (2016). Selection of *Streptomyces* against soil borne fungal pathogens by a standardized dual culture assay and evaluation of their effects on seed germination and plant growth. *BMC Microbiology*, 16(1), 272. <https://doi.org/10.1186/s12866-016-0886-1>
- Lee, S. H., H'ng, P. S., Chow, M. J., Sajap, A. S., Tey, B. T., Salmiah, U., & Sun, Y. L. (2011). Effectiveness of pyroligneous acids from vapour released in charcoal industry against biodegradable agent under laboratory condition. *Journal of Applied Sciences*, 11(24), 3848–3853. <https://doi.org/10.3923/jas.2011.3848.3853>
- Maheswari, S., & Rajagopal, K. (2013). Biodiversity of endophytic fungi in *Kigelia pinnata* during two different seasons. *Current Science*, 104(4), 515–518. Retrieved from <https://www.jstor.org/stable/24089753?seq=1>
- Marques, J. P. R., Soares, M. K. M., & Appezzato-Da-Gloria, B. (2013). New staining technique for fungal-infected plant tissues. *Turkish Journal of Botany*, 37, 1–4. <https://doi.org/10.3906/1204-9>
- Nami Kartal, S., Terzi, E., Kose, C., Hofmeyr, J., & Imamura, Y. (2011). Efficacy of tar oil recovered during slow pyrolysis of macadamia nut shells. *International Biodeterioration and Biodegradation*, 65(2), 369–373. <https://doi.org/10.1016/j.ibiod.2010.08.011>
- Okutucu, C., Duman, G., Ucar, S., Yasa, I., & Yanik, J. (2011). Production of fungicidal oil and activated carbon from pistachio shell. *Journal of Analytical and Applied Pyrolysis*, 91(1), 140–146. <https://doi.org/10.1016/j.jaap.2011.02.002>
- Pramudiyani, L., Sabran, M., & Noor, A. (2019). Agronomic performance and nutrition content of hiyung as local variety of cayenne pepper (*Capsicum frutescens*) at dry land and swamp land of South Kalimantan Province. *Buletin Plasma Nutfah*, 25(1), 43-52. <https://doi.org/10.21082/blpn.v25n1.2019.p43-52>
- Qiao, W., Ling, F., Yu, L., Huang, Y., & Wang, T. (2017). Enhancing taxol production in a novel endophytic fungus, *Aspergillus aculeatinus* Tax-6, isolated from *Taxus chinensis* var. *mairei*. *Fungal Biology*, 121(12), 1037–1044. <https://doi.org/10.1016/j.funbio.2017.08.011>
- Rosana, Y., Matsuzawa, T., Gonoi, T., & Karuniawati, A. (2014). Modified slide culture method for faster and easier identification of dermatophytes. *Microbiology Indonesia*, 8(3), 7. <https://doi.org/10.5454/imi.8.3.7>
- Seifert, K. A., & Gams, W. (2011). The genera of Hyphomycetes - 2011 update. *Persoonia: Molecular Phylogeny and Evolution of Fungi*, 27, 119–129. <https://doi.org/10.3767/003158511X617435>

Witiyasti Imaningsih *et al.*: Disease Protection of Hiyung Cayenne Pepper

- Selim, K., El-Beih, A., AbdEl-Rahman, T., & El-Diwany, A. (2012). Biology of endophytic fungi. *Current Research in Environmental & Applied Mycology*, 2(1), 31–82. <https://doi.org/10.5943/cream/2/1/3>
- Septiana, E., Sukarno, N., Sukarno, & Simanjuntak, P. (2017). Endophytic fungi associated with turmeric (*Curcuma longa* L.) can inhibit histamine-forming bacteria in fish. *HAYATI Journal of Biosciences*, 24(1), 46–52. <https://doi.org/10.1016/j.hjb.2017.05.004>
- Sieber, T. N., & Grünig, C. R. (2006). Biodiversity of fungal root-endophyte communities and populations, in particular of the dark septate endophyte *Phialocephala fortinii* s. l. In B. J. E. Schulz, C. J. C. Boyle, & T. N. Sieber (Eds.), *Microbial Root Endophytes* (pp. 107–132). Berlin, Heidelberg: Springer. https://doi.org/10.1007/3-540-33526-9_7
- Soesanto, L., Hartono, A. R. R., Mugiastuti, E., & Widarta, H. (2020). Seed-borne pathogenic fungi on some soybean varieties. *Biodiversitas*, 21(9), 4010–4015. <https://doi.org/10.13057/biodiv/d210911>
- Stone, J. K., Polishook, J. D., & White Jr, J. F. (2004). Endophytic fungi. In M. Foster & G. Bills (Eds.), *Biodiversity of Fungi: Inventory and Monitoring Methods* (1st ed., pp. 241–270). Elsevier Academic Press. <https://doi.org/10.13140/RG.2.1.2497.0726>
- Suresh, G., Pakdel, H., Rouissi, T., Brar, S. K., Fliss, I., & Roy, C. (2019). *In vitro* evaluation of antimicrobial efficacy of pyroligneous acid from softwood mixture. *Biotechnology Research and Innovation*, 3(1), 47–53. <https://doi.org/10.1016/j.biori.2019.02.004>
- Tomah, A. A., Abd Alamer, I. S., Li, B., & Zhang, J. Z. (2020). A new species of *Trichoderma* and gliotoxin role: A new observation in enhancing biocontrol potential of *T. virens* against *Phytophthora capsici* on chili pepper. *Biological Control*, 145, 104261. <https://doi.org/10.1016/j.biocontrol.2020.104261>
- Underwood, W., & Somerville, S. C. (2008). Focal accumulation of defences at sites of fungal pathogen attack. *Journal of Experimental Botany*, 59(13), 3501–3508. <https://doi.org/10.1093/jxb/ern205>
- Venkateswarulu, N., Shameer, S., Bramhachari, P. V., Basha, S. K. T., Nagaraju, C., & Vijaya, T. (2018). Isolation and characterization of plumbagin (5-hydroxyl-2-methylnaptalene-1,4-dione) producing endophytic fungi *Cladosporium delicatulum* from endemic medicinal plants: Isolation and characterization of plumbagin producing endophytic fungi from endemic medicinal plants. *Biotechnology Reports*, 20, e00282. <https://doi.org/10.1016/j.btre.2018.e00282>

Antifungal Activities of the Combination of Ulin Wood Liquid Smoke and Hiyung Cayenne Pepper Root Endophyte Fungi Against *Colletotrichum capsici*

ORIGINALITY REPORT

17%

SIMILARITY INDEX

14%

INTERNET SOURCES

12%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|---|---|----|
| 1 | iosrjournals.org Internet Source | 2% |
| 2 | Submitted to Universitas Brawijaya Student Paper | 1% |
| 3 | jurnal.fp.unila.ac.id Internet Source | 1% |
| 4 | 3lata.pl Internet Source | 1% |
| 5 | agrivita.ub.ac.id Internet Source | 1% |
| 6 | journal.ugm.ac.id Internet Source | 1% |
| 7 | innspub.net Internet Source | 1% |
| 8 | dokumen.pub Internet Source | 1% |

9

Internet Source

1 %

10

Dwi Ningsih Susilowati. "THE POTENTIAL OF ENDOPHYTIC FUNGI AS BIOCONTROL AND PHOSPATE SOLUBILIZATION AGENT IN *Capsicum anuum*", Fungal Territory, 2020

Publication

<1 %

11

journal.ipb.ac.id

Internet Source

<1 %

12

mail.scialert.net

Internet Source

<1 %

13

A Wachid, Sutarman. " Inhibitory Power Test of Two Isolates in In Vitro Way Againts The Cause of Red Chilli Stem Rot ", Journal of Physics: Conference Series, 2019

Publication

<1 %

14

N H Haryanti, H Wardhana, Suryajaya, R Ramadhan, Yunita. "Halaban (*Vitex Pubescens* vahl) charcoal and coal bottom ash briquettes with pressure variations", Journal of Physics: Conference Series, 2020

Publication

<1 %

15

Yuhan Zhou, Yuelian Liu, Qingfang Lv. "Suspensions Derived from Anaerobically Fermented Tilapia Offal Inhibit *Fusarium oxysporum* f. sp. *cubense* in Banana", Agricultural Sciences, 2018

<1 %

16

ojs3.unpatti.ac.id

Internet Source

<1 %

17

Suharso Suharso, Eka Setiososari, Agung Abadi Kiswandono, Buhani Buhani, Heri Satria. "Liquid smoke of coconut shell as green inhibitor of calcium carbonate scale formation", DESALINATION AND WATER TREATMENT, 2019

Publication

<1 %

18

"Endophytes and Secondary Metabolites", Springer Science and Business Media LLC, 2019

Publication

<1 %

19

twj.ulm.ac.id

Internet Source

<1 %

20

I Yuliana, H Khatimah, L Rosida, N S Skripsiana, E Suhartono. "Martapura river water leads to testes alteration in rats", Journal of Physics: Conference Series, 2019

Publication

<1 %

21

Charles R. Howell, Robert D. Stipanovic. "Gliovirin, a new antibiotic from , and its role in the biological control of ", Canadian Journal of Microbiology, 1983

Publication

<1 %

22

digilib.unimed.ac.id

<1 %

23

publikasiilmiah.ums.ac.id:8080

Internet Source

<1 %

24

"Industrial Crop Plants", Springer Science and Business Media LLC, 2024

Publication

<1 %

25

Jing Yao, Jiangying Zeng, Huinan Tang, Yirui Cheng, Jingjing Tan, Tianyuan Li, Xiangyu Li, Jiyan He, Yongjun Zhang. "Effect of deep eutectic solvent extraction on Auricularia auricula polysaccharide solubilization and antioxidant potential", Sustainable Chemistry and Pharmacy, 2023

Publication

<1 %

26

Maria del Refugio Castañeda Chávez, Violeta Pardo Seda, Erasmo Orrantia Borunda, Fabiola Lango Reynoso. "Influence of water temperature and salinity on seasonal occurrences of Vibrio cholerae and enteric bacteria in oyster-producing areas of Veracruz, México", Marine Pollution Bulletin, 2005

Publication

<1 %

27

Rebecca T. Indriawan, Edi Suryanto, Dewa G. Katja. "KOMPOSISI KIMIA ASAP CAIR DAGING BUAH PALA DAN EFEK KOMBINASI DENGAN

<1 %

SARI LEMON CUI TERHADAP *Streptococcus mutans*", CHEMISTRY PROGRESS, 2022

Publication

28

jerami.faperta.unand.ac.id

Internet Source

<1 %

29

jurnal.untan.ac.id

Internet Source

<1 %

30

pdffox.com

Internet Source

<1 %

31

real-j.mtak.hu

Internet Source

<1 %

32

revistabionatura.com

Internet Source

<1 %

33

www.scilit.net

Internet Source

<1 %

34

Alsayed Alfiky, Eliane Abou-Mansour, Mout De Vrieze, Floriane L'Haridon, Laure Weisskopf. " Newly Isolated spp. Show Multifaceted Biocontrol Strategies to Inhibit Potato Late Blight Causal Agent both In Vitro and In Planta ", *Phytobiomes Journal*, 2024

Publication

<1 %

35

C. Martínez-Arias, D. Macaya-Sanz, J. Witzell, J. A. Martín. "Enhancement of *Populus alba* tolerance to *Venturia tremulae* upon inoculation with endophytes showing in vitro

<1 %

biocontrol potential", European Journal of Plant Pathology, 2018

Publication

36

Liping Qiao, Yuzhen Jiao, Xiaokui Li, Yafei Zhang, Laifeng Lu, Xi Zhang, Xia Liu. "Herbal smoke fumigation for controlling *Penicillium crustosum* in fresh walnuts", Food Research International, 2023

Publication

<1 %

37

R. Thilagam, N. Hemalatha. "Plant growth promotion and chilli anthracnose disease suppression ability of rhizosphere soil actinobacteria", Journal of Applied Microbiology, 2019

Publication

<1 %

38

ann-clinmicrob.biomedcentral.com

Internet Source

<1 %

39

docksci.com

Internet Source

<1 %

40

jwoodscience.springeropen.com

Internet Source

<1 %

41

www.researchsquare.com

Internet Source

<1 %

42

Hasan Ashari Oramahi, Farah Diba, Juanita. "Anti-Termites Properties of Liquid Smoke from Bintangur Wood", Jurnal Sylva Lestari, 2021

<1 %

43

"Seed Endophytes", Springer Science and Business Media LLC, 2019

Publication

<1 %

44

S. Nami Kartal, Evren Terzi, Coskun Kose, John Hofmeyr, Yuji Imamura. "Efficacy of tar oil recovered during slow pyrolysis of macadamia nut shells", International Biodeterioration & Biodegradation, 2011

Publication

<1 %

45

Ting-ting Li, Jing-di Zhang, Jia-quan Tang, Zhi-cheng Liu, Ya-qian Li, Jie Chen, Li-wen Zou. " Combined Use of CCTCCSBW0199 and Brassinolide to Control Infection in Tomato ", Plant Disease, 2020

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On