

# BIOCONTROL FOR RHIZOCTONIA STEM ROT DISEASE BY USING COMBINATION OF SPECIFIC ENDOPHYTE IN PADDY TIDAL SWAMP

*by Mariana -*

---

**Submission date:** 18-Jun-2024 11:17AM (UTC+0700)

**Submission ID:** 2404595242

**File name:** Turnitin\_Agrivita\_2.pdf (253.49K)

**Word count:** 4390

**Character count:** 22738

## BIOCONTROL FOR RHIZOCTONIA STEM ROT DISEASE BY USING COMBINATION OF SPECIFIC ENDOPHYTE IN PADDY TIDAL SWAMP

4  
Ismed Setya Budi<sup>1)</sup> and Mariana

Faculty of Agriculture Lambung Mangkurat University, Banjarbaru, Indonesia  
Jl. A. Yani Po Box 1028, Banjarbaru 70714  
Corresponding author Phone: +62-81933753340 Email: isb\_unlam@yahoo.com

Received: January 12, 2012/ Accepted: September 29, 2013

### ABSTRACT

The use of combination of specific endophytic in tidal swamps to control stem root disease as biological control agents has not been done. It is expected that this combination is able to continuously protect plants from pathogen interference. The research was carried out in type C tidal swamp in Banjar regency of South Kalimantan, from March to November 2011, temperature 29-32°C, and pH 4.0-5.5. The method used was Split Plot design. Biocontrol preparation for both types of endophytic was applied in seeds in 7 days after planting (DAP). Observation on high intensity and plant diseases of planting stage on tidal swamps (taradak, ampak and lacak) was conducted. The results showed that there was a reduction of disease ranging from 58.70 to 87.29%. The application of combination of two biocontrol agents (*T. viride* PS-2.1 + *P. fluorescent* PS-4.8), (*Fusarium* non-pathogenic PS-1.5 + *P. fluorescent* PS-4.8) and (*T. viride* PS-2.1 + FNP PS-1.5) isolate gave the best inhibition result, reduced disease intensity, and increased plant height. The result of soil analysis before and after application of endophytic showed that there was an increase in soil fertility with the element addition of N, P, K and pH.

Keywords: stem rot, endophytic combination, paddy, tidal swamp

### INTRODUCTION

17  
*R. solani* is one of the most important phytopathogen which attacks rice cultivated on tidal swamps of South Kalimantan, causing stem root disease. *R. solani* intensity constantly increases because it is difficult to control flood (Budi and Mariana, 2012). Therefore, effective and

efficient controlling method safe for environment is required.

*R. solani* and other soil borne pathogens are controlled using synthetic, but the application is usually hampered by the coming tide which washes away the pesticide. On the other hand, planting local varieties in tidal swamps using three replacement stages can lead to inefficiency in pesticide use (Budi and Mariana, 2012). The use of specific biological agents should be done immediately because of consumers' demand for synthetic chemical free products.

According to Budi and Mariana (2009), in paddy antagonist isolation from three different types of tidal swamp (locally termed as: taradak, ampak and lacak) were founded 28 endophytic colonies consisting of 23 colonies of fungi and 4 colonies of bacteria. From the *in-vitro* test, there were 4 isolates having higher antagonistic potency, namely *Trichoderma viride* PS-2.1, *Fusarium* non-pathogenic PS-1.5, *Pseudomonas fluorescent* PS-4.8, and *Bacillus* sp PS-3.14.

On biological control, *R. solani* can be parasitized by mycoparasites such as *Gliocladium* spp., *Trichoderma* spp. and *Verticillium biguttatum* (Van den Boogert and Deacon, 1994). Fungus *V. biguttatum* is a mycoparasite with biological activity against the important plant pathogen *R. solani*. The biotrophic mycoparasite *V. biguttatum* was first isolated from sclerotia of *R. solani* on potato tubers in Netherlands (Jager and Velvis, 1988). This fungus is able to destroy sclerotia (Jager and Velvis, 1988) and hyphae (Van den Boogert and Deacon, 1994) of *R. solani*. When applied as a spore suspension, mycoparasite *V. biguttatum* has the ability to suppress stem canker and black scurf on potatoes (Jager and Velvis, 1988).

12  
According to Howell and Stipanovic (1995), the growth of *Rhizoctonia solani* on the cotton plant can be controlled by seed treatment

Accredited SK No.: 81/DIKTI/Kep/2011

11  
<http://dx.doi.org/10.17503/Agrivita-2013-35-3-p304-310>

using *Gliocladium virens*. Antagonists of non-pathogenic *Fusarium* strains isolated from suppressive soil have the capability of reducing the disruption caused by *Fusarium* wilt in some plant (Nel *et al.*, 2006). While the bacterium *Pseudomonas capacia*, *P. fluorescens* and *P. gladio* were also able to control the growth of *P. solanacearum* using wilt on tomatoes. Other bacteria such as *Bacillus mesentericus*, *B. megaterium*, *B. mycoides* and *Erwinia* sp. also act as biological control for wilt disease in several plants (Harman, 1999).

Combined antagonists might give a better result than a single antagonist beside its uncertain result due to influence from environmental condition and time of application (Yigit and Dikilitas, 2007). Use of *Trichoderma* spp. and *Gliocladium* sp. will provide optimal results in controlling soil borne and air borne pathogens when the application is done in low inoculum population (Cook and Baker, 1983).

The use of specific biological agents which have a coevolution will be able to stimulate the development of harmful rhizosphere microorganisms, and more than one antagonist is usually isolated in the field (Budi and Mariana, 2009). Therefore, selecting the best combination of antagonists that can protect plants better from various pathogenic disorders is needed.

This research was conducted to determine the efficacy combination of specific antagonist isolate in planting stage in tidal swamps for biological control of stem rot disease in paddy. It is expected that by using the correct combination of antagonist will give a better result.

## MATERIALS AND METHODS

Isolation and mass production of antagonist agents from plant samples were done by taking healthy plants as samples from the infested paddy planted areas in tidal swamp in Banjar regency of South Kalimantan; endophytic isolation was done on the root and the base of the plant (Budi dan Mariana, 2012). Isolation method was based on Homby methods (Fokkema *et al.*, 1959) where dilution plate method ( $10^{-4}$  to  $10^{-6}$ ) was applied. Isolation of *Pseudomonas fluorescens* group using selective media (Sands and Rovira, 1970). Each *Pseudomonas fluorescens* group was isolated then tested according to Dhingra and Sinclair method (1995).

Test on *in vitro* inhibition potency through the combination of endophytes towards *R. solani*

as well as the test measuring the suitability of antagonist combination were performed on potato dextrose agar (PDA) in a petri dish by growing isolates in pair. Then, measurements to figure out the inhibition potency were carried out by applying the formula of Fokkema (Fokkema *et al.*, 1959):

$$I = (r_1 - r_2) (r_1)^{-1} \times 100$$

Remark:

I is the percentage of inhibition;

r1 is the radius of A colony that grows in the opposite direction to B

r2 is the radius of A colony that grows in the direction of B

In vivo test measuring the hampering ability of antagonist isolate towards stem rot disease pathogen was conducted via split plot design in a farming land and a greenhouse on sterilized soil. Pathogen inoculation was performed when it was three-week-old by sprinkling formulated pathogens ( $40 \text{ kg} \cdot \text{ha}^{-1}$ ) in the soil around paddy stems. The application of antagonists was given one week before planting, and at the time of planting by soaking seeds for 24 hours into suspension with spore density of  $10^{-4}$  per ml. Observations were carried out the next three weeks in every planting stage on tidal swamps (locally termed as: taradak, ampak and lacak) by counting the number of plants that showed wilt or stem rot symptoms and measuring the height of the plant.

All treatments are given according to randomized block design (RBD) with five replications. To know the effect of the differences among the treatments, the median value of Duncan's test (DMRT) at 5% was performed.

## RESULTS AND DISCUSSION

The treatment performed could reduce the disease around 58.70 to 87.29% in comparison to control. The combination of *T. viride* PS - 2.1 + FNP PS - 1.5 represented the inhibition ability reaching up to 73.40%, and it reduced the disease intensity up to 79.67%. However, Combination of *T. viride* PS-2.1 + *P. fluorescens* PS-4.8 seemed to show better inhibition potency (78.00%) and disease intensity reduction (84.50%), while FNP PS-1.5 + *P. fluorescens* PS-4.8 managed to perform the best inhibition potency (82.00%) and disease intensity reduction potency (87.29%) (Table 1). But, the single isolate by Budi and Mariana

(2009) on inhibition ability pair test was known that all of the isolates had an ability to provide better inhibition towards *R. solani*, such as *Trichoderma viride* PS-2.1 (68.00%), *Fusarium* non-pathogenic PS-1.5 (65.98%), *Pseudomonas fluorescent* PS-4.8 (55.50%) and *Bacillus sp* PS-3.14 (40.25%). Thus the combined use of several antagonists will be able to increase the inhibition on the development of stem rot disease of rice. Guetsky *et al.* (2001) stated that the appropriate combination would be able to enhance antagonistic potency. Olivain *et al.* (2006) proved that the use of either combined antagonists or antagonist applied singly was capable of inhibiting the development of pathogens causing tomato root disease.

The results also showed it was effective to give the application in one week before planting and soak the seeds for 24 hours before planting in the antagonist suspension with a density of  $10^5$  spores/ml, where the combined application of *T. viride* PS-2.1, *P. fluorescens* + PS-4.8 could reduce attack intensity by 84.50%, while the combined application of FNP PS-1.5 + *P. fluorescens* PS-4.8 amounted to 87.29%, and *T. viride* PS-2.1 + FNP PS-1.5 reduced the attack intensity by 79.67%. However, the combined antagonists precisely reduced the capability of inhibiting disease progression (Table 1 and 2).

The results of this study are in line with Yigit and Dikilitas (2007), implying that the use of FNP + *P. fluorescens* and *T. harzianum* T-22 + *P. fluorescens* in the laboratory test showed the better potency in inhibiting the development

15

of *Fusarium oxysporum f. sp. lycopersici* Snyder and Hansen than singly applied antagonist. A combination of *Fusarium* isolates Fo47 + *P. fluorescent* strain C7 showed similar potency in inhibiting the development of *Fusarium* wilt on tomato. Combined application of *P. fluorescent* and *Bacillus subtilis* in sterilized soil test indicates the ability of the inhibition is much better than just one type of single application because the bacteria proved to work synergistically. Biocontrol efficacy arising from combined use of two biocontrol agents did not depend greatly on biocontrol mechanisms. Combined use with at least one competitive biocontrol agents resulted in better control than combined use of two mycoparasitic (Xu and Jeger, 2013).

The potency of *Trichoderma* spp was allegedly taken from many studies indicated the mechanism of action of enzymes such as  $\beta$ -1-3-glucanases and  $\beta$  1-4 glucanases. Those enzymes were capable of producing substances, spurring the development of resistance (induced resistance) and competing for space and nutrients of pathogens (Olivain *et al.*, 2006). Last but not least, this type of fungus was also capable of improving soil fertility, as proven by the growth of plant in the phase of taradak, ampak or lacak (Table 3). The presence of antagonistic fungi was able to suppress disease progression in addition their capability of supplying nutrients for plant growth (Altomare *et al.*, 1999; Hanson and Howell, 2004 and Harman, 2006).

Table 1. Percentage of inhibition in vitro test and Intensity of stem rot disease after application on sterilized soil in greenhouse

Treatments	In-vitro test Inhibition (%)	In-vivo test	
		Disease Intensity (%)	(Reduction) (%)
Control	00,00 <sup>a</sup>	78,7 <sup>a</sup>	00,00
<i>T. viride</i> PS-2.1	30,00 <sup>c</sup>	21,7 <sup>d</sup>	72,43
FNP PS-1.5	17,28 <sup>b</sup>	27,6 <sup>c</sup>	64,93
<i>P. fluorescens</i> PS-4.8	26,80 <sup>c</sup>	28,4 <sup>c</sup>	63,91
<i>Bacillus sp</i> PS-3.14	13,50 <sup>b</sup>	32,5 <sup>c</sup>	58,70
<i>T. viride</i> PS-2.1 + FNP PS-1.5	73,40 <sup>e</sup>	16,0 <sup>de</sup>	79,67
<i>T. viride</i> PS-2.1 + <i>P. fluorescens</i> PS-4.8	78,00 <sup>e</sup>	12,2 <sup>e</sup>	84,50
<i>T. viride</i> PS-2.1 + <i>Bacillus sp.</i> PS-3.14	25,00 <sup>c</sup>	24,8 <sup>d</sup>	68,49
FNP PS-1.5 + <i>P. fluorescens</i> PS-4.8	82,00 <sup>e</sup>	10,0 <sup>e</sup>	87,29
FNP PS-1.5 + <i>Bacillus sp.</i> PS-3.14	42,75 <sup>cd</sup>	31,5 <sup>c</sup>	59,97
<i>Bacillus sp.</i> PS-3.14 + <i>P. fluorescens</i> PS-4.8	50,00 <sup>d</sup>	29,8 <sup>c</sup>	62,13

Remarks: \*\*Within column, means followed by different letters are significantly different (P<0.01; LSD test).

Table 2. Stem rot disease intensity on some time treatments and antagonist combination in greenhouse

Formulation	Antagonist	Plant attacked		Plant height (cm)
	Application time	Disease intensity (%)	Reduction	
3 <i>T. viride</i> PS-2.1 + <i>P. fluorescen</i> PS-4.8	Control	85.14 a	0.00	28.20 a
	7 d before planting	25.18 b	70.43	37.64 b
	seed soaking	20.44 c	75.99	39.12 b
	7 d before planting + seed soaking	10.47 d	87.71	49.29 c
<i>T. viride</i> PS-2.1 + <i>Bacillus sp</i> PS-3.14	7 d before planting	31.20 b	63.35	30.25 ab
	seed soaking	27.25 b	67.99	39.00 b
	7 d before planting + seed soaking	20.50 c	75.92	45.25 c
	7 d before planting	18.39 c	78.41	38.56 b
2 FNP PS-1.5 + <i>P. fluorescen</i> PS-4.8	seed soaking	13.78 cd	83.82	31.92 ab
	7 d before planting + seed soaking	10.00 d	88.26	52.20 c
	7 d before planting	34.40 b	59.60	37.64 b
	seed soaking	31.00 b	63.59	39.50 b
1 FNP PS-1.5 + <i>Bacillus sp</i> PS-3.14	7 d before planting + seed soaking	22.47 c	73.61	45.70 c

Remarks: \*\*Within column, means followed by different letters are significantly different ( $P < 0.01$ ; LSD test).

FNP and *Trichoderma* sp. are similar in ability according to Benhamou *et al.* (2002) in a way that both of them could colonize the surface of plant roots quickly, protecting from any pathogenic attack. Almeida *et al.* (2007), however, agreed that those two were known for their role in producing chitinase,  $\beta$ -glucanase 1-3 and 1-4 glucosidase enzymes. Thomashow and Weller (1996) added that this ability was triggered by the presence of toxins, antibiotics and siderophores.

Various combination of antagonists proved to be able to improve fertility. This is in line with the test performed by Duijff *et al.* (1998) showing that there was synergism in the use of combination of *Pseudomonas fluorescens* WCS417 + FNP so as to inhibit the development of *Fusarium* wilt pathogen better than the antagonist given singly. The variability of control in different environmental conditions could be reduced by using two kinds of antagonist: *F. oxysporum* avirulen coupled with a group *Pseudomonas fluorescens* in controlling wilt disease in various crops. The combination of several bacterial isolates antagonistic to fungus *Trichoderma hamate* more effectively suppress disease caused by *R. solani* compared with any other antagonist isolates applied singly (Kwok *et al.*, 1987).

The intensity of disease which occurred during planting stage of cultivation (taradak, ampak

and lacak) revealed that there was significant difference compared with control. All treatment combinations give the ability to suppress the disease intensity more than 50%, the best treatment was 3 performed through the isolate combination of *T. viride* PS-2.1 + *P. fluorescen* PS-4.8 (Table 3).

The ability of the FNP in inhibiting progression of some diseases was pointed out by Weller (1988), in which plant defense mechanism was triggered by the availability of certain agents known as endophytic antagonists. Information previously presented by Pal and Gardener (2001) implies that endophyte as an induction agent of resistance (induced resistance) can stimulate plant resistance, which may be caused by the presence of certain chemicals, non-pathogenic microorganisms, or incompatible pathogenic or virulence.

The ability of endophytic fungus in inhibiting development of pathogen could be 4 performed with the presence of antibiosis mechanism, competition and microparasit. Ozbay and Newman (2004) proved that the fungus *Trichoderma* spp was capable of accumulating carbon monoxide in the competition for space and nutrients. *T. harzianum* strain T24 was even capable of producing cellulose in the form of 1,3-glucanase enzyme one hour after inoculation,

whereas in strains SC164, SC 167 and SC 168 tested in a greenhouse, *Trichoderma* was found capable of inhibiting the *R. solani* attacks on tomatoes because of accumulation in chitinase and highest glucanase enzyme.

Thus, it is obvious that the combination of endophyte, once released to root zone of plants, plays a role in endophytic microbial conservation in rhizosphere environment which is expected to contribute to the conservation in biological environment. Similarly, augmented crop land in the growth of *rhizosphere T. viride* caused the continuous resistance due to a number of fungal antagonists which was capable of protecting the plants from pathogenic attack which may cause wilt disease. The application of *T. viride* on banana plant was also capable of preventing wilt disease caused by *Fusarium sp.*

Based on the results of soil chemical analysis, it proved the effect of antagonist application of chemical elements in soil. There was a significant increase when compared to the plant before given any application. Longer period of application had a tendency to give more significant increase where the effect in taradak phase would be much lower than that of lacak.

The microbe enhanced nutrition and pH soil as shown in Table 4. This is because the

combination of fungi accelerated decomposition of organic material in soil. In addition, decomposition of compost may cause soil pH to become more alkaline, so the nutrient becomes available to plant. The increase of pH in soil could increase when added with glucose due to decarboxylation and amination of nitrogen compound in order to contribute to plant growth. According to Harman (2006), *Trichoderma* spp and plant symbionts are capable of enhancing the mechanism of resistance to root disease infected either directly or indirectly by changing the composition of microflora in roots. *Trichoderma* spp and *Gliocladium virens* produce chlamydospores, which actively grow as biocontrol fungus within 2 to 3 day period under no special aseptic condition. *G. virens* and *T. hamatum* with the weight of 1.5% (wt/wt) reduced damping-off disease on eggplant caused by *Rhizoctonia solani*. According to Harman (2006), antagonism involving cortex and epidermis of root can secrete bioactive molecules which lead to forming thallus cell wall of *Trichoderma*. Meandroteome plant transcript is able to spur plant resistance, increase nutrient absorption and plant growth.

Table 3. Disease intensity and plant height after application in each planting stage on type C tidal swamps in Banjar Regency of South Kalimantan

Treatments	Planting Stage on Tidal Swamps								
	Taradak			Ampak			Lacak		
	Symptom Intensity (%)	Reduction	Plant height (cm)	Symptom Intensity (%)	Reduction	Plant height (%)	Symptom Intensity (%)	Reduction	Plant height (%)
Control	33.50 c	0.00	15.25 a	42.00 c	0.00	47.20 a	55.25 c	0.00	93.50 a
<i>T. viride</i> PS-2.1 + <i>P. fluorescens</i> PS-4.8	12.50 a	62.69	36.00 c	10.40 a	75.24	69.50 c	16.00 a	71.04	145.50 c
<i>T. viride</i> PS-2.1 + <i>Bacillus sp</i> PS-3.14	20.25 b	39.50	20.25 b	25.42 b	39.48	56.25 b	20.18 b	63.48	130.75 b
FNP PS-1.5 + <i>P. fluorescens</i> PS-4.8	14.00 a	58.21	32.40 c	24.50 a	41.67	53.50 b	22.00 b	60.18	139.45 bc
FNP PS-1.5 + <i>Bacillus sp</i> PS-3.14	15.50 a	53.73	23.25 b	26.50 b	36.91	53.40 b	23.45 b	57.56	127.25 b

Remarks: \*\*Within column, means values followed by different letters are significantly different (P<0.01; LSD test).

Table 4. The results of chemical analysis of soil before and after application of endophyte combination in tidal swamps

Treatments	Result of soil nutrient analysis					
	Before treatment			After treatment		
	N	P	K	N	P	K
Control	0.237	0.018	0.352	0.201	0.010	0.394
<i>T. viride</i> + <i>P. fluorescens</i>	0.240	0.018	0.352	0.743	0.027	0.491
<i>T. viride</i> + <i>Bacillus</i> sp.	0.237	0.018	0.352	0.765	0.027	0.480
FNP + <i>P. fluorescens</i>	0.239	0.018	0.352	0.740	0.028	0.594
FNP + <i>Bacillus</i> sp.	0.237	0.021	0.352	0.680	0.027	0.431

### CONCLUSION

The combination of two specific endophytes in paddy tidal swamp showed that the inhibition ability is much better than singly applied endophyte. The best combination of two endophytes was *T. viride* PS-2.1 + *P. fluorescens* PS-4.8, *Fusarium* non-pathogenic PS-1.5 + *P. fluorescens* PS-4.8 and *T. viride* PS-2.1 + FNP PS-1.5 which reduced the intensity of stem root disease caused by *Rhizoctonia* sp, stimulated plant growth, soil pH, and soil fertility made of N, P and K which are available to plants.

### REFERENCES

- Alabouvette, C., P. Lemanceau and C. Steinberg. 1993. Recent advances in the biological control of *Fusarium* wilt. *Pestic. Sci.* 37:365-373
- Almeida, F. B. R., F. M. Cerqueira, R.N. Silva, C. J. Ulhoa, A.L. Lima. 2007. Mycoparasitism studies of *Trichoderma harzianum* strains against *Rhizoctonia solani*: evaluation of coiling and hydrolytic enzyme production. *Biotechnol Lett* 29: 1189–1193
- Altomare, C., W.A. Norvell, T. Björkmar and G.E. Harman. 1999. Solubilization of phosphates and micronutrients by the plant growth promoting and biocontrol fungus *Trichoderma harzianum* Rifai 1295-22. *Applied Environ. Microbiol.* 65: 2926-2933.
- Benhamou, N., C. Garand and A. Goulet. 2002. Ability of nonpathogenic *Fusarium oxysporum* Strain Fo47 to induce resistance against *Pythium ultimum* infection in cucumber. *Applied Environ. Microbiol.* 68: 4044-4060
- Budi, I.S. and Mariana. 2012. Bioecology of nonpathogenic *Fusarium* specific location of tidal swamps. *Proceeding International Seminar on University-Based Research for Wetland Development.* p.157-167.
- Budi, I.S. dan Mariana. 2009. Control on Wilt Disease on Paddy in Tidal Swamp Area, South Kalimantan by Using Antagonist and Botanic Pesticide. Faculty of Agriculture, Lambungmangkurat University, Banjarbaru
- Bollen, W.B. 1961. Interactions between pesticides and soil micro-organisms. *Ann. Rev. Microbiol.* 15, 69-92
- Cook, R.J. and K.F. Baker. 1983. The nature and practice of biological control of plant pathogens. APS Press, St. Paul, MN, USA.
- Duijff, B.J., D. Pouhair, C. Olivain, C. Alabouvette and P. Lemanceau. 1998. Implication of systemic induced resistance in the suppression of fusarium wilt of tomato by *Pseudomonas fluorescens* WCS417r and by nonpathogenic *Fusarium oxysporum* Fo47. *Eur. J. Plant Pathol.* 104: 903-910.
- Dhingra, O.D. and J.B. Sinclair. 1995. Basic plant pathology methods. Second edition. CRC Press, Inc., Boca Raton. pp.448
- Fokkema, N.J., J.H. Bond and H.A. Fribourg. 1959. Methods for studying soil microflora plant disease relationships. Burgess Publ. Co. USA. pp. 247.
- Fuchs, J.G., Y. Moenne-Loccoz and G. Defago. 1999. Nonpathogenic *Fusarium oxysporum* strain Fo47 to protect tomato against *Fusarium* wilt. *Bio. Control.* 4:105-110.

- Guetsky, R., D. Shtienberg, Y. Elad & A. Dinoor. 2001. Combining biocontrol Agents to reduce the variability of biological control. *Phytopathol* 91: 621-627.
- Howell, C.R. and Stipanovic. 1995. Mechanisms in the control of *Rhizoctonia solani* induced cotton seedling disease by *Gliocladium virens* antibiosis. *Phytopathol J.* 85: 469-472
- Hanson, L.E. and C.R. Howell. 2004. Elicitors of plant defense responses from biocontrol strains of *Trichoderma virens*. *Phytopathol.* 94:171-176.
- Harman G.E. 2006. Overview of mechanisms and uses of *Trichoderma* spp. *Phytopatol.* 96:190-194
- Jager G. and H. Velvis . 1988. Inactivation of sclerotia of *Rhizoctonia solani* on potato tubers by *Verticillium biguttatum*, a soil-borne mycoparasite. *Netherlands Journal of Plant Pathology* 94 (5) : 225-231
- Kwok, O.C.H., P.C. Gahy, H.A.J, Hoitink, and G.A. Kuter. 1987. Interactions between bacteria and *Trichoderma hamatum* in suppression of *Rhizoctonia* damping-off in bark compost media. *Phytopathology* 77: 1206-1212.
- Lewis., R.D. Lumsden and Locke. 1998. Biological control of damping-off caused by *Phytophthora ultimum* and *Rhizoctonia solani* with *Gliocladium virens* in soilless mix. *Phytopathology* 79:361-366
- Nel, B., C. Steinberg, N. Labuschagne and A. Viljoen. 2006. The potential of non-pathogenic *Fusarium oxysporum* and other biological control organisms for suppressing fusarium wilt of banana. *Plant Pathol. Journal*, 1(55): 217-223
- Olivain C., C. Humbert, J. Nahalkova, J. Fatehi, F. L'Haridon and C. Alabouvette. 2006. Colonization of tomato root by pathogenic and nonpathogenic *Fusarium oxysporum* strains inoculated together and separately into the soil. *Appl.Environ.Microbiol.*72 (2): 1523 - 1531
- Ozby, N and S. Newman. 2004. Biocontrol with *Trichoderma* spp with emphasis on *T. harzianum*. *Pakistan Journal of Biological Science* 7(4): 478-484
- Pal, K. K. and B. McSpadden Gardener, 2011. Biological control of plant pathogens. *The Plant Health Instructor* DOI: 10.1094/PHI-A-2006-1117-02. <https://www.apsnet.org/edcenter/advanced/topics/Pages/BiologicalControl.aspx>
- Sands, D.C. and A.D. Rovira. 1970. Isolation of *Pseudomonas fluorescens* with a selective medium. *Applied Microbiology* 20(3): 513-514.
- Thomashow, L.S. and D.M. Weller. 1996. Current concepts in the use of introduced bacteria for biological disease control: Mechanisms and antifungal metabolites. In: *Plant-Microbe Interactions*. Stacey, G., and N.T. Keen (Eds) New York: Chapman and Hill. p.236-271.
- Van Den Boogert, P. H. J. F. and J. W. Deacon. 1994. Biotrophic mycoparasitism by *Verticillium biguttatum* on *Rhizoctonia solani*. *Eur.J.Plant Pathol.*100(2):137-156.
- Van den Boogert P.H.J.F and Lutikholt A.J.G. 2004. Compatible biological and chemical control systems for *Rhizoctonia solani* in potato. *Euro. J. Plant Pathol.* 110: 111-118.
- Von Alten, H., A. Lindemann and F. Schönbeck. 1993. Stimulation of vesicular arbuscular mycorrhiza by fungicides or rhizosphere bacteria. *Mycorrhiza* 2 : 167-173
- Weller, D.M. 1988. Biological control of soilborne plant pathogens in the rhizosphere with bacteria. *Ann.Rev.Phytopathol.* 26: 379-407.
- Xu, X-M. and M.J. Jeger. 2013. Theoretical modelling suggests that synergy may result from combined use of two biocontrol agents for controlling foliar pathogens under spatial heterogeneous conditions. *Phytopathology*. 103(8): 768-775
- Xu, X-M., P. Jeffries, M. Pautasso, and M.J. Jeger. 2011. Combined use of biocontrol agents to manage plant disease in theory and practice. *Phytopathology* 101(9): 1024-1031
- Yigit, F and M. Dikilitas. 2007. Control of *Fusarium* wilt of tomato by combination of *Pseudomonas fluorescens*, non-pathogen *Fusarium* and *Trichoderma harzianum* T-22 in greenhouse conditions. *Plant Pathol. J.* 6(2):159-163

# BIOCONTROL FOR RHIZOCTONIA STEM ROT DISEASE BY USING COMBINATION OF SPECIFIC ENDOPHYTE IN PADDY TIDAL SWAMP

## ORIGINALITY REPORT

15%

SIMILARITY INDEX

12%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.academicjournals.org">www.academicjournals.org</a> Internet Source	4%
2	<a href="http://jhpttropika.fp.unila.ac.id">jhpttropika.fp.unila.ac.id</a> Internet Source	3%
3	<a href="http://sanningengevoelenspotrebujes.com">sanningengevoelenspotrebujes.com</a> Internet Source	2%
4	<a href="http://ajouronline.com">ajouronline.com</a> Internet Source	1%
5	Baharuddin, R Jahuddin, A Yani, M Tuwo. " In vitro testing for effectiveness of and MO Plus against (Blast Disease) and pv. oryzae (Bacterial Leaf Blight) ", IOP Conference Series: Earth and Environmental Science, 2019 Publication	1%
6	J. A. Lewis, R. P. Larkin, D. L. Rogers. " A Formulation of and to Reduce Damping-off Caused by and Saprophytic Growth of the	1%

# Pathogen in Soilless Mix ", Plant Disease, 1998

Publication

7

Soil Biology, 2015.

Publication

1 %

8

"How Research Can Stimulate the Development of Commercial Biological Control Against Plant Diseases", Springer Science and Business Media LLC, 2020

Publication

<1 %

9

X.-M. Xu, M. J. Jeger. "Theoretical Modeling Suggests that Synergy May Result from Combined Use of Two Biocontrol Agents for Controlling Foliar Pathogens Under Spatial Heterogeneous Conditions", Phytopathology, 2013

Publication

<1 %

10

[www.researchgate.net](http://www.researchgate.net)

Internet Source

<1 %

11

[dergipark.org.tr](http://dergipark.org.tr)

Internet Source

<1 %

12

[pjbt.org](http://pjbt.org)

Internet Source

<1 %

13

Biological Management of Diseases of Crops, 2013.

Publication

<1 %

14 Syed Sartaj Alam, Kazunori Sakamoto, Kazuyuki Inubushi. " Biocontrol efficiency of Fusarium wilt diseases by a root-colonizing fungus sp. ", Soil Science and Plant Nutrition, 2011  
Publication <1 %

---

15 scialert.net  
Internet Source <1 %

---

16 J.A Lewis, R.D Lumsden. "Biocontrol of damping-off of greenhouse-grown crops caused by Rhizoctonia solani with a formulation of Trichoderma spp.", Crop Protection, 2001  
Publication <1 %

---

17 Mohsin Tariq, Sumera Yasmin, Fauzia Y. Hafeez. "Biological control of potato black scurf by rhizosphere associated bacteria", Brazilian Journal of Microbiology, 2010  
Publication <1 %

---

18 ir.lib.uwo.ca  
Internet Source <1 %

---

19 mafiadoc.com  
Internet Source <1 %

---

20 www.iobc-wprs.org  
Internet Source <1 %

---

21

Fahri YIGIT ., Murat DIKILITAS .. "Control of Fusarium Wilt of Tomato by Combination of Fluorescent Pseudomonas, Non-pathogen Fusarium and Trichoderma harzianum T-22 in Greenhouse Conditions", Plant Pathology Journal, 2007

Publication

<1 %

22

[hdl.handle.net](http://hdl.handle.net)

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On