

# Prosiding Konferensi Internasional-2020

*by - -*

---

**Submission date:** 18-Jun-2024 08:27PM (UTC+0700)

**Submission ID:** 2404783379

**File name:** Prosiding\_Konferensi\_Internasional-2020.pdf (652.18K)

**Word count:** 2773

**Character count:** 14868

# Changes in Soil Chemical Properties and Growth Performance of Corn (*Zea mays* L.) Grown in an Acidic-tropical Soil Amended with Fly-ash

Akhmad Rizalli Saidy<sup>1,2</sup>, Bambang Joko Priatmadi<sup>1,2</sup> and Rizmi Yunita<sup>2</sup>

<sup>1</sup>Department of Soil, Faculty of Agriculture, Lambung Mangkurat University, Jalan Achmad Yani KM 36 Banjarbaru 70714, Indonesia.

<sup>2</sup>Doctoral Program of Agricultural Science, Postgraduate Program of Lambung Mangkurat University, Jalan Achmad Yani KM 36 Banjarbaru 70714, Indonesia

Keywords: Toxic elements, soil amendment, lime materials, heavy metals, exchangeable aluminium

**Abstract:** Fly-ash, by product in coal power-plant production, has many essential nutrients for plant growth along with toxic metals. Therefore, fly-ash is frequently applied for improving soil fertility. In this experiment, we applied four different amounts of fly-ash, viz. 0 (100% soil), 25, 50 and 75 Mg ha<sup>-1</sup> to an acidic-tropical soil and determined the changes in soil chemical properties and corn production in a green-house experiment. Results of study showed that fly-ash application improved soil pH, exchangeable cations (Na, K, Ca and Mg), cation exchangeable capacity (CEC), concentrations of aluminium and iron. The experiment also showed that fly-ash application to soils also improved corn growth (plant height and dried-weight shoot) and production. Fly-ash 50 Mg ha<sup>-1</sup> lead to higher corn production than the fly-ash application of 25 Mg ha<sup>-1</sup>, and increasing the amounts of added fly-ash did not significantly increase the corn production. Results of this study demonstrate that medium-level of fly-ash application resulted in the improvements of soil chemical properties and corn production.

## 1 INTRODUCTION

One of the problems corn cultivation in dry-land is high soil-acidity (low pH of the soil). Low soil pH results in low availability of soil phosphorus, calcium, magnesium, potassium and sodium (Bohn *et al.*, 2001), and this condition eventually results in unhealthy plant growth. The soil acidity problem can be reduced by the application of soil ameliorant. Soil amelioration materials such as lime and organic matter have been generally applied to soils for soil characteristic improvement (Ma *et al.*, 2018; Soong *et al.*, 2018). Soil amelioration increases soil pH (dos Santos *et al.*, 2018; Jayalath *et al.*, 2016; Li & Johnson, 2016; Soong *et al.*, 2018) and improves the content of nutrients such as N, P, K, Ca and Mg (Holland *et al.*, 2018; Rheinheimer *et al.*, 2018; Sod *et al.*, 2018).

Fly-ash is the result of the combustion process that contains elements such as Si, K, Ca, Mg and Na (Swain *et al.*, 2007), which is generally existed in oxide form so that is able to increase the soil pH and

cation exchange capacity (Clark *et al.*, 2001). Fly-ash generally contains 4.9 to 25.2% CaO, 1.3 to 5.1% MgO, 1.0 to 1.7% Na<sub>2</sub>O, 0.6 to 1.3% K<sub>2</sub>O and from 36.9 to 52.5 % SiO<sub>2</sub> (Kishor *et al.*, 2010); therefore, fly-ash is potential to be applied for soil amelioration. However, the use of fly-ash as a soil ameliorant is currently not done broadly because of the high concentration heavy metals in the fly-ash.

The use of fly-ash for improvement of soil properties is already done in some countries. Previous research showed that application of fly-ash to soils as a source of Si is able to increase the uptake of Si, P and K, and thereby increasing rice production (Lee *et al.*, 2006). The research also showed that no accumulation of heavy metals in rice plant was observed after fly-ash application. In this study, we applied different amounts of fly-ash to soils and then determined changes in chemical soil characteristics and the growth performance of corn grown at that amended soils.

## 2 MATERIALS AND METHODS

### 2.1 Sampling of Soil and Fly-ash

Soil samples for this experiment was collected from the Experimental Plot for Up-land Crops, Lambung Mangkurat University, Pelaihari, South Kalimantan, Indonesia. The soil was sampled at a depth of 0-30 cm at several different points, and then the samples were homogenized. Plant debris was carefully removed from the collected samples and then stored in polyethylene vessel at 5 °C. Fly-ash used for this study was sampled from the Asam-asam Steam Power Plant, South Kalimantan Province, Indonesia.

### 2.2 Greenhouse Experiment

Seven kilograms of soil was placed in a 10-L pot, then fly-ash in equal amount to 0, 25, 50 and 75 Mg ha<sup>-1</sup> was added to the pots, swirled gently until the fly-ash evenly distributed to the soils. Water was applied to the pots to acquire 70% of water field capacity, then the pots were incubated for 15 days. Each treatment of soil and fly-ash had five replicates. Water was added periodically to each pot during the incubation period to compensate for evaporative losses and ensure constant water content in each pot. At the end of incubation, a 250 g of soil sub-sample was taken from the pot for determination of soil pH, exchangeable cations (Na, K, Ca and Mg), cation exchangeable capacity, exchangeable aluminium and iron, mineral nitrogen and available phosphorus.

Seeds of corn (2-3 seeds) grown in the pots and only one corn plant was allowed to be grown in the pots after three weeks. Nitrogen, phosphorous and potassium fertilizers were applied to each pot at the rates of 150 kg, 100 kg and 100 kg per hectare, respectively. Plant maintenance was carried out by daily watering, weeding as well as pest and disease control by using pesticides if necessary. Measuring of corn growth and production were conducted after 90 days of planting.

### 2.3 Statistical Analysis

Influence of fly-ash application on changes in the soil properties and the growth of corn was quantified through analysis of variance (Anova) followed by the Least Significant Difference (LSD) test at level of  $P < 0.05$ . Prior to the analysis of variance, Shapiro-Wilk's and Bartlett's tests were carried out to ensure data to be analyzed have a normal distribution and equal variances, respectively. All

statistical tests were conducted using the package of GenStat 12<sup>th</sup> Edition (Payne, 2008).

## 3 RESULTS AND DISCUSSION

### 3.1 Changes in Soil Characteristics with Fly-ash Application

Results of study showed that fly-ash application increased soil CEC, soil pH, exchangeable Al, Fe, inorganic nitrogen and available P (Figure 1). Increased soil pH occurs because fly-ash containing CaO and MgO, which reacted with H<sup>+</sup> ions to neutralize soil acidity. The greater the amount of fly-ash application, the greater the amount of CaO and MgO are given into the soils, thus the greater the change in soil pH. It was reported in the previous study that the neutralization capacity of fly-ash ranged from 0.01 to 3.74 meq per gram H<sub>3</sub>O<sup>+</sup> (Kishor et al., 2010). Increasing the pH of the soil treated with fly-ash was also reported in several other studies (Clark et al., 2001; Lee et al., 2006; Sajwan et al., 2007; Swain et al., 2007).




Figure 1. Changes in soil pH (a), cation exchangeable capacity – CEC (b), mineral/inorganic nitrogen (c), exchangeable aluminium (d), exchangeable iron (e), and available P (f) as influenced by the different amount of fly-ash application.




Figure 2: Changes in exchangeable calcium (a), magnesium (b), sodium (c) and potassium as influenced by fly-ash application.

Concentrations of exchangeable aluminium and iron in soil also increased with fly-ash application. The activity and solubility of certain metals may change with changes in pH. For example, aluminium is relatively insoluble as  $Al(OH)_3$  at neutral pH, but it exists predominantly as highly soluble at pH range of 3.4 and 6.3 (Bohn et al., 2001; Sposito, 2004).

Iron is also observed in insoluble phase at neutral pH, and the solubility of iron increase considerably when pH drops to a range of 3.7 and 5.9 (Sposito, 2004). When fly-ash was added to soils with a range of acidic pH, the aluminium and iron contained in the fly-ash dissolved and increased the amount of exchangeable aluminium and iron in soils.

Available phosphor and inorganic nitrogen ( $NH_4^+$  and  $NO_3^-$ ) of the soils increased significantly with fly-ash application (Figure 1). Low concentration of phosphor is often reported for acidic soils due to strong retention of phosphorus ion by mineral soils (Vitousek et al., 2010). Soil pH is believed to be one of the most significant soil characteristics since it affects nutrient availability in soil system (Kemmitt et al., 2006; Lauber et al., 2009). Fly-ash application increased soil pH and eventually improved available phosphor (Figure 1). Another possible explanation for increasing available phosphor with fly-ash application is that fly-ash contains essential nutrients such K, Mg, P and S (Knapp & Insam, 2011). Consequently, fly-ash application to soils may provide soils with an abundant range of essential nutrients, among them phosphor. Previous study showed that significant increase in available phosphor for plants in soil systems following the incorporation of fly-ash to the soils (Schönegger et al., 2018).

Plant-available nitrogen ( $\text{NH}_4^+$  and  $\text{NO}_3^-$ ) also improved significantly after application of fly-ash to the soils. It is well documented that soil pH plays an important role in controlling the rate of nitrogen mineralization, which increasing soil pH enhance organic nitrogen mineralization (Kemmitt *et al.*, 2006; Ste-Marie & Paré, 1999; Thompson *et al.*, 1954). Fly-ash incorporation in this study was able to improve soil pH from 4.7 to 5.1-5.7, depending the amounts of added fly-ash (Figure 1). Increasing soil pH subsequently leads to increase in the concentration of plant-available nitrogen, consistent with previous study reported that application of 50  $\text{Mg ha}^{-1}$  fly-ash enhance plant-available inorganic nitrogen in paddy field (Singh *et al.*, 2011).

Results of the study showed that the fly-ash application affected the concentration of exchangeable Na, Ca, and Mg in soil. Fig. 2 showed that the concentration of exchangeable Ca and Mg was higher than other cations (Na, K, and Mg) in all soils applied with fly-ash. This is due to the high Ca and Mg contents than other cations (Na and K) of

fly-ash, so that the availability of Ca and Mg increased significantly when the fly-ash is added to the soils. This result is consistent with the previous study that reported that Ca is the most dominant cation of soils applied with fly-ash (Sharma & Kalra, 2006).

### 3.2 Growth Performance of Corn with Fly-ash Application

Growth performance of corn (plant height, dry shoot matter weight and yield) was also improved by fly-ash application (Figure 3). Increasing growth performance of corn is attributed to increase in nutrient uptake by corn with fly-ash application. This was supported by significant correlations between nitrogen, phosphor and potassium uptake and the parameters of corn growth performance (plant height, dry shoot matter weight and yield) (data not shown).




Figure 3: Influence of fly-ash application on corn growth and production

## 4 CONCLUSIONS

Observations on the chemical characteristics of soil showed that soil pH, exchangeable bases (Ca, Mg, Na and K), cation exchangeable capacity (CEC), mineral nitrogen, available P, exchangeable aluminium and iron improved with fly-ash application. The growth and production of corn also improve with fly-ash addition to soils. Increases in maize growth and production may relate to increasing the nutrient uptake by corn with fly-ash application. It could be concluded that fly-ash application to soils is able to improve the fertility of acidic soils and the growth of plant.

## ACKNOWLEDGEMENTS

The authors acknowledged the PT. Perusahaan Listrik Negara (PLN) (Persero) for funding this study through the cooperation between PT PLN (Persero) and Environmental Research Centre, Lambung Mangkurat University.

## REFERENCES

Bohn, H.L., McNeal, B.L., O'Connor, G., 2001. *Soil Chemistry*. John Wiley & Sons, New York.

- Clark, R., Zeto, S., Ritchey, K., Baligar, V., 2001. Mineral acquisition by maize grown in acidic soil amended with coal combustion products. *Communications in Soil Science and Plant Analysis* 32(11-12), 1861-1884.
- dos Santos, D.R., Tiecher, T., Gonzatto, R., Santanna, M.A., Brunetto, G., da Silva, L.S., 2018. Long-term effect of surface and incorporated liming in the conversion of natural grassland to no-till system for grain production in a highly acidic sandy-loam Ultisol from South Brazilian Campos. *Soil and Tillage Research* 180, 222-231.
- Holland, J.E., Bennett, A.E., Newton, A.C., White, P.J., McKenzie, B.M., George, T.S., Pakeman, R.J., Bailey, J.S., Fornara, D.A., Hayes, R.C., 2018. Liming impacts on soils, crops and biodiversity in the UK: A review. *Science of The Total Environment* 610-611, 316-332.
- Jayalath, N., Mosley, L., Fitzpatrick, R., Marschner, P., 2016. Addition of organic matter influences pH changes in reduced and oxidised acid sulfate soils. *Geoderma* 262, 125-132.
- Kemmitt, S.J., Wright, D., Goulding, K.W., Jones, D.L., 2006. pH regulation of carbon and nitrogen dynamics in two agricultural soils. *Soil Biology and Biochemistry* 38(5), 898-911.
- Kishor, P., Ghosh, A., Kumar, D. 2010. Use of fly ash in agriculture: A way to improve soil fertility and its productivity. *Asian Journal of Agricultural Research* 4(1), 1-14.
- Knapp, B.A., Insam, H. 2011. Recycling of biomass ashes: current technologies and future research needs. in: *Recycling of Biomass Ashes*, Springer, pp. 1-16.
- Lauber, C.L., Hamady, M., Knight, R., Fierer, N., 2009. Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Applied and Environmental Microbiology* 75(15), 5111-5120.
- Lee, H., Ha, H.S., Lee, C.H., Lee, Y.B., Kim, P.J., 2006. Fly ash effect on improving soil properties and rice productivity in Korean paddy soils. *Bioresource Technology* 97(13), 1490-1497.
- Li, W., Johnson, C.E., 2016. Relationships among pH, aluminum solubility and aluminum complexation with organic matter in acid forest soils of the Northeastern United States. *Geoderma* 271, 234-242.
- Ma, B., Lv, X., Cai, Y., Chang, S.X., Dyck, M.F., 2018. Liming does not counteract the influence of long-term fertilization on soil bacterial community structure and its co-occurrence pattern. *Soil Biology and Biochemistry* 123, 45-53.
- Payne, R., 2008. *A Guide to Anova and Design in Genstat*. VSN International, Hempstead, UK.
- Rheinheimer, D.S., Tiecher, T., Gonzatto, R., Zafar, M., Brunetto, G., 2018. Residual effect of surface-applied lime on soil acidity properties in a long-term experiment under no-till in a Southern Brazilian sandy Ultisol. *Geoderma* 313, 7-16.
- Sajwan, K., Paramasivam, S., Alva, A., 2007. Effects of different rates of fly ash and sewage sludge mixture amendments on cation availability and their leachability. *Journal of Environmental Science and Health Part A* 42(8), 1155-1160.
- Schönegger, D., Gómez-Brandón, M., Mazzier, T., Insam, H., Hermanns, R., Leijenhorst, E., Bardelli, T., Fernández-Delgado Juárez, M. 2018. Phosphorus fertilising potential of fly ash and effects on soil microbiota and crop. *Resources, Conservation and Recycling* 134, 262-270.
- Sharma, S.K., Kalra, N. 2006. Effect of flyash incorporation on soil properties and productivity of crops: A review. *Resources, Conservation and Recycling* 87, 45-65.
- Singh, J.S., Pandey, V.C., Singh, D., 2011. Coal fly ash and farmyard manure amendments in dry-land paddy agriculture field: effect on N-dynamics and paddy productivity. *Applied Soil Ecology* 47(2), 133-140.
- Soong, J.L., Marañon-Jimenez, S., Cotrufo, M.F., Boeckx, P., Bodé, S., Guenet, B., Peñuelas, J., Richter, A., Stahl, C., Verbruggen, E., 2018. Soil microbial CNP and respiration responses to organic matter and nutrient additions: Evidence from a tropical soil incubation. *Soil Biology and Biochemistry* 122, 141-149.
- Sposito, G., 2004. *The Surface Chemistry of Natural Particles*. Oxford University Press on Demand.
- Ste-Marie, C., Paré, D., 1999. Soil, pH and N availability effects on net nitrification in the forest floors of a range of boreal forest stands. *Soil Biology and Biochemistry* 31(11), 1579-1589.
- Swain, D., Rautaray, S., Ghosh, B., 2007. Alkaline coal fly ash amendments are recommended for improving rice-peanut crops. *Acta Agriculturae Scandinavica Section B-Soil and Plant Science* 57(3), 201-211.
- Thompson, L., Black, C., Zoellner, J., 1954. Occurrence and mineralization of organic phosphorus in soils, with particular reference to associations with nitrogen, carbon, and pH. *Soil Science* 77(3), 185-196.
- Vitousek, P.M., Porder, S., Houlton, B.Z., Chadwick, O.A., 2010. Terrestrial phosphorus limitation: mechanisms, implications, and nitrogen-phosphorus interactions. *Ecological Applications* 20(1), 5-15.

# Prosiding Konferensi Internasional-2020

---

## ORIGINALITY REPORT

---

16%

SIMILARITY INDEX

15%

INTERNET SOURCES

14%

PUBLICATIONS

3%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

1	Biogeochemistry of Trace Elements in Coal and Coal Combustion Byproducts, 1999. Publication	2%
2	Vimal Chandra Pandey, Nandita Singh. "Impact of fly ash incorporation in soil systems", Agriculture, Ecosystems & Environment, 2010 Publication	2%
3	<a href="http://www.nal.usda.gov">www.nal.usda.gov</a> Internet Source	2%
4	Submitted to Universitas Sebelas Maret Student Paper	1%
5	<a href="http://link.springer.com">link.springer.com</a> Internet Source	1%
6	<a href="http://journal.unila.ac.id">journal.unila.ac.id</a> Internet Source	1%
7	Muthia Elma, Ni Kadek Devi Ananda Saraswati, Paskah Fransiska Afrida Simatupang, Retno Febriyanti et al. "Hydrogel derived from water hyacinth and pectin from	1%

# banana peel as a membrane layer", Materials Today: Proceedings, 2023

Publication

8

Nadeesha L. Ukwattage, U.Vajini Lakmalie, Ranjith P. Gamage. "Soil and plant growth response and trace elements accumulation in sweet corn and snow pea grown under fresh and carbonated coal fly ash amendment", Agronomy Journal, 2021

Publication

1 %

9

Deborah Schönegger, María Gómez-Brandón, Thomas Mazzier, Heribert Insam et al. "Phosphorus fertilising potential of fly ash and effects on soil microbiota and crop", Resources, Conservation and Recycling, 2018

Publication

1 %

10

[doczz.net](https://doczz.net)

Internet Source

1 %

11

[journals.mums.ac.ir](https://journals.mums.ac.ir)

Internet Source

1 %

12

Adil Ameen Bhat, Adnan Shakeel, Hosny H. Kesba, Zafar Ahmad Handoo, Abrar Ahmed Khan. "Elucidating the role of fly ash in root-knot nematode (*Meloidogyne incognita*) suppression on okra (*Abelmoschus esculentus* L.): Insights into cellular viability and host defence system", Physiological and Molecular Plant Pathology, 2023

1 %



**13** Aqli Mursadin, Andi Yuwenda Iriyanto. "Modeling Steam Power Plant Condenser Vacuum under Off-design Conditions: A Statistical Approach", 2023 International Conference on Advanced Mechatronics, Intelligent Manufacture and Industrial Automation (ICAMIMIA), 2023 **1** %

Publication

---

**14** Vimal Chandra Pandey. "Opportunities and challenges in fly ash-aided paddy agriculture", Elsevier BV, 2020 **1** %

Publication

---

**15** [helda.helsinki.fi](https://helda.helsinki.fi) **1** %

Internet Source

---

**16** [indianecologicalsociety.com](https://indianecologicalsociety.com) **1** %

Internet Source

---

**17** [www.ncbi.nlm.nih.gov](https://www.ncbi.nlm.nih.gov) **1** %

Internet Source

---

Exclude quotes  On

Exclude matches  < 1%

Exclude bibliography  On