

ASSESSING THE HEALTH OF SOUTH KALIMANTAN COASTAL SWAMP WETLANDS USING MEASUREMENTS OF HEAVY METALS IN COMMERCIAL FISH SPECIES

by H.B. Santoso

Submission date: 27-Jun-2024 07:21PM (UTC+0700)

Submission ID: 2409391694

File name: IJCS-24-02_47_Santoso.pdf (1.02M)

Word count: 12814

Character count: 65388

ASSESSING THE HEALTH OF SOUTH KALIMANTAN COASTAL SWAMP WETLANDS USING MEASUREMENTS OF HEAVY METALS IN COMMERCIAL FISH SPECIES

Heri Budi SANTOSO^{1,*}, Rizmi YUNITA², KRISDIANTO¹

63

24

¹ Department of Biology, Faculty of Mathematics and Natural Sciences, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia, 70714

² Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Lambung Mangkurat University, Banjarbaru, South Kalimantan, Indonesia, 70714

Abstract

Heavy metals pollution damage coastal swamp ecosystems. This study's primary purpose was to determine the amounts of heavy metals in coastal swamp waters and the tissues of three fish species (*Mugil cephalus*, *Arius sagor* and *Plotosus lineatus*). This study determined that the water in the coastal swamps of South Kalimantan was contaminated with heavy metals in the following order: Fe > Cu > Zn > Cr > Pb > Hg. The heavy metal concentrations found in the three commercial fish species are as follows: Fe > Zn > Cu > Cr > Pb > Cd > Hg. The bioaccumulation of heavy metals was highest in *Mugil cephalus*, followed by *Arius sagor* and *Plotosus lineatus*. The Fe, Cu and Cr concentrations in *Mugil cephalus*, *Arius sagor* and *Plotosus lineatus* exceeded the international national MPLs. The liver tissue has the highest capacity to bioaccumulate heavy metals. Regular monitoring of these metals in fish and water sources is recommended to ensure the safety and conservation of coastal swamp wetlands.

Keywords: Fish metal accumulation; Bioindicators; Biodiversity threats; Coastal swamp conservation

Introduction

The coastal swamp wetland is a tidal swamp part of the estuary ecosystem. Wetland habitats are well recognized as highly productive ecosystems, providing favorable conditions for various fish species to flourish. Numerous fish species find refuge, nourishment and reproduction opportunities in coastal swamp wetland environments [1]. Nevertheless, wetland ecosystems also exhibit high sensitivity and ecological vulnerability [2]. Several wetland ecosystems, including coastal swamps, face severe threats from heavy metal contamination. It is essential to promptly address this issue [3]. Due to the ease with which these contaminants enter the food chain, coastal swamp ecosystems are susceptible to the negative impacts of heavy metal contamination. Heavy metals harm coastal swamp ecosystems due to their inherent toxicity, permanence, nonbiodegradability and tendency to bioaccumulate in organisms, a phenomenon known as biomagnification [4]. Heavy metals in coastal wetland habitats are caused by two primary sources of contamination: naturally occurring deposits and human activities, such as waste produced by households, industries and agricultural practices [5, 6]. The coastal swamp ecosystems of Kuala Lupak and Kuala Tambangan are prone to heavy metal contamination due to unregulated waste disposal. Unfortunately, monitoring water quality in

* Corresponding author: heribudisantoso@ulm.ac.id

these places has been neglected, necessitating the deployment of assessments to collect vital information regarding the health of these coastal swamp ecosystems. The presence of heavy metal contamination in wetland ecosystems not only accelerates the decline in water quality but also disrupts delicate biological equilibrium and affects biodiversity. Organisms inhabiting wetland ecosystems, mainly fish, can accumulate heavy metals over long periods, exposing them to the potential for both lethal and sublethal effects linked with the heavy metals [2]. The absence of routine monitoring of heavy metal contamination results in the increasing accumulation of heavy metals in fish, making their recovery a challenging problem. The presence of high levels of heavy metal contamination in fish has substantial long-term effects, including the development of diseases. This poses a significant threat to organisms' health, biological systems' viability and coastal ecosystems' core functioning [7].

Due to their outstanding capability to accumulate heavy metals, fish have been utilised to evaluate the ecological health of aquatic systems. This characteristic is essential because it is directly related to their trophic location, frequently at the top of the food chain. The presence of heavy metals in fish has histological and lethal significance [2]. Compared to other aquatic animals, fish are more sensitive to the toxicity of heavy metals, making them an ideal indicator of the ecological health of wetland ecosystems [8]. In contrast to organic pollutants, heavy metals are more resistant to decomposition by natural processes. The deposition of heavy metals in sediments is followed by their mobilization up the food chain, which results in biomagnification at higher trophic levels. This problem threatens the health and biodiversity of fish and human populations [3, 9]. Using fish as a biomonitoring instrument for assessing heavy metal concentrations and their subsequent bioaccumulation in aquatic environments is an appropriate and relevant method for determining the bioavailability of heavy metals [10]. Fish are valuable bioindicators in ecological research. They can reveal chronic and acute exposure patterns, provide the prognosis of pollution threats within ecosystems and thoroughly understand trophic and ecosystem health [11]. This assessment is vital to the mitigation procedure for effectively managing coastal swamp ecosystems. The study by Santoso *et al.* [12] concentrated on the mitigation measures utilized to mitigate the negative impacts of heavy metal pollution on giant mudskipper fish in the estuary of the Barito River. However, there has been no exploration of the presence of heavy metal contaminants in commercially consumed fish species, such as mullets (*Mugil cephalus*), eel-tailed catfish (*Plotosus lineatus*) and sea catfish (*Arisaeta sagor*), that coastal residents of South Kalimantan frequently consume. The evaluation of heavy metal deposition in commercial fish is essential for assessing the health of coastal wetland ecosystems and monitoring food safety in the community.

Mullets, or belanak (the local name), are a species of fish that inhabit maritime environments and estuaries and have substantial economic importance. Mullets have a remarkable tolerance for salinity and temperature fluctuations, allowing them to adapt to various dietary sources within their biological niche. Therefore, they commonly inhabit tropical and subtropical coastal and estuarine habitats. Most of a mullet's diet comprises zooplankton, decaying plants, algae and detritus. When the digestive tract is dissected, sometimes it is found that the stomach contents consist of bits of sand and dirt [13]. This species has a gray coloration and the typical length is between 10-47cm, accompanied by an average body mass ranging from 5-176g [14]. Eel-tailed catfish, known as Sembilang locally, is an economically valuable fish resource in estuarine ecosystems. The general public usually consumes this specific fish as a smoked fish, which is immensely popular due to its excellent flavor and nutritious benefits [15]. The average length of the carnivorous eel-tailed catfish is 25-60cm. Its distribution range encompasses maritime regions, river estuaries and freshwater habitats. This aquatic species demonstrates predatory behavior, preying primarily on juvenile fish and benthic creatures, such as gastropods, mollusks and crustaceans. The position of this fish's second dorsal fin along the

vertical axis between its anal and pelvic fins is one of its distinguishing characteristics. In addition, its upper lip can grow upward or forwards. The barbels can reach the back of the eye and are predominantly dark brown [16]. In estuarine and marine habitats of tropical and subtropical regions, sea catfish, sometimes known as otek in the local dialect, are frequently observed. This fish species demonstrates a dual habitat strategy, first inhabiting freshwater habitats, then migrating to estuary waters for breeding and finally relocating to the open sea. This particular kind of fish belongs to the group of large demersal fish. This type of fish is typically captured between 250-700mm in length, with the potential to reach a maximum size of 1,500mm. Sea catfish have a predominantly carnivorous and predatory diet, devouring mollusks, gastropods and crustaceans. In Indonesia, sea catfish can be spotted in various coastal settings, particularly where river estuaries meet the ocean. This species is typically found between 20 and 100 meters below the surface. It is distinguished by the presence of adipose fins on its dorsal fin and a stiff plate, such as a butterfly or sagor on its head structure [17].

In South Kalimantan, the coastal swamp wetland ecosystems of Kuala Lupak and Kuala Tambangan are characterized by considerable anthropogenic activities. High levels of maritime transportation activities are present in both ecosystems, including the shipping and unloading of coal and the collection of waste from coal mining operations. The discharge of coal mining waste and the occurrence of coal avalanches during loading and unloading/transportation procedures can lead to the introduction of pollutants into water bodies. The accumulation of these contaminants over a lengthy period has the potential to contaminate estuaries [56]. These procedures generate waste containing iron (Fe), manganese (Mn), mercury (Hg), cadmium (Cd), chromium (Cr), lead (Pb) and copper (Cu) [18]. In addition, this aquatic ecosystem is exposed to waste discharge from both industrial and residential sectors. The Kuala Lupak coastal wetland is distinguished by converting mangrove land into ponds, which serve as a hub for ship traffic and residential communities. Along the riverbanks are several industries, including wood/plywood processing facilities, rubber production, fisheries and oil palm plantations. Therefore, the discharge of waste from these sectors creates a substantial threat of contaminating estuarine waters [19, 20]. Our hypothesis contends that insufficient treatment [39] and subsequent discharge of this waste into water bodies may substantially threaten aquatic ecosystems and human health. Consequently [40] it is essential to undertake biomonitoring activities to effectively address and mitigate the adverse effects of heavy metal contamination. Accordingly, this study aims to evaluate the levels of heavy metals in aquatic environments and commercially consumed fish species to understand the ecological health of coastal swamp [21] land ecosystems. This study intends to shed light on the bioavailability and biomagnification [37] of heavy metals in commercial fish populations. In addition, the results of this study were compared to maximum permissible limits (MPL). This study has been carried out in the coastal swamp of Kuala Lupak and Kuala Tambangan river estuaries, South Kalimantan Province in July 2023.


Materials and Methods

Materials

The study was conducted in the coastal swamp waters of Kuala Lupak river estuaries in the Barito Kuala district and Kuala Tambangan river estuaries in the Tanah Laut district of South Kalimantan Province. The study locations are separated by approximately 60km along the coastline of South Kalimantan, which converges with the southern coast of the Java Sea. Table 1 provides a complete summary of each location and study station, encompassing relevant details. Additionally, Figure 1 graphically represents the sample locations.

Table 1. Study station description

Location	Station	Description
Kuala Lupak	Coastal swamp waters	Kuala Lupak is a wildlife conservation area distinguished by the presence of a mangrove swamp forest environment. The Kuala Lupak River traverses this region and residential areas can be found near its mouth. The waters of the river Kuala Lupak flow into the Java Sea. Samples were collected from the coastal swamp located at coordinates 3°27'28.223"S 114°22'04.780" E.
	Coastal waters	The coastal waters of Kuala Lupak are located approximately 8.0km west of the Barito River's mouth, the largest river in South Kalimantan. A high level of human activities characterizes this river. The coastal region of Kuala Lupak is physically connected to the Java Sea, with a coastline spanning approximately 30km. Mangrove swamp forests dominate the shoreline of Kuala Lupak. The study collected samples from the coastal waters where the Kuala Lupak River discharges at 3°28'02.383"S 114°21'31.282" E.
Kuala Tambangan	Coastal swamp waters	There are fishing settlements and mangrove woods in the vicinity of the Kuala Tambangan estuary, which empties into the Java Sea. In the coastal swamp area of this estuary, numerous fishing vessels operate and anchor. Samples were collected from coastal swamp waters of the river estuary located at coordinates 3.968195" S 114.632223" E.
	Coastal waters	Kuala Tambangan's coastal waters are near the Java Sea, with a coastline of approximately 20km. A large number of fishing vessels and coal ships use the coastal waterways. The samples were collected from the coastal waters located at coordinates 3.968005" S 114.629447" E, in the vicinity of the Kuala Tambangan River discharges.



16
 Fig. 1. Study area map and sampling locations in the Kuala Lupak estuary and Kuala Tambangan South Kalimantan, Indonesia

77 thods

Water and fish samples were collected in July 2023. The study employed the purposive sampling technique, which involves selecting samples based on predetermined criteria, such as identifying probable contamination sources. Various actions that cause contamination have

been considered to pinpoint these sources precisely. Figure 1 shows the collection of fish and water samples at the research location.

83 Surface water samples were obtained from two stations around the study locations using the Aqua trap water sampler. To facilitate subsequent analysis, a volume of approximately 2mL of concentrated nitric acid (HNO₃) was introduced to each 1.0L after a filtration process. The concentration level of dissolved heavy metals was determined using the liquid-liquid extraction method, as described by Jayaprakash *et al.* [21]. A 100mL unfiltered sample was added to a separating funnel and combined with 2mL of a 2% solution of ammonium pyrrolidine dithiocarbamate (APDC). After vigorous stirring, 10ml of isobutyl methyl ketone (IBMK) was used for the extraction process. The resulting aqueous phase was then treated for further extraction using strong HNO₃ and high-purity water. Atomic absorption spectrometry (AAS) 36 (Permo Scientific ICE 3500 series Germany) was used to examine the final solution after any traces of organic solvents in the solutes were evaporated on a low-temperature hot plate. The AAS was used to quantify 27 lead (Pb), mercury (Hg), cadmium (Cd), zinc (Zn), iron (Fe), copper (Cu) and chromium (Cr) concentrations in both water and fish samples. The concentration of water was quantified in micrograms per liter (µg L⁻¹). Finally, the samples for metal analysis were properly prepared following established procedures Jayaprakash *et al.* [21].

The fish samples included *Plotosus lineatus*, *Arius sagor* and *Mugil cephalus*. Local fishermen obtained these specimens from coastal swamp wetland waters (Fig. 2).



Fig. 2. Fish species used in the Study A: *Mugil cephalus*; B: *Arius sagor*; C: *Plotosus lineatus*

To ensure representativeness, a sample size of 15 fish per species was taken from each station (coastal swamp and coastal) and sampling location (Kuala Lupak and Kuala Tambangan). The identification of the fish samples was also used cross-referencing with the information from Kottelat *et al.* [22]. This was done to ensure that the classification of the species as either commercial or noncommercial and the determination of its name, family, diet, habitat and type were accurate. To preserve 39 and keep the samples in the best possible condition for further analysis, they were carefully transported to the lab in an insulated container filled with ice. In addition, fish dissection entails using ice cubes to protect the animals and surgical tools made of stainless steel to help remove the liver, muscles and skin tissues. A digestion tube preloaded with 5mL of HNO₃ and sulfuric acid (H₂SO₄) was used for each organ sample to activate the reaction. A hot block digestion equipment kept at 60°C for 30 minutes was used to process the samples. After the samples were cooled, 10mL of HNO₃ was added and the mixture was then heated at temperatures between 120 and 150°C until the solution developed a distinctly dark color. One milliliter of hydrogen peroxide (H₂O₂) was added to the mix to produce a clear solution, which aided the filtration procedure 48. These filtered samples were then subjected to an AAS analysis to determine the number of 29 heavy metals in the fish tissues, expressed in mg·kg⁻¹. The limits of detection used for analysis of Pb, Cd, Zn, Fe, Cr, Cu and Hg were 0.001, 0.001, 0.001, 0.006, 0.002, 0.007 and 0.00004mgkg⁻¹, respectively.

Statistical analysis 72

The collected data about heavy metal concentrations in water bodies and fish tissues met the assumptions required to analyze variance (ANOVA), specifically normality and

homogeneity. Therefore, factorial ANOVA can be used to diagnose the data correctly. Then, a post hoc test, precisely Tukey's honestly significant difference (Tukey's HSD) test, with a 95% confidence level was used to identify the difference in treatment ($P < 0.05$). The information is presented in tabular form, showing the means \pm SDs. To determine the relationship between heavy metal concentration variables in water and fish tissues, Pearson correlation analysis was used at a 95% confidence level. The statistical analysis was performed using version 9.4 of the Statistical Analysis System (SAS) software (SAS institute North Carolina State University).

58

Results and discussion

Heavy metal concentrations in the water column

The water samples collected at the research location exhibited a range of heavy metal concentrations, spanning from 0.075 to 620 μ gL⁻¹. At the Kuala Lupak location, the water shows varying concentrations of heavy metals. The Fe concentration is the highest at 620 μ gL⁻¹, while the Hg concentration is the lowest at 0.075 μ gL⁻¹. In the Kuala Tambangan location, the highest concentration is Fe at 370 μ gL⁻¹ while the lowest is Cd at 0.19 μ gL⁻¹, as indicated in Table 2. Heavy metal concentrations in water were significantly different at the locations of Kuala Lupak and Kuala Tambangan ($P < 0.05$). Cd, Cr, Cu, Fe, Pb and Zn concentrations in the water at the Kuala Lupak location are higher than those at the Kuala Tambangan location. Except for Cr, the results of comparing the concentrations of seven different heavy metals in water at two designated stations suggest no statistically significant changes. The concentration of Cr in the coastal swamp is more significant than that of the coast, as seen in Table 2.

59

Table 2. The concentration of heavy metals in water bodies at the study location

Station	Concentration (μ gL ⁻¹)						
	Cd	Cr	Cu	Fe	Hg	Pb	Zn
Coastal swamp	0.20 \pm 0.00 ^a	15 \pm 0.00 ^a	43 \pm 0.024 ^a	495 \pm 0.137 ^a	0.198 \pm 0.135 ^a	0.97 \pm 0.00002 ^a	21 \pm 0.016 ^a
Coastal	0.20 \pm 0.00 ^a	14 \pm 0.001 ^b	43 \pm 0.025 ^a	494 \pm 0.133 ^a	0.186 \pm 0.122 ^a	0.95 \pm 0.00004 ^a	20 \pm 0.015 ^a
Location							
Kuala Lupak	0.21 \pm 0 ^a	15 \pm 0 ^a	65 \pm 0.001 ^a	620 \pm 0.00 ^a	0.075 \pm 0.001 ^b	0.98 \pm 0.00001 ^a	35 \pm 0.001 ^a
Kuala Tambangan	0.19 \pm 0 ^b	14 \pm 0 ^b	21 \pm 0.001 ^b	370 \pm 0.00 ^b	0.309 \pm 0.013 ^a	0.93 \pm 0.00002 ^b	6 \pm 0.00 ^b

Values are the mean and SD. According to the Honestly Significant Difference test with a 95% confidence level, values followed by various letters on the same type of metal reveal a significant difference

This indicates that the coastal swamp and coastal stations do not significantly impact the amounts of heavy metals in the water ($P > 0.05$). In this study, the general order of heavy metal concentrations in water was Fe > Cu > Zn > Cr > Pb > Cd > Hg. Table 3 demonstrates that Fe has the highest concentration in water bodies based on the findings of this study and prior research conducted at estuaries or coastal stations in various parts of the world. Even though the concentrations of just three heavy metals (Fe, Cu and Zn) were found to be excessive in comparison to earlier studies conducted in estuaries in other countries, the total results of this study, except for Fe, remained below the limits set by the World Health Organization (WHO) and the United States Environmental Protection Agency (USEPA). The provisions of Indonesia Government Regulation No. 22 of 2021 on the Implementation of Environmental Protection and Management specifically regarding the seawater quality criteria for marine biota, determined that the concentrations of Fe, Cu and Cr are above the MPL listed in Table 3.

It is necessary to assess heavy metal pollution in coastal wetland ecosystems to comprehend the status of contamination, emission sources and environmental behavior, given the importance of wetland ecosystems in maintaining environmental stability and providing essential ecological services, particularly for protecting rich biodiversity and conservation. In addition, knowledge regarding the amounts of heavy metals in water and commercial fish is vital to ensuring the safety of human food consumption [23].

Table 3. Heavy metal concentrations (μgL^{-1}) in coastal swamps of Kuala Lupak and Kuala Tambangan and other parts of the world

Study site	Cd	Cr	Cu	Fe	Hg	Pb	Zn	References
Kuala Lupak estuary, Indonesia	0.21	15	65	620	0.075	0.98	35	Present study
Kuala Tambangan estuary, Indonesia	0.19	14	21	370	0.309	0.93	6	Present study
Southeast coast of India	1.34	2.16	3.04	67.16	-	3.65	31.72	Adani <i>et al.</i> [7]
Karachi coasts of Pakistan	5.64	6.83	8.12	24.21	-	21.21	-	Mehar <i>et al.</i> [24]
Great Vedarnyam Swamp, India	0.1	2.9	0.1	-	1.5	1.4	0.3	Pandiyan <i>et al.</i> [1]
Yangtze River Estuary, China	0.0413	0.279	1.47	-	0.0363	0.829	8.91	Fan <i>et al.</i> [25]
San Carlo, Southern Atlantic coast of Spain	0.1	0.5	1.6	8	< 0.1	0.5	12	Usero <i>et al.</i> [26]
USEPA	5	100	1300	300	-	15	5000	USEPA (2018)
WHO	3	50	2000	300	-	10	3000	WHO (2008)
Indonesia Gov. Reg. No. 22 of 2021	1	5	8	300	1	8	50	Indonesia Gov. Reg. No. 22 of 2021

International and National Guidelines values: WHO-World Health Organization; USEPA-United States Environmental Protection Agency; Indonesia Government Regulation No. 22 of 2021 on the Implementation of Environmental Protection and Management specifically regarding the seawater quality criteria for marine biota

According to the findings, it has been observed that the coastal swam⁷⁴ waters of South Kalimantan exhibit the highest concentration of Fe (370 and $620\mu\text{gL}^{-1}$). This observation is consistent with the earlier findings of researchers from three other countries, implying ¹⁵ increased Fe levels are the primary characteristic of estuary water pollution. Nonetheless, the concentration of Fe in this study exceeds the Fe values found on the southeastern coast of India ($67.16\mu\text{gL}^{-1}$) [7], Karachi coast of Pakistan ($24.21\mu\text{gL}^{-1}$) [24] and San Carlos coast of Spain ($8\mu\text{gL}^{-1}$) [26] (Table 3). Increased Fe concentrations in estuarine and coastal waters may result from natural events such as terrestrial geological processes (weathering and decomposition of rocks and ore materials) and volcanic eruptions⁷³ which are released into water bodies through runoff, erosion and flooding. In addition, Fe is produced by human activities such as coal transportation loading and unloading and mining operations [27]. Other anthropogenic activities on land, such as domestic waste, industrial waste and corrosion of water pipelines delivered by rivers to estuaries, also increase Fe concentrations. Iron (Fe^{2+}) is released by the oxidation of iron pyrite (FeS_2) in coal seams to ⁶ sulfuric acid and sulfur [28]. Toxic heavy metals also come from agricultural waste disposal, agricultural runoff, ⁴ fine surface drainage, irrigation return flow, urban rainwater discharge and leachate water [2], the removal of mangrove vegetation and coastal development (ports, industrial districts and civil engineering works) [3]. Near the study location, shipping transportation activities, loading and unloading of coal on barges, coal transportation, industrial areas and activities on riverbank land, residential areas, agricultural activities¹⁰ and the conversion of mangrove land into fishing/farming land were observed.

Heavy metal concentrations in various tissues of different fish species

Table 4 shows that the highest Cd concentration in the liver of *Arius sagor* was found in the Kuala Tambangan estuary (0.072mgkg^{-1}), which was substantially different from the coast ($P < 0.05$). The lowest Cd concentration (0.001mgkg^{-1}) was found in the muscle and skin tissue of the examined fish species. The liver of *Mugil cephalus* in the Kuala Tambangan estuary had the highest Cr concentration (6.70mgkg^{-1}). The skin of *Plotosus lineatus* in the estuary (0.49mgkg^{-1}) and the coast of Kuala Tambangan (0.46mgkg^{-1}) exhibited the lowest Cr concentration. The liver of *Plotosus lineatus* from the Kuala Lupak estuary had the most Cu (14.48mgkg^{-1}). In comparison, the skin and muscle of *P. lineatus* from the Kuala Tambangan estuary and coast contained the least (0.01mgkg^{-1}). The liver of *M. cephalus* in the estuary (800.97mgkg^{-1}) and coast (800.91mgkg^{-1}) of Kuala Tambangan contained the highest quantity of Fe ($P > 0.05$). In contrast, the skin of *P. lineatus* in the estuary and coast of Kuala

Tambangan had the lowest Fe concentrations (17.73 and 17.72mgkg⁻¹, respectively). The same Hg concentration (0.00004mgkg⁻¹) was identified in three tissues from three test species at two locations and two stations (Table 5). The highest Pb concentration was found in the liver of *M. cephalus* from the Kuala Lupak coast (0.701mgkg⁻¹). Nevertheless, the skin and muscles of the three test species collected Pb at the same concentration between the estuary and coastal stations, precisely 0.001mgkg⁻¹ (Table 6). Table 6 depicts the estuary (64.41mgkg⁻¹) and coast (64.39mgkg⁻¹) of Kuala Lupak as having the highest Zn concentration in the liver of *A. sagor* ($P > 0.05$). The muscle of *P. lineatus* from the estuary and coast of Kuala Lupak had the lowest Zn concentration (26.72 and 0.70mgkg⁻¹; $P > 0.05$). The findings of this study indicate that liver tissue exhibited the highest bioaccumulation of heavy metals, with skin and muscle follow suit in terms of their respective levels of bioaccumulation. The amounts of Fe, Cu, Cr, Zn and Pb in the liver tissue of *M. cephalus* were significantly higher than those detected in the liver tissue of *A. sagor* and *P. lineatus*. Based on the geographical positioning, it may be inferred that the coastal swamps of Kuala Lupak are more heavily contaminated with heavy metals than those of Kuala Tambangan. Based on a combination of different locations, stations, species and fish tissues, the order of heavy metal concentrations in this finding is Fe > Zn > Cu > Cr > Pb > Cd > Hg. Iron is the predominant heavy metal polluting the coastal swamp waters of South Kalimantan's Kuala Lupak and Kuala Tambangan.

Table 4. Concentrations of Cd, Cr and Cu in various tissues of different fish species

Species	Loc.	Sta.	Concentration (mgkg ⁻¹)									
			Cd			Cr			Cu			
			Muscles	Liver	Skin	Muscles	Liver	Skin	Muscles	Liver	Skin	
<i>Mugil cephalus</i>	KL	E	0.001±0.0e	0.048±0.001c	0.001±0.0e	1.68±0.00	1.63±0.01	1.53±0.00	6.87±0.00	1.92±0.00	1.86±0.00	
		C	0.001±0.0e	0.041±0.001d	0.001±0.0e	1.68±0.01	1.6±0.03	1.51±0.00	6.85±0.01	1.91±0.02	1.83±0.01	
	KT	E	0.001±0.0e	0.001±0.0e	0.001±0.0e	1.48±0.00	6.70±0.00a	0.96±0.01	1.22±0.01	10.98±0.01c	0.45±0.00	
		C	0.001±0.0e	0.001±0.0e	0.001±0.0e	1.47±0.01	6.67±0.01a	0.91±0.01	1.16±0.03	10.87±0.01d	0.44±0.01	
	<i>Arius sagor</i>	KL	E	0.001±0.0e	0.001±0.0e	0.001±0.0e	1.81±0.00	1.25±0.03	0.88±0.01	2.63±0.01	10.11±0.01	1.51±0.00
			C	0.001±0.0e	0.001±0.0e	0.001±0.0e	1.79±0.02	1.23±0.00	0.81±0.01	2.62±0.01	10.09±0.03	1.49±0.00
KT		E	0.001±0.0e	0.072±0a	0.001±0.0e	2.58±0.00b	2.12±0.00d	2.03±0.01	0.77±0.01	4.03±0.00	1.1±0.00	
		C	0.001±0.0e	0.062±0b	0.001±0.0e	2.52±0.01c	2.11±0.01d	2.01±0.01	0.57±0.01	4.01±0.00	1.09±0.01	
KL		E	0.001±0.0e	0.001±0.0e	0.001±0.0e	0.74±0.00	1.54±0.01	1.47±0.00	2.18±0.01	14.48±0.01a	2.93±0.01	
		C	0.001±0.0e	0.001±0.0e	0.001±0.0e	0.72±0.01	1.44±0.03	1.45±0.01	2.15±0.01	14.43±0.02b	2.87±0.01	
KT	E	0.001±0.0e	0.001±0.0e	0.001±0.0e	1.2±0.00	1.63±0.00	0.49±0.01u	0.01±0.00w	9.39±0.00	0.01±0.00w		
	C	0.001±0.0e	0.001±0.0e	0.001±0.0e	1.19±0.01	1.61±0.01	0.46±0.01u	0.01±0.00w	9.36±0.00	0.01±0.00w		

Loc.: location; KL: Kuala Lupak; KT: Kuala Tambangan; Sta.: station; E: estuarine/coastal swamp; C: coastal. Values are the mean and SD. Mean±std.dev followed by different letters for the same type of metal shows a significant difference based on Tukey's HSD post hoc test with a 95% confidence level; FAO (Food and Agriculture Organization of the United Nations, 1983); WHO (World Health Organization) Expert Committee on Food Additives (1989); US FDA (United States Food and Drug Administration); BPOM RI (Food and Drug Supervisory Agency of the Republic of Indonesia, 2018)

Table 5. Concentrations of Fe and Hg in various tissues of different fish species

Species	Loc.	Sta.	Concentration (mgkg ⁻¹)						
			Fe			Hg			
			Muscles	Liver	Skin	Muscles	Liver	Skin	
<i>Mugil cephalus</i>	KL	E	21.83±0.00	515.55±0.01b	31.12±0.00	0.00004±0	0.00004±0	0.00004±0	
		C	21.82±0.01	515.47±0.02b	31.09±0.00	0.00004±0	0.00004±0	0.00004±0	
	KT	E	26.96±0.01	800.97±0.00a	47.31±0.01	0.00004±0	0.00004±0	0.00004±0	
		C	26.92±0.01	800.91±0.01a	47.29±0.01	0.00004±0	0.00004±0	0.00004±0	
	<i>Arius sagor</i>	KL	E	22.24±0.01	300.28±0.06c	22.87±0.00	0.00004±0	0.00004±0	0.00004±0
			C	22.21±0.01	300.34±0.03c	22.83±0.01	0.00004±0	0.00004±0	0.00004±0
KT	E	25.28±0.00	25.61±0.00	26.45±0.01	0.00004±0	0.00004±0	0.00004±0		
	C	25.23±0.01	25.55±0.01	26.43±0.01	0.00004±0	0.00004±0	0.00004±0		
<i>Plotosus lineatus</i>	KL	E	24.91±0.00	266.33±0.01	20.18±0.00u	0.00004±0	0.00004±0	0.00004±0	
		C	24.2±0.07	265.92±0.52	20.15±0.01u	0.00004±0	0.00004±0	0.00004±0	
KT	E	21.48±0.01	147.12±0.00	17.73±0.01v	0.00004±0	0.00004±0	0.00004±0		
	C	21.41±0.01	145.12±0.01	17.72±0.01v	0.00004±0	0.00004±0	0.00004±0		

Loc.: location; KL: Kuala Lupak; KT: Kuala Tambangan; Sta.: station; E: estuarine/coastal swamp; C: coastal. Values are the mean and SD. Mean±std.dev followed by different letters for the same type of metal shows a significant difference based on Tukey's HSD post hoc test with a 95% confidence level; WHO (World Health Organization) Expert Committee on Food Additives (1989); SNI 7387:2009 (Indonesian National Standard: Maximum limits for heavy metal contamination in food); US FDA (United States Food and Drug Administration)

Table 6. Concentrations of Pb and Zn in various tissues of different fish species

Species	Loc.	Sta.	Concentration (mgkg ⁻¹)						
			Pb			Zn			
			Muscles	Liver	Skin	Muscles	Liver	Skin	
<i>Mugil cephalus</i>	KL	E	0.001±0c	0.698±0b	0.001±0c	17.98±0.01	17.95±0.04	26.01±0.01	
		C	0.001±0c	0.701±0a	0.001±0c	17.92±0.01	17.93±0.02	25.89±0.02	
	KT	E	0.001±0c	0.001±0c	0.001±0c	9.61±0.01	20.69±0.01	6.22±0.01	
		C	0.001±0c	0.001±0c	0.001±0c	9.46±0.01	20.51±0.02	6.21±0.01	
	<i>Arius sagor</i>	KL	E	0.001±0c	0.001±0c	0.001±0c	7.91±0.00	64.41±0.01 a	17.41±0.00
			C	0.001±0c	0.001±0c	0.001±0c	7.84±0.03	64.39±0.02 a	17.24±0.01
KT		E	0.001±0c	0.001±0c	0.001±0c	12.85±0.01	14.97±0.00	23.67±0.01	
		C	0.001±0c	0.001±0c	0.001±0c	12.65±0.01	14.93±0.00	23.45±0.02	
<i>Plotosus lineatus</i>		KL	E	0.001±0c	0.001±0c	0.001±0c	0.72±0.00z	34.62±0.01 b	8.41±0.00
			C	0.001±0c	0.001±0c	0.001±0c	0.70±0.02z	34.59±0.03 b	8.35±0.01
KT	E	0.001±0c	0.001±0c	0.001±0c	4.61±0.01	31.08±0.00	4.43±0.00z		
	C	0.001±0c	0.001±0c	0.001±0c	4.59±0.01	32.08±0.01 c	4.42±0.01z		
Permissible limits in fishes			0.5 (FAO) 2 (WHO) 0.3 (SNI 7387:2009)			30 (FAO) 100 (WHO) 100 (BPOM RI)			

Loc.: station; KL: Kuala Lupak; KT: Kuala Tambangan; Sta.: station; E: estuarine/coastal swamp; C: coastal. Values are the mean and SD. Mean±std.dev follow different letters for the type of metal shows a significant difference based on Tukey's HSD post hoc test with a 95% confidence level. FAO (Food and Agriculture Organization of the United Nations, 1983) WHO (World Health Organization) Expert Committee on Food Additives (1989) BPOM RI (Food and Drug Supervisory Agency of the Republic of Indonesia, 2018) SNI 7387:2009 (Indonesian National Standard: Maximum limits for heavy metal contamination in food)

According to the Pearson correlation analysis presented in Table 7, a significant and positive association is observed between heavy metal concentrations in water bodies and the tissues of *M. cephalus*, specifically the liver (r = 0.782), muscles (r = 0.665) and skin (r = 0.683). These findings indicate that the concentration of heavy metals in the water increases, there is a corresponding increase in the concentration of heavy metals in the tissues above. A significant and positive correlation also exists between the concentration of heavy metals in water and the corresponding concentration found in the liver (r = 0.780), muscles (r = 0.732) and skin (r = 0.560) of *Arius sagor*. A significant and positive correlation between the concentrations of heavy metals found in water bodies and the corresponding concentrations was also observed in the tissues of *Plotosus lineatus*, specifically the liver (r = 0.885), muscles (r = 0.864) and skin (r = 0.812). In general, the liver tissue of the three test fish species exhibited the most significant and positive correlation with concentrations of heavy metals in aquatic environments.

Table 7. Correlation of heavy metal concentrations in water and heavy metal concentrations in different tissues in different fish species

	<i>Mugil cephalus</i>	Liver	Muscles	Skin	Water
Liver		1			
Muscles		.842**	1		
Skin		.901**	.972**	1	
Water		.782**	.665**	.683**	1
<i>Arius sagor</i>					
Liver		1			
Muscles		.647**	1		
Skin		.565**	.942**	1	
Water		.780**	.732**	.560**	1
<i>Plotosus lineatus</i>					
Liver		1			
Muscles		.964**	1		
Skin		.948**	.953**	1	
Water		.885**	.864**	.812**	1

The numbers displayed are correlation coefficient values; coefficient followed by * indicate a relationship between two variables at the 95% confidence level. Correlation at level P < 0.05; ** Correlation at level P < 0.001

Estuaries are known as "sinks," dilution sites and storage grounds for heavy metal-containing waste from land and rivers [29]. Due to dredging activities, massive transportation, waves and storms, silt is removed and particles containing heavy metals are mixed back up from sediment polluted with heavy metals, increasing the concentration of dissolved heavy metals in water bodies [29]. Therefore, aquatic biota, including fish, readily absorb and accumulate heavy metals through their food, skin and gills from contaminated water bodies and sediments. Concentrations of heavy metals in fish mainly refer to heavy metal concentrations in water bodies and sediments that fish typically inhabit [30]. This is consistent with the finding that there is a significant and positive correlation between the concentration of heavy metals in the water and the concentration of heavy metals in the tissues of the test fish (Table 7). Yogeshwaran *et al.* [31] and Soltani *et al.* [32] revealed that the bioaccumulation of heavy metals in estuarine aquatic biota is closely correlated with water contamination. Heavy metals are present in marine biota due to their abundance in water bodies, sediments and fish diets. Deposits include diverse forms of heavy metals, including dissolved, bonded to suspended particles and accumulated, which increases their assimilation into biological processes. The accumulation of heavy metals in fish at concentrations surpassing the threshold value (MPL) can harm human health [5, 33]. It has been reported that the accumulation of metals in fish tissues beyond permissible limits has severe implications that negatively affect fish populations and humans [34]. To reduce the potential health risks connected with fish consumption, it is essential to detect and quantify the levels of heavy metals in commonly consumed commercial fish species.

These data suggest that the liver of *M. cephalus* contains the highest concentration of heavy metals, while the skin of *P. lineatus* contains the lowest. Despite living in identical ecological settings, the three test fish species exhibited differences in metal accumulation that might be attributed to differences in eating preferences, trophic levels, living behavior and heavy metal contamination gradients [21]. Due to its diet of zooplankton, dead plants and sedimentary material, *Mugil cephalus* accumulates the heaviest metals. Upon dissection, the digestive tract of *M. cephalus* carries sand or dirt [13]. The study by Santoso *et al.* [12] revealed that Fe (13.704.410mgkg⁻¹), Zn (81.863mgkg⁻¹), Cu (37.635mgkg⁻¹), Pb (23.772mgkg⁻¹) and Cd (1.142mgkg⁻¹) were found in high concentrations in sediments in the coastal swamp waters of South Kalimantan, indicating that fish that consume sediment detritus and live in a polluted environment will accumulate more heavy metals than other fish species. Fish may easily bioaccumulate several types of heavy metals from water, sediments and aquatic foods [24].

The Fe concentrations in the test fish tissues ranged between 17.72 and 800.94mgkg⁻¹, above the MPL set by the WHO (100mgkg⁻¹) and SNI 7387:2009 (1mg/kg). The liver of *Mugil cephalus* contained the highest concentration of Fe (800.97mgkg⁻¹), while the skin of *Plotosus lineatus* contained the lowest concentration (17.72mgkg⁻¹). The Fe concentration in the liver of *M. cephalus* in this study (800.97mg/kg) was more significant than that in previous similar studies conducted by Fazio *et al.* [35] in the Black Sea of Bulgaria (502.300mgkg⁻¹), Karadede *et al.* [36] in Turkey (200.86mgkg⁻¹), Usero *et al.* [26] in the southern Atlantic coast of Spain (199mgkg⁻¹), El-hak *et al.* [37] in Lake Manzala in Egypt (85.08mgkg⁻¹) and Fazio *et al.* [35] in the Ionian Sea of Italy (33.150mgkg⁻¹). These studies indicated that Fe is the primary pollutant and that the liver is the organ that bioaccumulates heavy metals. The liver of *M. cephalus* accumulated the most Fe, Cu, Cr, Zn and Pb in this study, followed by the liver of *A. sagor* and *P. lineatus*. According to Zaoui *et al.* [30], the liver accumulates the highest amount of heavy metals as fish absorb them from their environment. The accumulation of heavy metals in the liver may be related to its metabolic function. High amounts of heavy metals in liver tissues are often associated with a natural binding protein, such as metallothionein, which serve

as a necessary metal store (i.e., Zn and Cu) to meet enzymatic and other metabolic requirements. Fe also tends to accumulate in liver tissues due to the physiological role of the liver in the formation of red blood cells and hemoglobin [6]. In addition, the elevated metal concentrations identified in the liver may result from its metabolic activity, allowing it to accumulate high metal concentrations. High metal concentrations in the liver may indicate that this organ stores sequestered products. When assessing metal contamination in aquatic environments, the liver is often recommended as a target tissue [38]. The liver is the primary organ responsible for metal accumulation among fish organs. The liver is involved in the natural binding of metallothionein (MT) proteins and plays an essential role in trace metal metabolism in fish. Due to its more significant metabolic activity and function of storing and transporting metals throughout the body, the liver accumulates more than the muscle. In addition, the liver is essential for detoxification [39]. Lower metal accumulation in skin and muscle results from a decreased susceptibility for fat to absorb or combine with heavy metals and a reduced metabolic rate [40].

The highest concentration of Fe in the muscles was found in *M. cephalus* (26.96mgkg^{-1}), while *P. lineatus* had the lowest (21.02mgkg^{-1}). This result is higher than previous findings on *M. cephalus* muscles by El-hak *et al.* [37] in Lake Manzala in Egypt (13.29mgkg^{-1}), Fazio *et al.* [35] in the Black Sea of Bulgaria (11.06mgkg^{-1}) and in the Ionian Sea of Italy (5.92mgkg^{-1}), Karadede *et al.* [36] in Ataturk Dam Lake in Turkey (6.88mgkg^{-1}), Usero *et al.* [26] in the San Carlos estuary of Spain (4.11mgkg^{-1}) and Albuquerque *et al.* [3] in the Maranhao estuary of Brazil (0.72mgkg^{-1}). Muscles were examined for metal contamination since it is the final site of metal contamination and the portion of fish most frequently consumed by humans [2]. The concentration of Fe in the skin of *M. cephalus* was the highest (47.31mgkg^{-1}), while that of *P. lineatus* was the lowest (17.73mgkg^{-1}). Lakra *et al.* [34] discovered lower Fe concentrations in catfish skin (36.18mgkg^{-1}) from India's coal mine waste. Possibly due to the continual release of mucus and shedding of dead cells, the buildup of heavy metals in the skin was not as significant as that reported in the liver. The skin possesses mucogenic activity and vigorous mucus exfoliation to remove clinging toxins; as a result, its concentration is lower than that of the liver [41].

In this study, Zn concentrations were detected in all fish tissue samples. The Zn concentration ranges from 0.70 to 64.41mgkg^{-1} , above the FAO MPL of 30mgkg^{-1} but still below the BPOM RI MPL of 700mgkg^{-1} . The highest Zn content was found in the liver of *A. sagor* (64.41mgkg^{-1}), while the lowest concentration was found in the muscle of *P. lineatus* (0.70mgkg^{-1}). This finding is higher than previous findings in *M. cephalus* liver by Usero *et al.* [26] in San Carlos estuary of Spain (38.40mgkg^{-1}), Fazio *et al.* [35] in the Ionian Sea of Italy (30.00mgkg^{-1}) and Black sea of Bulgaria (18.48mgkg^{-1}), Al-Kazaghly *et al.* [42] off the Zliten coast of Libya (25.27mgkg^{-1}), but these findings are relatively similar to previous results by Hauser-Davis *et al.* [43] in the liver of *M. cephalus* in Itaipu beach of Brazil (64.61mgkg^{-1}). In this study, the Zn concentration in muscles was highest in *A. sagor* (17.98mgkg^{-1}), higher than previous findings in *A. sagor* muscles by Pandion *et al.* [44] in Kalpakkam coast of India (0.90mgkg^{-1}), Usero *et al.* [26] in San Carlos estuary of Spain (3.87mgkg^{-1}), Albuquerque *et al.* [3] in Maranhao estuary of Brazil (0.09mgkg^{-1}), Fazio *et al.* [35] in the Black Sea of Bulgaria (4.53mgkg^{-1}), Al-Kazaghly *et al.* [42] off the Zliten coast of Libya (5.90mgkg^{-1}), Karadede *et al.* [36] in Ataturk Dam Lake in Turkey (7.74mgkg^{-1}). However, this finding is lower than previous findings by Fazio *et al.* [35] on *M. cephalus* muscles in the Ionian Sea Italy (19.30mgkg^{-1}) and Darvish *et al.* [45] in the Persian Gulf (18.75mgkg^{-1}). *M. cephalus* has the highest levels of Zn in its skin (26.01mgkg^{-1}). The majority of fish species receive Zn through

their diet. Zn is crucial for metabolic activities and the proper functioning of many enzymes; hence, fish consume vast amounts of Zn, resulting in increased levels of Zn in fish tissues [46]. Zn is a necessary element for metabolic activities and a cofactor for several enzymes involved in these processes. Despite this, Zn poisoning is marked by nausea, diarrhea, fever, lethargy and fatigue. Zinc metalloenzymes and Zn²⁺-protonated enzymes can also cause Zn accumulation by generating five- or six-membered stable ring chelates [47]. The primary sources of zinc in estuarine waterways include wastewater from the textile industry, detergent waste, waste from crude oil production, refineries, brass manufacture, metal plating and immersion of painted idols [46, 48].

Cu concentrations in the fish tissues ranged from 0.01 to 14.48 mg kg⁻¹, above the WHO MPL (3 mg kg⁻¹). The highest Cu content was identified in the liver of *P. lineatus* (14.48 mg kg⁻¹), while the lowest concentration was found in the skin and muscle of *P. lineatus* (0.01 mg kg⁻¹). This finding is lower than previous findings in *M. cephalus* liver by Usero *et al.* [26] in the San Carlos estuary of Spain (17.00 mg kg⁻¹), El-hak *et al.* [37] in Lake Manzala in Egypt (19.79 mg kg⁻¹), Fazio *et al.* [35] in the Black Sea of Bulgaria (68.62 mg kg⁻¹) and Karadede *et al.* [36] in Ataturk Dam Lake in Turkey (267.86 mg kg⁻¹). In this study, the muscles of *M. cephalus* had the highest Cu concentration (6.87 mg kg⁻¹), while *P. lineatus* skin tissue had the highest Cu concentration (2.88 mg kg⁻¹). Cu is a known essential element in fish, plays a vital role in enzymatic activities and is required to synthesize hemoglobin. However, excessive intake will have negative health consequences [49]. Cu is an essential metal that facilitates iron release during hemoglobin synthesis. Cu can cause liver, kidney and brain damage and even death at high doses and with prolonged exposure [50]. Through mercapto and disulfide, Cu deposited in the liver forms tetrahedral metallothionein and metalloenzyme complex species [47]. Increased water transportation by boats and ships, repeated application of antifouling paint, oil discharge/leakage from ship engines, mining, electroplating, pesticides and commercial fishing operations are all associated with the rise in Cu concentrations in water [48, 51].

In this study, the Cr concentration in fish tissues ranged from 0.46 to 6.70 mg kg⁻¹, above the WHO MPL (0.15 mg kg⁻¹), US FDA (1 mg kg⁻¹) and BPOM RI (2.5 mg kg⁻¹). The highest Cr concentration was discovered in the liver of *M. cephalus* (6.70 mg kg⁻¹), while the lowest was in the skin of *P. lineatus* (0.46 mg kg⁻¹). The results of this study were higher than previous findings in *M. cephalus* liver by Usero *et al.* [26] in the San Carlos estuary of Spain (0.029 mg/kg) and Fazio *et al.* [35] in the Ionian Sea of Italy (0.1506 mg kg⁻¹) and in the Black Sea of Bulgaria (0.1303 mg/kg). *Arius sagor* had the highest Cr concentration in its muscles (2.58 mg kg⁻¹), whereas *P. lineatus* had the lowest (0.50 mg kg⁻¹). This result is higher than previous findings in *M. cephalus* muscles by Albuquerque *et al.* [3] in the Maranhao estuary of Brazil (0.11 mg kg⁻¹) and Fazio *et al.* [35] in the Ionian Sea of Italy (0.0213 mg kg⁻¹) and in the Black Sea of Bulgaria (0.1809 mg kg⁻¹). *Arius sagor* has the highest concentration of Cr in its skin (2.03 mg kg⁻¹). As a microelement with a critical role in glucose metabolism, Cr is considered a particularly hazardous pollutant. Although Cr is needed for glucose, lipid and protein metabolism, high concentrations of Cr are toxic to the liver, kidney and endocrine system [3]. Due to its higher availability, Cr is more readily absorbed by fish in estuaries, either directly from the water or indirectly through the food ingested. The rise in Cr concentration in waterways indicates that Cr is present in various oxidation states and is increasingly utilized in industrial applications [51]. Cr is produced by the mining, electroplating, textile and tannery industries [48]. For coastal communities that rely on local catches for sustenance, high concentrations of Cr in estuaries from anthropogenic causes are of great concern [3].

[35] study found that Pb concentrations in fish tissue ranged from 0.001 to 0.701mgkg⁻¹. The Pb concentration in the liver of *M. cephalus* was above the FAO MPL (0.5mgkg⁻¹) and [9] NI 7387:2009 (0.3mgkg⁻¹), but the values in *A. sagor* and *P. lineatus* were within the MPL. The highest Pb concentration was found in the liver of *Mugil cephalus* (0.701mgkg⁻¹) [13] while the lowest value was found in the muscle and skin of all fish species (0.81 mgkg⁻¹). The liver is presumably the target organ for Pb accumulation, resulting in low Pb concentrations in the muscles and skin. These data show lower Pb concentrations in the liver of *A. sagor* than in prior studies by Abdel-Warith *et al.* [52] in Saudi Arabia (5.54mgkg⁻¹), in *M. cephalus* liver by El-hak *et al.* [37] in Lake Manzala in Egypt (6.83mgkg⁻¹), but higher when compared to *M. cephalus* liver by Al Kazaghly *et al.* [42] off the Zliten coast of Libya (0.64mgkg⁻¹). This study found that the Pb concentrations in the muscles of *A. sagor* were lower than those in prior studies by Abdel-Warith *et al.* [52] in Saudi Arabia (1.27mgkg⁻¹), Pandion *et al.* [44] on the Kalpakkam coast of India (0.30mgkg⁻¹), El-hak *et al.* [37] in *M. cephalus* muscles (7.04mgkg⁻¹) in Lake Manzala in Egypt and Usero *et al.* [26] in the San Carlos estuary of Spain (0.04mgkg⁻¹). This study revealed lower Pb concentration [4] in the skin compared to an earlier study on *A. sagor* in Saudi Arabia (0.32mgkg⁻¹) [52]. Pb is the most common and widespread hazardous metal and it can be found in the civil construction industry, metal alloys, rainfall-runoff, refineries, marinas, fish processing industries, paint, pesticides, batteries, car emissions, mining and coal combustion [48]. Pb is highly toxic and harmful to human health. Toxicity in marine and estuarine organisms is proportional to the chemical form that is available and absorbed [7] with organic form absorption posing the most significant threat [23].

The Cd concentration [7] est fish tissues ranged between 0.001 and 0.072mgkg⁻¹, below the FAO MPL (0.50mgkg⁻¹). The Cd concentration in the muscles and skin of all fish samples was the same (0.001mgkg⁻¹); however, it differed in the livers of *Mugil cephalus* (0.041 and 0.048mgkg⁻¹) and *Arius sagor* (0.062 and 0.072mgkg⁻¹). Presumably, Cd is momentarily kept in the liver, where its concentration rises, before accumulating in the muscles and skin. According to Jakimska *et al.* [46], Cd is briefly stored in the liver; however, if the exposure is prolonged, it is transferred to the kidneys, where it is absorbed and bioaccumulated. This occurs because Cd forms a mixture with metallothionein. The results of this study were lower than previous findings in *M. cephalus* livers in the San Carlos estuary of Spain by Usero *et al.* [26] (0.14mgkg⁻¹), Al-Kazaghly *et al.* [42] off the Zliten coast of Libya (0.44mgkg⁻¹), Fazio *et al.* [35] in the Black Sea of Bulgaria (0.239mgkg⁻¹) and Hauser-Davis *et al.* [34] in the Itaipu beach of Brazil (0.13mgkg⁻¹) but higher than the findings of Fazio *et al.* [26] in the liver [20] *M. cephalus* in the Ionian Sea of Italy (0.0351mgkg⁻¹). The Cd concentration in muscle in this study was lower than previous findings by Fazio *et al.* [26] in *M. cephalus* muscle in the Black Sea estuary of Bulgaria (0.0391mgkg⁻¹) and Al-Kazaghly *et al.* [42] (0.06mgkg⁻¹) but higher than previous findings by Fazio *et al.* [35] in *M. cephalus* muscle in the Ionian Sea of Italy (0.0008mgkg⁻¹). Cd [44] is a highly hazardous element that is mainly transported by airborne particles. Because they are at the top of the marine food chain, fish, seabirds and marine mammals tend to collect large amounts of heavy metals in their livers. Cd poses a significant threat to the survival of estuarine and marine creatures since it can disrupt the reproductive system or even cause mortality. Its bioaccumulation in the human body can negatively affect the lungs, liver, bones, reproduction, kidneys and cancer [50]. The burning [3] fossil fuels (such as power plants, incineration and waste disposal sites, phosphate fertilizer factories and nonferrous metal smelting operations such as Cu, Pb and Zn), battery production, pigments, plastic stabilizers, welding, electroplating, pesticides, fertilizer and nuclear fission plants are associated with elevated concentrations of cadmium in water [33,48].

In this study, the Hg concentration in all tested tissues and fish species was 0.00004mgkg^{-1} , below the US FDA MPL (0.0005mgkg^{-1}). This suggests that Hg does not pollute coastal swamp waters and the concentration is very small in the fish tissues at the study location, because gold mining that uses Hg is few in number in South Kalimantan. Gold mining is more common in Central Kalimantan. In this study, Hg concentrations in the liver were lower than previous findings by Looi *et al.* [53] in the liver of *A. sagor* in the Strait of Malacca (0.48mgkg^{-1}), Usero *et al.* [26] in the liver of *M. cephalus* in San Carlos estuary of Spain (0.035mgkg^{-1}) and Al-Kazaghly *et al.* [42] in the liver of *M. cephalus* off the Zliten coast of Libya (0.31mgkg^{-1}). Meanwhile, the Hg concentration in muscles in this study was lower than previous findings by Morcillo *et al.* [54] in the muscles of *A. sagor* in Murcia of Spain (0.049mgkg^{-1}), in the Strait of Malacca (0.34mgkg^{-1}) by Looi *et al.* [53], in *M. cephalus* muscles in San Carlos estuary of Spain (0.010mgkg^{-1}) by Usero *et al.* [26], in Maranhao estuary of Brazil (0.17mgkg^{-1}) by Albuquerque *et al.* [3], off the Zliten coast of Libya (0.07mgkg^{-1}) by Al-Kazaghly *et al.* [42] and in Murcia of Spain (0.05mgkg^{-1}) by Morcillo *et al.* [54]. Even at low concentrations, Hg is highly toxic to numerous organisms. As the organ responsible for detoxifying, the liver typically has the most significant Hg concentrations [46]. The higher Hg concentrations in these species are presumably due to biomagnification along the food chain. Exposure to Hg can have unwanted effects, such as decreased movement coordination, vision loss, speech, hearing, walking impairments, memory loss and muscle weakness [50]. Waste from the hydroelectric, mining, pulp and paper, pesticide and battery industries increases Hg concentrations in estuaries [48]. In addition to emissions from coal-fired power plants, municipal and medical waste combustion contributes to high mercury concentrations [50]. Given the significance of coastal swamp habitats to biodiversity conservation, long-term monitoring and evaluation of Hg must remain a top priority.

In the study area, Fe, Cu and Cr concentrations that exceed the MPL suggested by national and international recommendations can be attributed to uncontrolled human activities. There is a need for additional research to identify point sources and assess their detrimental effects on fish. Depending on the species and organ, the concentration of heavy metals in fish can vary and excessive amounts can accumulate to dangerous levels. Two factors, known as abiotic and biotic factors, influence the bioaccumulation of heavy metals in fish bodies. The properties of heavy metals are influenced by abiotic variables, such as the nature of chemical elements, their concentration, speciation and the presence or absence of their biological function in the body. Although biotic factors pertain to species-specific vulnerability, life history, sex, age and diet, they are species-specific (including diet composition and metabolism). Active and passive processes eventually absorb metal contaminants. These factors influence the affinity or distribution, localization and bioaccumulation of metal contaminants in various tissues or organs, including the liver, kidneys, gills and to a lesser extent, muscles [6]. Physiochemical and biological factors, including pH, temperature, hardness, exposure duration, species feeding habits and habitat complexity, further influence the accumulation of heavy metals in fish organs. In addition to size, sex, reproductive cycle, feeding preferences, swimming patterns and environment, other characteristics affect heavy metal bioaccumulation [6]. Cunningham *et al.* [33] added that species differences (intrinsic) and environmental factors (extrinsic) influence the accumulation of metals in fish. Intrinsic species-dependent characteristics, such as trophic state and feeding tactics, are associated with fundamental food choice patterns (e.g., algae, zooplankton, invertebrates, fish). In addition, intrinsic characteristics include the fish's age and the sexes and phases of sexual reproduction, which vary depending on the season in which the fish were captured. Extrinsic factors that influence metal bioaccumulation include the chemical form of the metal, the degree of contamination and the presence of other pollutants in

the aquatic environment. Extrinsic factors include water quality features such as salinity, temperature, dissolved oxygen content, pH, total suspended particles and bottom sediment quality, which can serve as reservoirs for metal adsorption and resuspension [50].

Conclusions

South Kalimantan's Kuala Lupak and Kuala Tambangan coastal swamp waters are contaminated with heavy metals in the following order: Fe > Cu > Zn > Cr > Pb > Cd > Hg. The Fe concentration surpasses the international and national MPLs, whereas the Cu and Cr amounts only exceed the national MPL. Heavy metal concentration examinations were also conducted on commercial fish typically consumed by coastal communities, such as *Mugil cephalus*, *Arius sagus* and *Plotosus lineatus*, with the following findings: Fe > Zn > Cu > Cr > Pb > Cd > Hg. Based on these data, *Mugil cephalus* has the largest bioaccumulation of heavy metals, followed by *Arius sagor* and *Plotosus lineatus*. In the bodies of *Mugil cephalus*, *Arius sagor* and *Plotosus lineatus*, the Fe, Cu and Cr concentrations surpassed the international and national MPLs. The patterns of heavy metal concentrations in the liver, muscle and skin tissues of the three test fish species were also analyzed. According to these findings, liver is the most bio-accumulative of heavy metals, followed by skin and muscle, particularly in the liver tissue of *M. cephalus*, which collects the highest concentrations of Fe, Cu, Cr, Zn and Pb. In the coastal swamp waters of Kuala Lupak and Kuala Tambangan, South Kalimantan, Indonesia, the predominant heavy metal is Fe, which pollutes the water and commercial fish species.

Long-term ingestion of fish with a high concentration of heavy metals can result in neurological, carcinogenic, nephrological and immunological issues, among others. Therefore, the concentrations of heavy metals, particularly Fe, Cu and Cr, should be monitored continually in the potentially polluted coastal swamps of South Kalimantan, as they tend to concentrate on the liver, skin and muscles. Furthermore, waste disposal from industrial activity and other sources must be regulated. In addition, routine evaluations of heavy metal concentrations in commercial fish species are required to better comprehend the pollution conditions of estuaries and coastlines and the accompanying human health hazards. To protect public health, it is necessary to prohibit fish consumption from water bodies contaminated with mining effluent. The severity of heavy metal bioaccumulation in fish tissues necessitates immediate, sustained and targeted actions by both government authorities and the local scientific community of research and development of mining agencies to assist in preventing and mitigating the situation and ensuring the physical well-being of the local population. Consideration must also be given to enforcing laws requiring the detoxification of effluents before their release into natural water bodies to protect aquatic ecosystems. To prevent the impacts of hazardous trace elements on human health, there is an immediate need for strong government regulation and continuous monitoring of trace metals in fish from mining areas.

Acknowledgments

This study is a part of the Regular Fundamental Research grant during the initial year of a two-year project focused on the mitigation of heavy metal pollution in the coastal swamp ecosystems of South Kalimantan. This research specifically investigates the use of biomonitoring methods and fish bioindicators to address this issue. The present study was financially supported by the Directorate of Research, Technology and Community Service, Ministry of Education, Culture, Research and Technology of the Republic of Indonesia, during the fiscal year 2023. The research was conducted under contract number 130/E5/PG.02.00.

PL/2023, issued on June 19, 2023. Best regards to Auliya Reni Hadisa, Dinda Triana and Muhammad Faiza Rahman for their assistance with collecting and preparing samples.

References

- [1] J. Pandiyan, S. Mahboob, M. Govindarajan, K.A. Al-Ghanim, Z. Ahmed, N. Al-Mulhm, R. Jagadheesan, K. Krishnappa, *An assessment of level of heavy metals pollution in the water, sediment and aquatic organisms: A perspective of tackling environmental threats for food security*, **Saudi Journal of Biological Sciences**, **28**(2), 2021, pp. 1218–1225. DOI: 10.1016/j.sjbs.2020.11.072.
- [2] E. Solgi, M.M. Galangashi, *Assessing the health of marine and lacustrine wetland using measurement of heavy metals in fish species: Case study from two Iranian international wetland (Gomishan and Zarivar)*, **Environmental Nanotechnology, Monitoring and Management**, **10**, 2018, pp. 73–78. <https://doi.org/10.1016/j.enmm.2018.05.004>.
- [3] K.F.D. Albuquerque, M.H.L. Silva, J.W.D. Azevedo, L.S. Soares, A.M. Bandeira, L.A. Soares, A.C.L. de Castro, *Assessment of water quality and concentration of heavy metals in fishes in the estuary of the Perizes River, Gulf of Maranhao, Brazil*, **Marine Pollution Bulletin**, **186**, 2023, Article Number: 114420. DOI: 10.1016/j.marpolbul.2022.114420.
- [4] M.H.L. Silva, A.C.L. de Castro, I.S. da Silva, P.F.P. Cabral, J.W.D. Azevedo, L.S. Soares, A.M. Bandeira, M.J. Basso, J.L.S. Nunes, *Determination of metals in estuarine fishes in a metropolitan region of the coastal zone of the Brazilian Amazon*, **Marine Pollution Bulletin**, **186**, 2023, Article Number: 114477. DOI: 10.1016/j.marpolbul.2022.114477.
- [5] T. Aytekin, D. Kargin, H.Y. Çogun, O. Temiz, H.S. Varkal, F. Kargin, *Accumulation and health risk assessment of heavy metals in tissues of the shrimp and fish species from the Yumurtalik coast of Iskenderun Gulf, Turkey*, **Heliyon**, **5**(8), 2019, Article Number: e02131. DOI: 10.1016/j.heliyon.2019.e02131
- [6] M. Pragnya, S.D. Kumar, A.J.S. Raju, L.N. Murthy, *Bioaccumulation of heavy metals in different organs of Labeo rohita, Pangasius hypophthalmus, and Katsuwonus pelamis from Visakhapatnam, India*, **Marine Pollution Bulletin**, **157**, 2020, Article Number: 111326. DOI: 10.1016/j.marpolbul.2020.111326.
- [7] P. Adani, A.A. Sawale, G. Nandhagopal, *Bioaccumulation of heavy metals in the food components from water and sediments in the coastal waters of Kalpakkam, Southeast coast of India*, **Environmental Nanotechnology Monitoring and Management**, **17**, 2022, Article Number: 100627. <https://doi.org/10.1016/j.enmm.2021.100627>.
- [8] M.M.N. Authman, M.S. Zaki, E.A. Khallaf, H.H. Abbas, *Use of Fish as Bio-indicator of the Effects of Heavy Metals Pollution*, **Journal of Aquaculture Research and Development**, **6**(4), 2015, pp. 328-340. DOI:10.4172/2155-9546.1000328
- [9] P. Tanhan, K. Imsilp, N. Lansubsakul, W. Thong-asa, *Oxidative response to Cd and Pb accumulation in coastal fishes of Pattani Bay*, **Italian Journal of Animal Science**, **22**(1), 2023, pp. 148–156. DOI: 10.1080/1828051X.2023.2166430.
- [10] S.A. Mustafa, *Histopathology and heavy metal bioaccumulation in some tissues of Luciobarbus xanthopterus collected from Tigris River of Baghdad, Iraq*, **Egyptian Journal of Aquatic Research**, **46**(2), 2020, pp. 123-129. DOI: 10.1016/j.ejar.2020.01.004.
- [11] C.N. Amadi, C. Frazzoli, O.E. Orisakwe, *Sentinel species for biomonitoring and biosurveillance of environmental heavy metals in Nigeria*, **Journal of Environmental Science and Health, Part C-Toxicology and Carcinogenesis**, **38**(1), 2020, pp. 21–60. DOI: 10.1080/26896583.2020.1714370.

- [12] H.B. Santoso, Hidayaturrahmah, E. Suhartono, R. Yunita, D. Biyatmoko, D. Anggraini, *Heavy metal concentrations in water, sediment and giant mudskipper (*Periophthalmodon schlosseri*) in the coastal wetlands of Kuala Lupak estuary of the Barito River, Indonesia*, **AAFL Bioflux**, **14**(5), 2021, pp. 2878–2893.
- [13] Z. Kljaković-Gašpić, A. Sekovanić, T. Orct, D. Šebešćen, E. Klasiček, D. Zanella, *Potentially Toxic Elements in Water, Sediments and Fish from the Karstic River (Raša River, Croatia) Located in the Former Coal-Mining Area*, **Toxics**, **11**(1), 2023, Article Number: 42. DOI: 10.3390/toxics11010042.
- [14] A.E. Nuringtyas, A.P. Larasati, F. Septian, I. Mulyana, W. Israwati, A.Z.A. Mourniaty, W. Nainggolan, R. Suharti, M.A. Jabbar, *Aspek Biologi Ikan Belanak (*Mugil cephalus*) di Perairan Teluk Banten*, **Buletin Jalanidhitah Sarva Jivitam**, **1**(2), 2019, pp. 81-87. DOI: 10.15578/bjsj.v1i2.8423
- [15] Y.R. Dewanti, Irwani, S. Rejeki, *Studi reproduksi dan morfometri ikan sembilang (*Plotosus canius*) betina yang didaratkan di pengepul wilayah Krobokan Semarang*, **Journal of Marine Research**, **1**(2), 2012, pp. 135–144. <https://doi.org/10.14710/jmr.v1i2.2030>.
- [16] A.P.A. Samad, M. Jamil, C. Gustiana, S. Anzitha, T. Mahyuddin, B. Baihaqi, *Evaluation on the growth pattern and biological conditions of Sembilang (*Plotosus canius*) in East Aceh*, **IOP Conference Series: Earth Environment Science**, **1115**(1), 2022, Article Number: 012006. DOI: 10.1088/1755-1315/1115/1/012006.
- [17] Y. Rustandi, A. Fahrudin, Zairion, M.N. Arkham, A. Zahid, A. Hamdani, A. Ramli, A. Trihandoyo, A. *Distribution fisheries resources of Small Island in Estuary Area: An assessment in Bunyu Island, North Kalimantan, Indonesia*, **IOP Conference Series: Earth and Environment Science**, **241**, 2019, Article Number: 012003. DOI: 10.1088/1755-1315/241/1/012003.
- [18] G. Kasmianti, R.A. Sakinah, B. Yudono, *The Analysis of Manganese (Mn) in Waste Water Treatment (IPAL) of Coal Mine of PT Bukit Asam Indonesia*, **Indonesian Journal of Fundamental and Applied Chemistry**, **6**(2), 2021, pp. 53–58. DOI: 10.24845/ijfac.v6.i2.53.
- [19] D. Sofarini, A. Rahman, I. Ridwan, *Permodelan Uji Logam Berat pada Badan Air, Biota dan Sedimen di Perairan Muara DAS Barito*, **Bumi Lestari**, **12**(1), 2012, pp. 32–44. <https://jurnal.harianregional.com/blje/id-1532>.
- [20] R.S.B. Waspodo, M.I. Sahana, *River water quality modelling in Barito watershed*, **IOP Conference Series: Earth and Environment Science**, **399**, 2019, Article Number: 012018. DOI: 10.1088/1755-1315/399/1/012018.
- [21] M. Jayaprakash, R.S. Kumar, L. Giridharan, S.B. Sujitha, S.K. Sarkar, M.P. Jonathan, *Bioaccumulation of metals in fish species from water and sediments in macrotidal Ennore creek, Chennai, SE coast of India: A metropolitan city effect*, **Ecotoxicology Environmental Safety**, **120**, 2015, pp. 243–255. DOI: 10.1016/j.ecoenv.2015.05.042.
- [22] M. Kottelat, A.J. Whitten, S.N. Kartikasari, S. Wirjoatmodjo, **Freshwater Fishes of Western Indonesia and Sulawesi**, Periplus Editions, Hong Kong, 1993, 221p. ISBN: 0945971605
- [23] M. Akila, S. Anbalagan, N.M. Lakshmisri, V. Janaki, T. Ramesh, R.J. Merlin, S. Kamala-Kannan, *Heavy metal accumulation in selected fish species from Pulicat Lake, India, and health risk assessment*, **Environmental Technology and Innovation**, **26**, 2022, Article Number: 102744. DOI: 10.1016/j.eti.2022.102744.
- [24] S. Mehar, I. Anam, Z. Masood, S. Alvi, W. Khan, M. Kabir, M. Shahbaz, T. Khan, *Original Bioaccumulation of heavy metals in the different tissues of Mackerel scad, *Decapterus macarellus* (Cuvier, 1833) collected from Karachi and Gwadar Coasts of*

- Pakistan*, **Saudi Journal of Biological Sciences**, **30**(2), 2023, Article Number: 103540. DOI: 10.1016/j.sjbs.2022.103540.
- [25] H. Fan, S. Chen, Z. Li, P. Liu, C. Xu, X. Yang, *Assessment of heavy metals in water, sediment and shellfish organisms in typical areas of the Yangtze River Estuary, China*, **Marine Pollution Bulletin**, **151**, 2020, Article Number: 110864. DOI: 10.1016/j.marpolbul.2019.110864.
- [26] J. Usero, C. Izquierdo, J. Morillo, I. Gracia, *Heavy metals in fish (Solea vulgaris, Anguilla anguilla and Liza aurata) from salt marshes on the southern Atlantic coast of Spain*, **Environment International**, **29**(7), 2004, pp. 949–956. DOI: 10.1016/S0160-4120(03)00061-8.
- [27] T. Dalu, R. Tshivhase, R.N. Cuthbert, F.M. Murungweni, R.J. Wasserman, *Metal Distribution and Sediment Quality Variation across Sediment Depths of a Subtropical Ramsar Declared Wetland*, **Water**, **12**(10), 2020, Article Number: 2779. DOI: 10.3390/w12102779.
- [28] M. Jaishankar, T. Tseten, N. Anbalagan, B.B. Mathew, K.N. Beeregowda, *Toxicity mechanism and health effects of some heavy metals*, **Interdisciplin Toxicology**, **7**(2), 2014, pp. 60–72. DOI: 10.2478/intox-2014-0009.
- [29] A.A.D. Machado, K. Spencer, W. Kloas, M. Toffolon, C. Zarfl, *Metal fate and effects in estuaries: A review and conceptual model for better understanding of toxicity*, **Science of the Total Environment**, **541**, 2016, pp. 268–281. DOI: 10.1016/j.scitotenv.2015.09.045.
- [30] N. Zaoui, N. Bouriga, I. Louiz, W.R. Bahri, N. Saadaoui, J.P. Quiganrd, M. Trabelsi, M. Ben-Attia, A.A.B. Shahin, *Assessment of effects of metal contamination and abiotic factors on fatty acid composition and biochemical biomarkers activity in the liver of Chelon ramada collected from two ecosystems on the Mahdia coast, Tunisia*, **Biochemical Systematics and Ecology**, **107**, 2023, Article Number: 104608. DOI: 10.1016/j.bse.2023.104608.
- [31] A. Yogeshwaran, K. Gayathiri, T. Muralisankar, V. Gayathri, J.I. Monica, R. Rajaram, K. Marimuthu, P.S. Bhavan, *Bioaccumulation of heavy metals, antioxidants, and metabolic enzymes in the crab Scylla serrata from different regions of Tuticorin, Southeast Coast of India*, **Marine Pollution Bulletin**, **158**, 2020, Article Number: 111443. <https://doi.org/10.1016/j.marpolbul.2020.111443>.
- [32] N. Soltani, M. Marengo, B. Keshavarzi, F. Moore, P.S. Hooda, M.R. Mahmoudi, S. Gobert, *Occurrence of trace elements (TEs) in seafood from the North Persian Gulf: Implications for human health*, **Journal of Food Composition and Analysis**, **97**, 2021, Article Number: 103754. DOI: 10.1016/j.jfca.2020.103754.
- [33] P.A. Cunningham, E.E. Sullivan, K.H. Everett, S.S. Kovach, A. Rajan, M.C. Barber, *Assessment of metal contamination in Arabian/Persian Gulf fish: A review*, **Marine Pollution Bulletin**, **143**, 2019, pp. 264–283. DOI: 10.1016/j.marpolbul.2019.04.007.
- [34] K.C. Lakra, T.K. Banerjee, B. Lal, *Coal mine effluent-induced metal bioaccumulation, biochemical, oxidative stress, metallothionein, and histopathological alterations in vital tissues of the catfish, Clarias batrachus*, **Environmental Science and Pollution Research**, **28**(20), 2021, pp. 25300–25315. DOI: 10.1007/s11356-021-12381-3.
- [35] F. Fazio, C. D'Iglio, G. Capillo, C. Saoca, K. Peycheva, G. Piccione, L. Makedonski, *Environmental investigations and tissue bioaccumulation of heavy metals in grey mullet from the Black Sea (Bulgaria) and the Ionian Sea (Italy)*, **Animals**, **10**(10), 2020 Article Number: 1739. DOI: 10.3390/ani10101739.
- [36] H. Karadede, S.A. Oymak, E. Ünlü, *Heavy metals in mullet, Liza abu, and catfish, Silurus triostegus, from the Atatürk Dam Lake (Euphrates), Turkey*, **Environment International**, **30**(2), 2004, pp. 183–188. DOI: 10.1016/S0160-4120(03)00169-7.

- [37] H.N.G. El-hak, M.I.S. El-Din, R.A. Elrayess, *Bioaccumulation of heavy metals and their histopathological impact on Mugil cephalus from the North Eastern Region of Manzala Lake, Egypt*, **Regional Studies in Marine Science**, **45**, 2021, Article Number: 101841. DOI: 10.1016/j.rsma.2021.101841.
- [38] M.M. Ikram, A. Ismail, C.K. Yap, A.A.N. Azwady, *Levels of heavy metals (Zn, Cu, Cd, and Pb) in mudskippers (Periophthalmodon schlosseri) and sediments collected from intertidal areas at Morib and Remis, Peninsular Malaysia*, **Toxicological and Environmental Chemistry**, **92**(8), 2010, pp. 1471–1486. DOI: 10.1080/02772241003614304.
- [39] A. Anandkumar, R. Nagarajan, K. Prabakaran, C.H. Bing, R. Rajaram, *Human health risk assessment and bioaccumulation of trace metals in fish species collected from the Miri coast, Sarawak, Borneo*, **Marine Pollution Bulletin**, **133**, 2018, pp. 655–663. DOI: 10.1016/j.marpolbul.2018.06.033.
- [40] E. Uluturhan, F. Kucuksezgin, *Heavy metal contaminants in Red Pandora (Pagellus erythrinus) tissues from the Eastern Aegean Sea, Turkey*, **Water Research**, **41**(6), 2007, pp. 1185–1192. DOI: 10.1016/j.watres.2006.11.044.
- [41] T.K. Banerjee, A.K. Mittal, *Histopathological studies on the repair of the excised skin wounds of the air-breathing scalyfish Channa striata (Bloch)*, **Current Science**, **77**(8), 1999, pp. 1067–1075.
- [42] R.F. Al-Kazaghly, M. Hamid, K.A. Ighwela, *Bioaccumulation of some heavy metals in Red mullet (Mullus barbatus), and common Pandora (Pagellus erythrinus) from Zliten coast, Libya*, **Jurnal Ilmiah Perikanan dan Kelautan**, **13**(1), 2021, pp. 81–86. DOI: 10.20473/jipk.v13i1.22037.
- [43] R.A. Hauser-Davis, I.C.A.C. Bordon, T.F. Oliveira, R.L. Ziolli, *Metal bioaccumulation in edible target tissues of mullet (Mugil liza) from a tropical bay in Southeastern Brazil*, **Journal of Trace Elements in Medicine and Biology**, **36**, 2016, pp. 38–43. DOI: 10.1016/j.jtemb.2016.03.016.
- [44] K. Pandion, K.D. Arunachalam, R. Rajagopal, D. Ali, S. Alarifi, S.W. Chang, B. Ravindran, *Health risk assessment of heavy metals in the seafood at Kalpakkam coast, Southeast Bay of Bengal*, **Marine Pollution Bulletin**, **189**, 2023, Article Number: 114766. DOI: 10.1016/j.marpolbul.2023.114766.
- [45] K.D. Bastami, M. Afkhami, M. Mohammadzadeh, M. Ehsanpour, S. Chambari, S. Aghaei, M. Esmailzadeh, M.R. Neyestani, F. Lagzaee, M. Baniamam, *Bioaccumulation and ecological risk assessment of heavy metals in the sediments and mullet Liza klunzingeri in the northern part of the Persian Gulf*, **Marine Pollution Bulletin**, **94**(1–2), 2015, pp. 329–334. DOI: 10.1016/j.marpolbul.2015.01.019.
- [46] A. Jakimska, P. Konieczka, K. Skóra, J. Namieśnik, *Bioaccumulation of Metals in Tissues of Marine Animals, Part I: the Role and Impact of Heavy Metals on Organisms*, **Polish Journal of Environmental Studies**, **20**(5), 2011, pp. 1117–1125.
- [47] M.H. Al-Yousuf, M.S. El-Shahawi, S.M. Al-Ghais, *Trace metals in liver, skin and muscle of Lethrinus lentjan fish species in relation to body length and sex*, **Science of the Total Environment**, **256**(2–3), 2000, pp. 87–94. DOI: 10.1016/S0048-9697(99)00363-0.
- [48] D. Paul, *Research on heavy metal pollution of river Ganga: A review*, **Annals of Agrarian Science**, **15**(2), 2017, pp. 278–286. <https://doi.org/10.1016/j.aasci.2017.04.001>.
- [49] A. Velusamy, P.S. Kumar, A. Ram, S. Chinnadurai, *Bioaccumulation of heavy metals in commercially important marine fishes from Mumbai Harbor, India*, **Marine Pollution Bulletin**, **81**(1), 2014, pp. 218–224. DOI: 10.1016/j.marpolbul.2014.01.049.
- [50] N. Alizada, S. Malik, S. Bin Muzaffar, *Bioaccumulation of heavy metals in tissues of Indian anchovy (Stolephorus indicus) from the UAE coast, Arabian Gulf*, **Marine**

- Pollution Bulletin**, **154**, 2020, Article Number: 111033. DOI: 10.1016/j.marpolbul.2020.111033.
- [51] S. Rajeshkumar, X. Li, *Bioaccumulation of heavy metals in fish species from the Meiliang Bay, Taihu Lake, China*, **Toxicology Reports**, **5**, 2018, pp. 288–295. DOI: 10.1016/j.toxrep.2018.01.007.
- [52] A.W.A. Abdel-Warith, E.S.M.I. Younis, N.A. Al-Asgah, A.M. Rady, H.Y. Allam, *Bioaccumulation of lead nitrate in tissues and its effects on hematological and biochemical parameters of Clarias gariepinus*, **Saudi Journal of Biological Sciences**, **27**(3), 2020, pp. 840–845. DOI: 10.1016/j.sjbs.2020.01.015.
- [53] L.J. Looi, A.Z. Aris, H. Haris, F.M. Yusoff, Z. Hashim, *The levels of mercury, methylmercury and selenium and the selenium health benefit value in grey-eel catfish (Plotosus canius) and giant mudskipper (Periophthalmodon schlosseri) from the Strait of Malacca*, **Chemosphere**, **152**, 2016, pp. 265–273. DOI: 10.1016/j.chemosphere.2016.02.126.
- [54] P. Morcillo, M.A. Esteban, A. Cuesta, *Mercury and its toxic effects on fish*, **AIMS Environmental Science**, **4**(3), 2017, pp. 386–402. DOI: 10.3934/environsci.2017.3.386.
-

Received: October 02, 2023

Accepted: May 24, 2024

ASSESSING THE HEALTH OF SOUTH KALIMANTAN COASTAL SWAMP WETLANDS USING MEASUREMENTS OF HEAVY METALS IN COMMERCIAL FISH SPECIES

ORIGINALITY REPORT

17%

SIMILARITY INDEX

0%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Nuray Alizada, Shaima Malik, Sabir Bin Muzaffar. "Bioaccumulation of heavy metals in tissues of Indian anchovy (*Stolephorus indicus*) from the UAE coast, Arabian Gulf", *Marine Pollution Bulletin*, 2020
Publication 1%
- 2 "Crop Production under Stressful Conditions", Springer Science and Business Media LLC, 2018
Publication 1%
- 3 Patricia A. Cunningham, Elizabeth E. Sullivan, Kibri H. Everett, Samuel S. Kovach, Anbiah Rajan, Mary C. Barber. "Assessment of metal contamination in Arabian/Persian Gulf fish: A review", *Marine Pollution Bulletin*, 2019
Publication 1%
- 4 Keila Fernanda de Melo Albuquerque, Marcelo Henrique Lopes Silva, James Werllen de Jesus Azevedo, Leonardo Silva Soares et al. "Assessment of water quality and concentration of heavy metals in fishes

in the estuary of the Perizes River, Gulf of Maranhão, Brazil", Marine Pollution Bulletin, 2023

Publication

5

Kalpana Chhaya Lakra, Tarun Kumar Banerjee, Bechan Lal. "Coal mine effluent-induced metal bioaccumulation, biochemical, oxidative stress, metallothionein, and histopathological alterations in vital tissues of the catfish, *Clarias batrachus*", Environmental Science and Pollution Research, 2021

1 %

Publication

6

Eisa Solgi, Mohsen Mohammadi Galangashi. "Assessing the health of marine and lacustrine wetland using measurement of heavy metals in fish species: Case study from two Iranian international wetland (Gomishan and Zarivar)", Environmental Nanotechnology, Monitoring & Management, 2018

<1 %

Publication

7

Anandkumar A., Nagarajan R., Prabakaran K., Chua Han Bing, Rajaram R.. "Human health risk assessment and bioaccumulation of trace metals in fish species collected from the Miri coast, Sarawak, Borneo", Marine Pollution Bulletin, 2018

<1 %

Publication

8

Megha Chhabra, Bhagwati Sharan, Manoj Kumar. "A fuzzy cognitive map of the quality of user experience determinants in mobile application design", Journal of Intelligent & Fuzzy Systems, 2023

Publication

<1 %

9

Luana S. Moreira, Ana P. R. Santana, Floriatan S. Costa, Josiane B. Vera et al. "Enhanced extraction of arsenic and cadmium from environmental samples using a natural deep eutectic solvent and determination by inductively coupled plasma mass spectrometry", International Journal of Environmental Analytical Chemistry, 2020

Publication

<1 %

10

Aldoghachi, Mohammed Abdulridha Jasim. "Toxicity Assessment and Bioaccumulation of Heavy Metals in Red Tilapia", University of Malaya (Malaysia), 2023

Publication

<1 %

11

M. Akila, S. Anbalagan, N.M. Lakshmisri, V. Janaki, T. Ramesh, R. Jancy Merlin, S. Kamala-Kannan. "Heavy metal accumulation in selected fish species from Pulicat Lake, India, and health risk assessment", Environmental Technology & Innovation, 2022

Publication

<1 %

12

M. Pragnya, S. Dinesh Kumar, A.J. Solomon Raju, L.N. Murthy. "Bioaccumulation of heavy metals in different organs of *Labeo rohita*, *Pangasius hypophthalmus*, and *Katsuwonus pelamis* from Visakhapatnam, India", *Marine Pollution Bulletin*, 2020

Publication

<1 %

13

Hassan Sheikhzadeh, Amir Hossein Hamidian. "Bioaccumulation of heavy metals in fish species of Iran: a review", *Environmental Geochemistry and Health*, 2021

Publication

<1 %

14

Zohra Ben Salem, Nicolas Capelli, Xavier Laffray, Grisey Elise, Habib Ayadi, Lotfi Aleya. "Seasonal variation of heavy metals in water, sediment and roach tissues in a landfill draining system pond (Etueffont, France)", *Ecological Engineering*, 2014

Publication

<1 %

15

Hashem Khandan Barani, Mohammad Sadegh Alavi-Yeganeh, Alireza Riyahi Bakhtiari. "Metals bioaccumulation, possible sources and consumption risk assessment in five Sillaginid species, a case study: Bandar Abbas (Persian Gulf) and Chabahar Bay (Oman Sea), Iran", *Marine Pollution Bulletin*, 2023

Publication

<1 %

16

Jayaprakash, M., R. Senthil Kumar, L. Giridharan, S.B. Sujitha, S.K. Sarkar, and M.P. Jonathan. "Bioaccumulation of metals in fish species from water and sediments in macrotidal Ennore creek, Chennai, SE coast of India: A metropolitan city effect", *Ecotoxicology and Environmental Safety*, 2015.

Publication

<1 %

17

Md. Abu Rayhan Khan, Muhammad Shamim Al Mamun, Mosummath Hosna Ara. "Health Risk Assessment of Liza parsia Fish Lipids: Heavy Metals and DDTs Contaminations", *Iranian Journal of Toxicology*, 2021

Publication

<1 %

18

Sivakumar Rajeshkumar, Xiaoyu Li. "Bioaccumulation of heavy metals in fish species from the Meiliang Bay, Taihu Lake, China", *Toxicology Reports*, 2018

Publication

<1 %

19

Amir Hossein Hamidian, Hassan Sheikhzadeh, Azadeh Boujari, Soheil Eagderi, Sohrab Ashrafi. "Comparative assessment of human health risk associated with heavy metals bioaccumulation in fish species (*Barbus grypus* and *Tenualosa ilisha*) from the Karoon River, Iran: Elucidating the role of habitat and feeding habits", *Marine Pollution Bulletin*, 2023

Publication

<1 %

20

Fazel Abdolahpur Monikh, Alireza Safahieh, Ahmad Savari, Abdolmajid Doraghi. "Heavy metal concentration in sediment, benthic, benthopelagic, and pelagic fish species from Musa Estuary (Persian Gulf)", Environmental Monitoring and Assessment, 2012

Publication

<1 %

21

Hülya Karadede, Seyit Ahmet Oymak, Erhan Ünlü. "Heavy metals in mullet, Liza abu, and catfish, Silurus triostegus, from the Atatürk Dam Lake (Euphrates), Turkey", Environment International, 2004

Publication

<1 %

22

Andi Kurniawan, Siti Mariyah Ulfa, Chamidah Chamidah. "The Biosorption of Copper(II) Using a Natural Biofilm Formed on the Stones from the Metro River, Malang City, Indonesia", International Journal of Microbiology, 2022

Publication

<1 %

23

Ary Marchellina, Agoes Soegianto, Trisnadi Widyaleksono Catur Putranto, Wildanun Mukholladun et al. "An assessment of the potential health hazards associated with metal contamination in a variety of consumable species living along the industrialized coastline of East Java, Indonesia", Marine Pollution Bulletin, 2024

Publication

<1 %

24

Heri B. Santoso, Eko Suhartono, Rizmi Yunita, Danang Biyatmoko. "Mudskipper Fish as a Bio-indicator for Heavy Metals Pollution in a Coastal Wetland", Egyptian Journal of Aquatic Biology and Fisheries, 2020

Publication

<1 %

25

Meltem Dural, M. Ziya Lugal Göksu, Argun Akif Özak. "Investigation of heavy metal levels in economically important fish species captured from the Tuzla lagoon", Food Chemistry, 2007

Publication

<1 %

26

Abdolahpur Monikh, F., A. Maryamabadi, A. Savari, and K. Ghanemi. "Heavy metals' concentration in sediment, shrimp and two fish species from the northwest Persian Gulf", Toxicology and Industrial Health, 2013.

Publication

<1 %

27

Arockia Vasanthi, Lourduraj, Peranandam Revathi, Jayaprakash Mini, and Natesan Munuswamy. "Integrated use of histological and ultrastructural biomarkers in Mugil cephalus for assessing heavy metal pollution in Ennore estuary, Chennai", Chemosphere, 2013.

Publication

<1 %

28

Bragazza, L.. "Heavy metals in bog waters: An alternative way to assess atmospheric

<1 %

precipitation quality?", Global and Planetary Change, 200610

Publication

29

Ping Zhuang, Zhi-an Li, Murray B. McBride, Bi Zou, Gang Wang. "Health risk assessment for consumption of fish originating from ponds near Dabaoshan mine, South China", Environmental Science and Pollution Research, 2013

Publication

<1 %

30

Bibhu Prasad Panda, Yugal Kishore Mohanta, Siba Prasad Parida, Abanti Pradhan et al. "Metal pollution in freshwater fish: A key indicator of contamination and carcinogenic risk to public health", Environmental Pollution, 2023

Publication

<1 %

31

Md. Simul Bhuyan, Sayeed Mahmood Belal Haider, Gowhar Meraj, Muhammad Abu Bakar et al. "Assessment of Heavy Metal Contamination in Beach Sediments of Eastern St. Martin's Island, Bangladesh: Implications for Environmental and Human Health Risks", Water, 2023

Publication

<1 %

32

Oluniyi Solomon Ogunola, Olawale Ahmed Onada, Augustine Eyiunmi Falaye. "Ecological Risk Evaluation of Biological and Geochemical Trace Metals in Okrika

<1 %

33

Bilandzic, N.. "Metal content determination in four fish species from the Adriatic Sea", *Food Chemistry*, 20110201

Publication

<1 %

34

Francesco Fazio, Claudio D'Iglio, Gioele Capillo, Concetta Saoca, Katya Peycheva, Giuseppe Piccione, Lubomir Makedonski. "Environmental Investigations and Tissue Bioaccumulation of Heavy Metals in Grey Mullet from the Black Sea (Bulgaria) and the Ionian Sea (Italy)", *Animals*, 2020

Publication

<1 %

35

Francesco Fazio, Concetta Saoca, Marilena Sanfilippo, Gioele Capillo, Nunziacarla Spanò, Giuseppe Piccione. "Response of vanadium bioaccumulation in tissues of *Mugil cephalus* (Linnaeus 1758)", *Science of The Total Environment*, 2019

Publication

<1 %

36

Sivakumar Rajeshkumar, Yang Liu, Xiangyang Zhang, Boopalan Ravikumar, Ge Bai, Xiaoyu Li. "Studies on seasonal pollution of heavy metals in water, sediment, fish and oyster from the Meiliang Bay of Taihu Lake in China", *Chemosphere*, 2018

Publication

<1 %

37

Marcelo Henrique Lopes Silva, Antonio Carlos Leal de Castro, Iranaldo Santos da Silva, Patrícia Fernanda Pereira Cabral et al. "Determination of metals in estuarine fishes in a metropolitan region of the coastal zone of the Brazilian Amazon", *Marine Pollution Bulletin*, 2023

Publication

<1 %

38

Param Adani, Amol Ashok Sawale, Ganesan Nandhagopal. "Bioaccumulation of heavy metals in the food components from water and sediments in the coastal waters of Kalpakkam, Southeast coast of India", *Environmental Nanotechnology, Monitoring & Management*, 2022

Publication

<1 %

39

Roveri, Vinicius. "An Integrated Environmental Assessment of the Water and Sediments from the Coastal Areas of Guarujá, São Paulo, Brazil: a Physico-Chemical, Biological and Ecotoxicological Approach", *Universidade Fernando Pessoa (Portugal)*, 2024

Publication

<1 %

40

Sintje Liline, Dominggus Rumahlatu, Sriyanti Salmanu, Mery Pattipeilohy, Kristin Sangur. "Bioaccumulation of Chromium, Iron, and the Expression of TNF- α and Caspase-3 in Mudskipper (*Periophthalmus* spp.) from

<1 %

41

Cecilia Nwadiuto Amadi, Chiara Frazzoli, Orish Ebere Orisakwe. "Sentinel species for biomonitoring and biosurveillance of environmental heavy metals in Nigeria", Journal of Environmental Science and Health, Part C, 2020

Publication

<1 %

42

Kazim Uysal. "The comparison of heavy metal accumulation ratios of some fish species in Enne Dame Lake (Kütahya/Turkey)", Environmental Monitoring and Assessment, 10/09/2008

Publication

<1 %

43

Oluwadamilare Emmanuel Obayemi, Mary Adebola Ayoade, Olaniyi Olusola Komolafe. "Health risk assessment of heavy metals in Coptodon zillii and Parachanna obscura from a tropical reservoir", Heliyon, 2023

Publication

<1 %

44

Said Muhammad, Kashif Ahmad. "Heavy metal contamination in water and fish of the Hunza River and its tributaries in Gilgit-Baltistan: Evaluation of potential risks and provenance", Environmental Technology & Innovation, 2020

Publication

<1 %

45

Zohra Ben Salem, Habib Ayadi. "Heavy metal accumulation in *Diplodus annularis*, *Liza aurata*, and *Solea vulgaris* relevant to their concentration in water and sediment from the southwestern Mediterranean (coast of Sfax)", *Environmental Science and Pollution Research*, 2016

Publication

<1 %

46

Ajeng Kurniasari Putri, Giri Rohmad Barokah, Nuri Andarwulan. "Human Health Risk Assessment of Heavy Metals Bioaccumulation In Fish and Mussels from Jakarta Bay", *Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology*, 2017

Publication

<1 %

47

Mohsen Dehghani, Sana Sharifian, Mohammad Reza Taherizadeh, Moein Nabavi. "Tracing the heavy metals zinc, lead and nickel in banana shrimp (*Penaeus merguensis*) from the Persian Gulf and human health risk assessment", *Environmental Science and Pollution Research*, 2021

Publication

<1 %

48

Abdullahi Abubakar, Adamu Uzairu, Patricia Adamma Ekwumemgbo, Oluwole Joshua Okunola. " Risk Assessment of Heavy Metals in Imported Frozen Fish Species Sold

<1 %

in Nigeria: A Case Study in Zaria Metropolis", *Advances in Toxicology*, 2015

Publication

49

Haifeng Gao, Junhong Bai, Rong Xiao, Peipei Liu, Wei Jiang, Junjing Wang. "Levels, sources and risk assessment of trace elements in wetland soils of a typical shallow freshwater lake, China", *Stochastic Environmental Research and Risk Assessment*, 2012

Publication

<1 %

50

Julieta Griboff, Micha Horacek, Daniel A. Wunderlin, Magdalena V. Monferran. "Bioaccumulation and trophic transfer of metals, As and Se through a freshwater food web affected by anthropic pollution in Córdoba, Argentina", *Ecotoxicology and Environmental Safety*, 2018

Publication

<1 %

51

Mamdouh S. Masoud, Mohamed I. El-Samra, Manal M. El-Sadawy. "Heavy-metal distribution and risk assessment of sediment and fish from El-Mex Bay, Alexandria, Egypt", *Chemistry and Ecology*, 2007

Publication

<1 %

52

Arkajyoti Mukherjee, Gourav Dhar Bhowmick. "Sources of Heavy Metal Contamination in Surface Waters of Indian Freshwater Ecosystems and Their Effect on

<1 %

Aquatic Biota: A State of the Art Review",
Journal of Hazardous, Toxic, and
Radioactive Waste, 2024

Publication

53

Hang, X.. "Characteristics and accumulation of heavy metals in sediments originated from an electroplating plant", Journal of Hazardous Materials, 20090430

Publication

<1 %

54

Hülya Karadede, Erhan Ünlü.
"Concentrations of some heavy metals in water, sediment and fish species from the Atatürk Dam Lake (Euphrates), Turkey", Chemosphere, 2000

Publication

<1 %

55

Manyin Zhang, Lijuan Cui, Lianxi Sheng, Yifei Wang. "Distribution and enrichment of heavy metals among sediments, water body and plants in Hengshuihu Wetland of Northern China", Ecological Engineering, 2009

Publication

<1 %

56

Owhonda Chikeru Ihunwo, Amalo Ndu Dibofori-Orji, Clifford Olowu, Millicent Uzoamaka Ibezim-Ezeani. "Distribution and risk assessment of some heavy metals in surface water, sediment and grey mullet (Mugil cephalus) from contaminated creek in Woji, southern Nigeria", Marine Pollution Bulletin, 2020

Publication

<1 %

57

Adnorita Fandah Oktariani, Putu Eka Sudaryatma, Yan Ramona, I Made Gelgel Wirasuta et al. "Heavy metals content in fresh tuna and swordfish caught from Hindian and Pacific Oceans: Health risk assessment of dietary exposure", *Veterinary World*, 2023

Publication

<1 %

58

Ahmet Demirak, Fevzi Yilmaz, A. Levent Tuna, Nedim Ozdemir. "Heavy metals in water, sediment and tissues of *Leuciscus cephalus* from a stream in southwestern Turkey", *Chemosphere*, 2006

Publication

<1 %

59

Ali A. Al-Halani et al.. "The Seasonal Assessment of Heavy Metals Bioaccumulation in European Seabass (*Dicentrarchus labrax*) Inhabiting Damietta Fishing Harbor, Egypt", *Egyptian Journal of Aquatic Biology and Fisheries*, 2021

Publication

<1 %

60

Ayşe Bahar Yilmaz. "Heavy metals in water and in tissues of himri (*Carasobarbus luteus*) from Orontes (Asi) River, Turkey", *Environmental Monitoring and Assessment*, 09/2008

Publication

<1 %

61

Jun Yi, Linus Shing Him Lo, Hongbin Liu, Pei-Yuan Qian, Jinping Cheng. "Study of Heavy Metals and Microbial Communities in

<1 %

Contaminated Sediments Along an Urban Estuary", *Frontiers in Marine Science*, 2021

Publication

62

Naghmeh Soltani, Farid Moore, Behnam Keshavarzi, Armin Sorooshian, Reza Javid. "Potentially toxic elements (PTEs) and polycyclic aromatic hydrocarbons (PAHs) in fish and prawn in the Persian Gulf, Iran", *Ecotoxicology and Environmental Safety*, 2019

Publication

<1 %

63

Noverita Dian Takarina, Anna Ida Sunaryo Purwiyanto, Yulianto Suteja. "Cadmium (Cd), Copper (Cu), and Zinc (Zn) levels in commercial and non-commercial fishes in the Blanakan River Estuary, Indonesia: A preliminary study", *Marine Pollution Bulletin*, 2021

Publication

<1 %

64

Shabanda, Ibrahim Sani. "In Vitro Bioaccessibility for Health Risk Assessment of Metal Contamination in Urban Dust and Soil Samples: A Chemometric Approach", University of Malaya (Malaysia), 2023

Publication

<1 %

65

Zhihua Tang, Xinyu Liu, Xiaojun Niu, Hua Yin, Minru Liu, Dongqing Zhang, Huafang Guo. "Ecological risk assessment of aquatic organisms induced by heavy metals in the

<1 %

estuarine waters of the Pearl River",
Scientific Reports, 2023

Publication

66

Zita Naangmenyele, Somandla Ncube, Felix J. Akpabey, Simiso Dube, Mathew M. Nindi. "Bioaccumulation and Human Risk Assessment of Heavy Metals in Oreochromis niloticus and Clarias gariepinus Fish Species from the Golinga Reservoir, Ghana", South African Journal of Chemistry, 2021

Publication

<1 %

67

Mohamed E. Shaheen, Joel E. Gagnon, J.C. Barrette, Amr E. Keshta. "Evaluation of pollution levels in sediments from Lake Edku, Egypt using laser ablation inductively coupled plasma mass spectrometry", Marine Pollution Bulletin, 2024

Publication

<1 %

68

Nazanin Kalani, Borhan Riazi, Abdol Reza Karbassi, Faramarz Moattar. "Measurement and ecological risk assessment of heavy metals accumulated in sediment and water collected from Gomishan international wetland, Iran", Water Science and Technology, 2021

Publication

<1 %

69

O. E. Aina, L. L. Mugivhisa, J. O. Olowoyo, C. L. Obi. "Heavy metals and potential health risk assessment of Lactuca sativa and

<1 %

Daucus carota from soil treated with organic manures and chemical fertilizer", Environmental Monitoring and Assessment, 2024

Publication

70

Phanwimol Tanhan, Kanjana Imsilp, Niyada Lansubsakul, Wachiryah Thong-asa.

"Oxidative response to Cd and Pb accumulation in coastal fishes of Pattani Bay", Italian Journal of Animal Science, 2023

Publication

71

Pradip Kumar Maurya, D.S. Malik, Krishna Kumar Yadav, Amit Kumar, Sandeep Kumar, Hesam Kamyab.

"Bioaccumulation and potential sources of heavy metal contamination in fish species in River Ganga basin: Possible human health risks evaluation", Toxicology Reports, 2019

Publication

72

Sonam Tashi, Dhan Badhur Gurung, Ugyen Dorji, Karma Wangchuk.

"Trace elements in fish species: Schizothorax richardsonii, Salmo trutta and Neolissochilus hexagonolepis in relation to sizes and tissues of Punatsang Chhu", Environmental Chemistry and Ecotoxicology, 2022

Publication

73

Ulaganathan Arisekar, Rajendran Shalini, Robinson Jeya Shakila, Soundar Abuthagir Ibrahlim et al. "Selenium and mercury

<1 %

<1 %

<1 %

<1 %

concentration, Se/Hg molar ratio and risk-benefit assessment of marine fish consumption: Human health risks and protective role of Se against Hg toxicity", Food Research International, 2024

Publication

74

Ulaganathan Arisekar, Rajendran Shalini, Soundar Abuthagir Ibrahahim, S. Deepika et al. "Biomonitoring of mercury and selenium in commercially important shellfish: Distribution pattern, health benefit assessment and consumption advisories", Environmental Geochemistry and Health, 2024

Publication

<1 %

75

Zahra Shaabani, Abbas Esmaili-sari, Ali Mashinchian Moradi, Lobat Taghavi, Forough Farsad. "Possible health risk assessment for heavy metal concentrations in water, sediment, and fish species and Turkmen pregnant women's biomonitoring in Miankaleh Peninsula, Iran", Environmental Science and Pollution Research, 2022

Publication

<1 %

76

E. Sherly Williams, V. Lekshmi Priya, L. Razeena Karim. "Bioaccumulation of heavy metals in edible tissue of crab (*Scylla serrata*) from an estuarine Ramsar site in

<1 %

Kerala, South India", Watershed Ecology and the Environment, 2022

Publication

77 Josephine Omowumi Olayinka-Olagunju. "Heavy Metal Bioaccumulation and Histopathological Studies of Fish Tissues from Ose River, Ondo State, Nigeria", European Journal of Environment and Earth Sciences, 2022

Publication

78 Selda Tekin-Özan. "Determination of heavy metal levels in water, sediment and tissues of tench (*Tinca tinca* L., 1758) from Beyşehir Lake (Turkey)", Environmental Monitoring and Assessment, 10/2008

Publication

79 Thomas Dippong, Maria-Alexandra Resz. "Heavy metal contamination assessment and potential human health risk of water quality of lakes situated in the protected area of Tisa, Romania", Heliyon, 2024

Publication

80 "Biotechnological Approaches to Sustainable Development Goals", Springer Science and Business Media LLC, 2023

Publication

81 Maryam Mohammadi, Abolfazl Askary Sary, Mojgan Khodadadi. "Determination of heavy metals in two barbs, *Barbus grypus* and *Barbus xanthopterus* in Karoon and

Dez Rivers, Khoozestan, Iran", Bulletin of Environmental Contamination and Toxicology, 2011

Publication

82

Memet Varol, Muhammet Raşit Sünbül. "Multiple approaches to assess human health risks from carcinogenic and non-carcinogenic metals via consumption of five fish species from a large reservoir in Turkey", Science of The Total Environment, 2018

Publication

<1 %

83

N. Rubalingeswari, D. Thulasimala, L. Giridharan, V. Gopal, N.S. Magesh, M. Jayaprakash. "Bioaccumulation of heavy metals in water, sediment, and tissues of major fisheries from Adyar estuary, southeast coast of India: An ecotoxicological impact of a metropolitan city", Marine Pollution Bulletin, 2021

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

ASSESSING THE HEALTH OF SOUTH KALIMANTAN COASTAL SWAMP WETLANDS USING MEASUREMENTS OF HEAVY METALS IN COMMERCIAL FISH SPECIES

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15

PAGE 16

PAGE 17

PAGE 18

PAGE 19

PAGE 20
