



Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

FR-ICTCRED-007 - Article Production

11 messages

Food Research <foodresearch.my@outlook.com>
To: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

Thu, Jul 8, 2021 at 8:42 PM

Dear Dr Riyadi,

Manuscript ID: FR-ICTCRED-007

Manuscript Title: Chemical profiles and antioxidant properties of *Bruguiera gymnorrhiza* fruit extracts from Central Sulawesi, Indonesia

Before we can proceed with the article production, I would like to clarify a few points that I have commented in the manuscript. Please refer to the attachment. Please address the issues raised in the comments.

Please use the attached copy to make your revisions as it has been corrected to the Journal's format. Once you have done, kindly revert the copy to me as soon as possible. Please note the faster you respond, the quicker we will process your manuscript.

Thanks & Regards,
Vivian New
Editor
Food Research

 **FR-ICTCRED-007.docx**
955K

Putut Har Riyadi <putut.riyadi@live.undip.ac.id>
To: wendytanod@gmail.com

Fri, Jul 9, 2021 at 10:56 AM

[Quoted text hidden]

 **FR-ICTCRED-007.docx**
955K

Putut Har Riyadi <putut.riyadi@live.undip.ac.id>
To: Food Research <foodresearch.my@outlook.com>

Fri, Jul 9, 2021 at 2:54 PM

Dear Dr. Vivian New
Editor
Food Research

Manuscript ID: FR-ICTCRED-007

Manuscript Title: Chemical profiles and antioxidant properties of *Bruguiera gymnorrhiza* fruit extracts from Central Sulawesi, Indonesia

Thanks & Regards,

Dr Riyadi

[Quoted text hidden]

 **FR-ICTCRED-007.docx**
953K

Food Research <foodresearch.my@outlook.com>
To: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

Sat, Jul 10, 2021 at 2:12 PM

Dear Dr Riyadi,

Received with thanks.

Thanks & Regards,
Vivian New
Editor
Food Research

From: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>
Sent: Friday, 9 July, 2021 3:54 PM
To: Food Research <foodresearch.my@outlook.com>
Subject: Re: FR-ICTCRED-007 - Article Production

[Quoted text hidden]

Food Research <foodresearch.my@outlook.com>
To: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

Wed, Jul 14, 2021 at 8:10 PM

Dear Dr Riyadi,

Please address the issue raised in the comment.

Thanks & Regards,
Vivian New
Editor
Food Research

From: Food Research <foodresearch.my@outlook.com>
Sent: Saturday, 10 July, 2021 3:12 PM
To: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

[Quoted text hidden]

[Quoted text hidden]

 **FR-ICTCRED-007.docx**
955K

Putut Har Riyadi <putut.riyadi@live.undip.ac.id>
To: Food Research <foodresearch.my@outlook.com>

Fri, Jul 16, 2021 at 3:57 PM

Dear Dr. Vivian New
Editor
Food Research

I have corrected the manuscript. Thank you very much.

Best regards

Dr. Riyadi

[Quoted text hidden]

 **FR-ICTCRED-007 (fix).docx**
949K

Food Research <foodresearch.my@outlook.com>
To: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

Sat, Jul 17, 2021 at 10:04 AM

Dear Dr Riyadi,

Please refer to the attachment for the galley proof of your manuscript FR-ICTCRED-007 entitled 'Bioactive compound profiles and antioxidant properties of *Bruguiera gymnorrhiza* fruits extract from Central Sulawesi, Indonesia'. Please check the content of the galley proof. If there are any mistakes, please comment and highlight in the PDF itself and revert to us within two (2) days of receipt. Once we have finalized the PDF version, your manuscript will be published online for early viewing.

Thanks & Regards,
Vivian New
Editor
Food Research

From: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

Sent: Friday, 16 July, 2021 4:57 PM

[Quoted text hidden]

[Quoted text hidden]

 **FR-ICTCRED-007.pdf**
826K

Putut Har Riyadi <putut.riyadi@live.undip.ac.id>
To: Food Research <foodresearch.my@outlook.com>

Sat, Jul 17, 2021 at 4:56 PM

Dear Dr. Vivian New
Editor
Food Research

One writing error, and we have commented and highlighted it in the PDF.
Thank you very much.

Best regards

Dr. Riyadi

[Quoted text hidden]

 **FR-ICTCRED-007.pdf**
800K

Food Research <foodresearch.my@outlook.com>
To: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

Sat, Jul 17, 2021 at 7:37 PM

Dear Dr Riyadi,

Please refer to the attachment for the edited galley proof. If the galley proof is fine, please approve the galley proof.

Thanks & Regards,
Vivian New
Editor
Food Research

From: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

Sent: Saturday, 17 July, 2021 5:56 PM

[Quoted text hidden]

[Quoted text hidden]

 **FR-ICTCRED-007.pdf**
826K

Putut Har Riyadi <putut.riyadi@live.undip.ac.id>
To: Food Research <foodresearch.my@outlook.com>

Sun, Jul 18, 2021 at 10:53 AM

Dear Dr. Vivian New
Editor
Food Research

Approved, please proceed.

Thanks & Regards,
Dr. Riyadi

[Quoted text hidden]

Food Research <foodresearch.my@outlook.com>
To: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

Sun, Jul 18, 2021 at 2:23 PM

Dear Dr Riyadi,

Noted with thanks.

Thanks & Regards,
Vivian New
Editor
Food Research

From: Putut Har Riyadi <putut.riyadi@live.undip.ac.id>

Sent: Sunday, 18 July, 2021 11:53 AM

[Quoted text hidden]

[Quoted text hidden]

Chemical profiles and antioxidant properties of *Bruguiera gymnorrhiza* fruit extracts from Central Sulawesi, Indonesia

^{1,*}Riyadi, P.H., ²Tanod, W.A., ³Dewanto, D.K., ⁴Herawati, V.E., ¹Susanto, E. and ⁵Aisiah, S.

¹Department of Fish Product Technology, Faculty of Fisheries and Marine Science, Universitas Diponegoro, Semarang, Central Java 50275 Indonesia

²Department of Fisheries and Marine, Politeknik Negeri Nusa Utara, Tahuna, North Sulawesi 95812 Indonesia

³Department of Fishing Technology, Sekolah Tinggi Perikanan dan Kelautan, Palu, Central Sulawesi 94118 Indonesia

⁴Department of Aquaculture, Faculty of Fisheries and Marine Science, Universitas Diponegoro, Semarang, Central Java 50275 Indonesia

⁵Study Program of Aquaculture, Faculty of Fisheries and Marine, Universitas Lambung Mangkurat, Banjarbaru, South Kalimantan 70714, Indonesia

Article history:

Received: 9 February 2021

Received in revised form: 21 March 2021

Accepted: 24 March 2021

Available Online: 28 July 2021

Keywords:

ADMET,

Antioxidant,

Bruguiera gymnorrhiza,

GCMS,

Nutraceutical

DOI:

[https://doi.org/10.26656/fr.2017.5\(S3\).007](https://doi.org/10.26656/fr.2017.5(S3).007)

Abstract

Bruguiera gymnorrhiza is one of the mangrove species that a source of antioxidants. Antioxidant substances are able to protect cells from oxidative stress and other related diseases. This study aimed to evaluate the chemical profiles, predicted biological activity, and antioxidant activity of the *B. gymnorrhiza* fruit extracts. The research methods included sampling, extraction (maceration with MeOH:DCM), identification of chemical profiles (GCMS spectra analysis), assaying for computational analysis (PASS server and ADMET), antioxidants (DPPH radical scavenging), and total phenolic content (Folin-Ciocalteu). *Bruguiera gymnorrhiza* fruits was collected from Central Sulawesi, Indonesia. The chemical profiles detected in the *B. gymnorrhiza* fruit extracts, namely isopimaradiene (64.20%); 4-(2-Aminopropyl) phenol (19.06%); dimethylamino-dimethylphosphene oxide (9.40%); 3-amino-2-benzylbutanoic acid (5.46%); and 1,4-dideuteriooctane (1.89%). PASS server analysis showed that the five compounds detected from *B. gymnorrhiza* fruit have the potential as an NF-E2-related factor (Nrf2) stimulant and oxygen scavenger. The ADMET analysis results indicated that *B. gymnorrhiza* fruits could be developed as folk medicine and nutraceutical products.

1. Introduction

Mangroves are a forested ecosystem widely distributed in tropical and subtropical areas (Zhu *et al.*, 2012; Al-Maqtari and Nagi, 2014; Suh *et al.*, 2014). Indonesia is recognized as a centre for biodiversity globally as an archipelago country due to its rich natural habitats, including mangroves (Kusmana, 2014). About 23% of total world mangrove forests are located in Indonesia (Armitage, 2020), with 202 mangroves species, e.g., *Avicennia* sp., *Bruguiera* sp., *Rhizophora* sp., and *Xylocarpus* sp. (Noor *et al.*, 2006; Kusmana, 2014). They have been utilized for several purposes, such as wood for house constructions, charcoal (Dahdouh-Guebas *et al.*, 2000), food ingredients (Jariyah *et al.*, 2014; Analuddin *et al.*, 2019; Amin *et al.*, 2021),

complementary foods (Kardiman *et al.*, 2017), food products (Situmorang and Barus, 2015; Wibawanti *et al.*, 2018), and folk medicine (Rout and Basak, 2014; Suh *et al.*, 2014).

Mangroves have long been used as folk medicine and therapeutics for diarrhoea, indigestion, and inflammation in coastal communities (Rout and Basak, 2014; Suh *et al.*, 2014). Their efficacies are related to the level and composition of bioactive compounds. It is expected to have several physiological activities such as anti-fungal (Acharya *et al.*, 2020), anti-bacteria (Behbahani *et al.*, 2018; Eswaraiah *et al.*, 2020), antioxidants (Arulkumar *et al.*, 2020; Karim *et al.*, 2020), and anti-hemolytic (Karim *et al.*, 2020).

*Corresponding author.

Email: putut.riyadi@live.undip.ac.id

The coast of Central Sulawesi, Indonesia, is the home of various mangrove species such as *Rhizophora mucronata*, *Sonneratia alba*, *Avicennia marina*, and *Bruguiera gymnorrhiza* (Damanik and Djamaludin, 2012; Lisna et al., 2017; Dewanto et al., 2018; Alhaffaf et al., 2019). *B. gymnorrhiza*, belonging to Rhizophoraceae, is reported as an essential and major mangrove species in the Asia Pacific, including Central Sulawesi (Zhu et al., 2012; Kusmana, 2014). It is traditionally used as a medicinal plant (Lin et al., 2020). The previous study has revealed that methanol leaves extract of *B. gymnorrhiza* and *Heritiera littoralis* revealed free radical scavenging, anti-haemolytic, cytotoxic, and antibacterial activity (Karim et al., 2020). Meanwhile, the aqueous extract of *B. gymnorrhiza* exhibited a protective effect against dextran sulfate sodium (DSS)-induced ulcerative colitis (Chen et al., 2020). In addition, the *B. gymnorrhiza* roots extract has been reported to inhibit the *Escherichia coli*, *Staphylococcus aureus*, and urinary tract infections (UTI) bacterial pathogens, e.g., *Pseudomonas aeruginosa* growth (Acharya et al., 2020). However, explorations of *B. gymnorrhiza* fruits have been limited. Even though *B. gymnorrhiza* fruits have reported possessed biological activities, it is therefore related to the chemical profiles and their antioxidant activities.

Therefore, in this study, we reported *B. gymnorrhiza* fruit extract's chemical profiles with GCMS, the prediction of biological activity with PASS server and ADMET analysis, and antioxidant activity with DPPH radical scavenging and total phenol content with Folin-Ciocalteu.

2. Materials and methods

2.1 Sampling location

Bruguiera gymnorrhiza fruits were collected from the coastal area of Tomini Bay in Laemanta, Parigi Moutong, Central Sulawesi, Indonesia (-0.1847 S and 120.0088 E) (Figure 1) in April 2020. Mangrove fruits were randomly taken from a mangrove tree *B. gymnorrhiza* followed by identification based on Noor et al. (2006). After collection, the *B. gymnorrhiza* fruits were washed with water to eliminate the presence of contaminants.

2.2 Materials

Free radical 2,2-Diphenyl-1-Picrylhydrazyl (DPPH, Merck-1898664), distilled water, dichloromethane (DCM, CH₂Cl₂; (Merck- 106050), methanol (MeOH, CH₃OH; Merck- 107018), Folin-Ciocalteu reagent (Merck-109001), gallic acid (C₆H₂(OH)₃CO₂H, Merck-842649; purity ≥ 98%), Na₂CO₃ were purchased from CV Amani Media Malang and CV Intraco Makassar,

Indonesia.




Figure 1. Sample coordinates for *Bruguiera gymnorrhiza* from Laemanta, Central Sulawesi, Indonesia

2.3 Preparation of *Bruguiera gymnorrhiza* extracts

Before extractions, 500 g of *B. gymnorrhiza* fruits were dried at 50-60°C using an oven (Finco OV50) for 15-20 hrs. After drying, the samples were pounded into flour. The crude extract of *B. gymnorrhiza* (100 g) was obtained by extraction with the combination of MeOH and DCM (1: 1) for 48 hrs (Hsiao et al., 2015; Putra et al., 2016). The filtrate obtained by the extraction was filtered with filter paper (Whatman No. 42). The filtrate was vacuum-concentrated at 40-45°C using a rotary evaporator (EYELA N-1100) to obtain the crude extracts. The extraction process was carried out three times. Then, the extracts were weighed and dissolved in 30% MeOH then stored at 4-5°C for further analysis.

2.4 Chemical profiles screening using GCMS

The chemical components of *B. gymnorrhiza* fruits extract were profiled using Gas Chromatography-Mass Spectrometry (GCMS). The analysis was performed on an HP 6890 GCMS system (Hewlett-Packard, California, USA) equipped with a capillary column (Agilent 19091S -433 HP-5MS; 30 m x 250 µm i.d.; Santa-Clara, California, USA). The carrier gas was helium at a flow rate of 1 mL/min. The oven temperature was set at 325°C. The pre-oven temperature was 150°C held at 2°C/min. It ran for 10°C/min and was then increased to 240°C, holding time for 11 mins. The total running time was 24 mins. The scanning range was 50 - 550 amu. Structural assignments were based on analysis of mass spectral fragmentation patterns and compared with mass spectra in the National Institute of Standards and Technology (NIST) and Wiley's compound profile database (Tanod et al., 2019).

2.5 Prediction of biological activity

The chemical profiles from *B. gymnorrhiza* fruits extract by GCMS were predicted through Prediction of

Activity Spectra for Substances (PASS)-way2Drug server (<http://www.pharmaexpert.ru/passonline/index.php>). PASS server is used to predict the chemical compounds' biological activity (Aisiah *et al.*, 2020; Riyadi, Romadhon, Anggo *et al.*, 2020). The structural target compounds obtained from the National Centre for Biotechnology Information (<https://pubchem.ncbi.nlm.nih.gov/>) in the form of canonical SMILE were used to predict their biological activity. The chemicals profile structure was also used to estimate their ADMET (absorption, distribution, metabolism, excretion, and toxicity) through AdmetSAR (<http://lmmd.ecust.edu.cn/admetar2>).

2.6 Determination of total phenol contents

Total phenolic content (TPC) of *B. gymnorrhiza* fruits extract was determined by the Folin-Ciocalteu (FC) reagent method (Blainski *et al.*, 2013 and Lamuela-Raventós, 2017). The gallic acids (1 mg/mL) were used as a standard by diluting ethanol: distilled water (1: 1). Gallic acid solutions were diluted in serial dilution with concentrations 5, 20, 40, 60, 80, and 100 µg/mL. From each dilution concentration, 1 mL of gallic acid was taken, and 9 mL of distilled water was added. Then those aliquots were mixed with 1 mL of FC reagent (homogenization) and incubated for 8 mins at room temperature. After incubation, 3 mL of 20% Na₂CO₃ was added, and the mixture was allowed to stand at room temperature for 2 hrs. The absorbance was measured at 750 nm. The standard curve for gallic acid was prepared with gallic acid concentration (µg/mL) with an absorbance value.

The aliquot (1 mg/mL extract) was added with 10 mL of distilled water + 1 mL of FC reagent and incubated as described above. Then TPC content was measured at 750 nm. *B. gymnorrhiza* fruits extract's total phenolic content was then calculated based on a gallic acid standard curve and expressed as mg of gallic acids equivalent (GAE)/g extract.

2.7 DPPH radical scavenging activity

DPPH is a free radical, stable at room temperature, which produces a violet solution in ethanol. Antioxidants' presence will reduce the colour and change into uncoloured ethanol solution (Clarke *et al.*, 2013). The total free radical scavenging capacity of Free radical scavenging capacity was assessed according to the previously reported method using a stable DPPH (Molyneux, 2004; Dewanto *et al.*, 2019). The extract was expressed as the IC₅₀, which is the antioxidant concentration required to quench 50% of the initial DPPH under the experiment treatment. Vitamin E, known as the strong DPPH radical scavenger (Yu *et al.*,

2002), was used as the positive control. The IC₅₀ value was converted to the vitamin E equivalent. Each evaluation was performed in triplicate.

Bruguiera gymnorrhiza fruit extracts (200 µg/mL) in ethanol solution was prepared and diluted into several concentrations (20, 40, 60, 80, 100 µg/mL). 2 mL aliquot of each concentration was reacted with 50 µM DPPH ethanol solution. The mixtures were homogenized and stand for 30 mins in the dark at room temperature before reading against a blank at 517 nm (UV-VIS spectrophotometer T90 + PG Instruments Ltd).

The DPPH radical scavenging effect in percentage (%) was calculated according to the following equation:

$$\text{DPPH Scavenging Effect (\%)} = \frac{A_{\text{blank}} - A_{\text{sample}}}{A_{\text{blank}}} \times 100\%$$

Where A_{sample} is the absorbance of a sample solution and A_{blank} is the absorbance of the blank solution (containing all reagents except the test sample).

The half of maximum concentration in reducing initial DPPH was calculated from the graph by plotting inhibition concentrations against initial DPPH to obtain a linear regression equation, $y = a + bx$. Each measurement was carried out in triplicates.

3. Results

3.1 Sample identification

Identification was carried out according to Noor *et al.* (2006). The samples included in the *B. gymnorrhiza* based on the leaves' shape and tip, the types of roots, fruit, and flowers. BG's bark has lenticels, and the surface is smooth to rough, dark gray to brown. BG roots are like planks extending to the side at the tree's base and developing several knee roots. Characteristics of BG leaves are skinned green on top and yellow on bottom. The BG leaves shape is elliptical to elliptical-lanceolate and had average length and width 4.5 to 7 cm and 8.5 to 22 cm, respectively. Characteristics of BG fruit are straight, blunt, and dark green-purple when fresh with length (12-30 cm) and diameter (1.5-2 cm) (Figure 2).




Figure 2. *Bruguiera gymnorrhiza* mangrove fruits from Laemanta, Central Sulawesi, Indonesia

3.2 Chemical profiles of *Bruguiera gymnorrhiza* fruit extracts

The GCMS analysis showed the presence of 5 compounds from *B. gymnorrhiza* fruit extracts (Table 1). The chemical profiles were identified through mass spectrometry attached with GC. The total ion chromatogram (TIC) of the GCMS analysis confirms that five compounds and their different retention time (4,700 to 22,824 mins) are presented in Figure 3. The chemical profiles contained in *B. gymnorrhiza* fruits extracts are isopimaradiene (64.20%), 4-(2-Aminopropyl) phenol (19.06%), dimethylamino-dimethylphosphene oxide (9.40%), 3-amino-2-benzylbutanoic acid (5.40), and 1,4-dideuteriooctane (1.89 %).

Abundance



Time->

Figure 3. GCMS chromatogram of mangrove fruit extracts *Bruguiera gymnorrhiza*

3.3 Biological activity prediction of fruits extracts

The prediction of biological activity was performed with chemical profiles in *B. gymnorrhiza* fruits extracts and presented in Table 2. The biological activity prediction was used the PASS server. The PASS server could predict compounds' biological activity based on their formula with 95 % accuracy (Filimonov and Poroikov, 2008; Riyadi, Tanod, Wahyudi *et al.*, 2020).

The biological activity prediction value of the *B. gymnorrhiza* fruits extracts' chemical profiles was expressed as a "probability to be active" (Pa). The chemical profiles are probable to act as antioxidants, free radical scavengers, oxygen scavengers, and NF-E2-related factor (Nrf2) stimulant properties (Figure 4). The PASS server was analysed and dominated by Nrf2 stimulant and oxygen scavenger, which play a role in protecting oxidative stress.




Figure 4. Probability to be active from *Bruguiera gymnorrhiza* fruit extracts profiles with PASS Server

ADMET analysis is an important parameter to determine whether a natural product will reach the body's target when administered orally (Dong *et al.*, 2018; Natesh *et al.*, 2021). The ADMET properties of compounds are important to the development process of nutraceutical and pharmaceutical products (van de Waterbeemd and Gifford, 2003; Guan *et al.*, 2019). The utilization of *B. gymnorrhiza* fruits extracts for human health benefits needs to be satisfied with the ADMET properties before further applications. Therefore, we evaluated the ADMET properties of *B. gymnorrhiza* fruits extracts chemical profiles by using admetSAR (Table 2).

The chemical profiles of *B. gymnorrhiza* fruit extracts showed good human intestinal absorption. However, only dimethylamino-dimethyl phosphene oxide did not show well on the caco-2 permeability (Table 2). Interestingly, none of the selected chemical profile functions is acted as both P-glycoprotein (P-gp) substrate and inhibitor. Furthermore, 4-(2-Aminopropyl) phenol), isopimaradiene, and 3-amino-2-benzyl butanoic acid from the *B. gymnorrhiza* fruits extracts act as substrate of CYP450 2D6, CYP450 3A4, and CYP450 2C9, respectively. In contrast, none of the chemical profiles acts as inhibitors of CYP450 enzymes, which

Table 1. Chemical profiles of *Bruguiera gymnorrhiza* fruit extracts by GCMS

RT (Min)	Chemical Profiles	Formula	Molecular Weight (g/mol)	Area (%)	Pubchem CID
4.700	1,4-dideuteriooctane	C ₈ H ₁₈	116.24	1.89	151945284
9.158	4-(2-Aminopropyl)phenol	C ₉ H ₁₃ NO	151.21	15.93	3651
11.295				3.13	
11.940	Isopimaradiene	C ₂₀ H ₃₂	272.5	64.2	13969536
12.453	dimethylamino-dimethylphosphene oxide	C ₄ H ₁₂ NOP	121.12	9.4	521290
22.824	3-amino-2-benzylbutanoic acid	C ₁₁ H ₁₅ NO ₂	193.24	5.46	541835

Table 2. Admet profile of *Bruguiera gymnorrhiza* fruit extracts

ADMET Profile	1,4-dideuterio Octane	4-(2-Aminopropyl) phenol)	Isopimara-diene	dimethylamino-pdimethyl phosphene oxide	3-amino-2-benzyl butanoic acid
Absorption properties					
Human Oral Bioavailability	+ (0.5286)	- (0.5857)	+ (0.6571)	+ (0.6000)	+ (0.8286)
Human Intestinal Absorption	+ (0.8865)	+ (0.9861)	+ (0.9828)	+ (0.8805)	+ (0.9945)
Caco-2 Permeability	+(0.9825)	+(0.9499)	+(0.8457)	-(0.5244)	+ (0.5957)
P-glycoprotein Substrate	-(0.9516)	-(0.8011)	-(0.9000)	-(0.98)	- (0.9485)
P-glycoprotein Inhibitor	-(0.9850)	-(0.9903)	-(0.8387)	-(0.988)	- (0.9903)
Distribution properties					
Blood-Brain Barrier (BBB) B	+(1.000)	-(0.993)		+(0.9911)	+ (0.8686)
Subcellular localization					
Metabolism properties					
CYP450 2C9 Substrate	-(0.8415)	-(1.0000)	-(0.7731)	-(1.0000)	+ (0.6409)
CYP450 2D6 Substrate	-(0.7215)	+(0.5870)	-(0.7252)	-(0.7913)	-(0.7568)
CYP450 3A4 Substrate	-(0.7947)	-(0.7593)	+(0.5636)	-(0.6488)	-(0.7595)
CYP450 2C9 Inhibitor	-(0.9349)	-(0.8930)	-(0.5747)	-(0.8930)	-(0.967)
CYP450 2D6 Inhibitor	-(0.9373)	-(0.7163)	-(0.9203)	-(0.9491)	-(0.944)
CYP450 2C19 Inhibitor	-(0.9540)	-(0.7723)	-(0.500)	-(0.8930)	-(0.8805)
CYP450 3A4 Inhibitor	-(0.9877)	-(0.8767)	-(0.8545)	-(0.9724)	-(0.9463)
CYP inhibitory promiscuity	-(0.8149)	-(0.9737)	-(0.6444)	-(0.9737)	-(0.9818)
Toxicity properties					
Hepatotoxicity	- (0.9500)	- (0.7750)	- (0.7500)	- (0.9500)	+ (0.825)
Acute Oral Toxicity		II (0.5359)	II (0.8150)	III (0.6949)	III (0.7484)
AMES (Ames mutagenesis)		-(0.9300)	-(0.9400)	-(0.5800)	

can interfere degradation process. In addition, all chemical profiles showed a positive response as oral bioavailability, water-soluble, and could bind to proteins in the blood.

3.4 Total phenol content

Phenolic compounds are mainly found in plants. Phenolic compounds have a protective role in human health and are beneficial in food and pharmaceutical products due to their benefits (Salar *et al.*, 2017). The present study evaluated the quantitative estimation of phenolic compounds in BGF extract collected from North Sulawesi, Indonesia. The TPC value of BGF extract was determined by the FC method reported by Blainski *et al.* (2013) and Lamuela-Raventós (2017). Gallic acid was used as a standard for the calibration curve to calculate the TPC with equation $y = 0.0101x - 0.0318$. The calculated result of TPC was accounted for 27539.60 ± 35.01 mg GAE/100 g dry extract. The results indicated that *B. gymnorrhiza* fruit extracts contained a high level of phenolic compounds.

3.5 Antioxidant activity analysis using the DPPH method

The DPPH method has been widely used for antioxidant evaluation due to a stable free radical to evaluate radical scavenging activity (Liu *et al.*, 2004). The level of discoloration exhibits the potency of

antioxidants to scavenge free radicals. These phenomena due to the reaction between the 2,2-Diphenyl-1-Picrylhydrazyl and antioxidant compound to form the hydrazine diphenyl picrite (Huliselan *et al.*, 2015; Tanod *et al.*, 2019). DPPH could reduce the exposure of radical proton scavengers (Yamaguchi *et al.*, 1998). The antioxidants reduced the stable DPPH radicals, resulting in decreased absorbance at 517 nm (Behgar *et al.*, 2011). The present study evaluated *B. gymnorrhiza* fruit extracts' antioxidant activity using the DPPH radical scavenging method with more than one concentration and compared the results with vitamin E (Figure 5).




Figure 5. DPPH scavenging effect from *Bruguiera gymnorrhiza* mangrove fruit extracts

The *B. gymnorrhiza* fruit extracts were also

evaluated for the inhibitory effect of DPPH radicals for IC₅₀ determination (Table 3). The *B. gymnorrhiza* fruit extracts possessed antioxidant activity because they could donate hydrogen atoms/electrons to react with DPPH radicals. The results showed a *B. gymnorrhiza* fruit extracts concentration dependent on free radical scavenging and comparable with synthetic antioxidants (vitamin E). The results indicated that *B. gymnorrhiza* fruit extracts have strong antioxidant activity (IC₅₀: 70.45±0.49 µg/mL), however its activity lower than vitamin E (IC₅₀: 13.72±1.02 µg/mL).

Table 3. IC₅₀ of *Bruguiera gymnorrhiza* mangrove fruit extracts using the DPPH Radical Scavenging, compared with vitamin E.

Mangrove fruits extract	IC ₅₀ (µg/mL)
<i>Bruguiera gymnorrhiza</i>	70.45±0.49
Vitamin E	13.72±1.02

4. Discussion

Bruguiera gymnorrhiza, a medicinal food plant, is widely distributed in Indonesia (Kusmana, 2014). *Bruguiera gymnorrhiza* has been used for several purposes, such as vegetables, firewoods (Kusmana et al., 2018), and folk medicine (Ahmed et al., 2007). The present study reported the characteristics of chemical profiles and antioxidants properties of *B. gymnorrhiza* fruit extracts. This study also predicted the biological activity related to antioxidant and ADMET properties of chemical profiles in *B. gymnorrhiza* fruit extracts. The major chemical profiles identified by GCMS from the *B. gymnorrhiza* fruit extracts were dominated by Isopimaradiene (terpenoids component) and 4-(2-Aminopropyl) phenol (phenolics component) (Figure 3 and Table 1). Plant phenolics denote a major group of compounds with antioxidants properties (Adesegun et al., 2009; Pandey and Rizvi, 2009; Lin et al., 2016). Phenolics reported have the ability to strong reactive oxygen species (ROS)-scavenging activity, dropping nitrite levels, and downregulated cyclooxygenase-2 (COX-2) expression, which can play a vital role in human health (Lin et al., 2009; Rodríguez-López et al., 2020). In addition, terpenoids are also reported as antioxidant potential (Mohandas and Kumaraswamy, 2018). Terpenoids have a hydroxyl group that may act as an antioxidant due to their relatively complex cyclic structure (consisting of alcohol, aldehyde, or carboxylic acid) (Dewanto et al., 2019).

The predictive of *B. gymnorrhiza* fruit extract chemical profiles by PASS-Server of the major compounds are related to antioxidant properties. Isopimaradiene, a diterpenoid, has been reported to have anti-oxidative and anti-inflammatory properties (Kim

and Kim, 2009; Tungharoen et al., 2019) and anti-bacterial activity (Rijo et al., 2009). Meanwhile, 4-(2-Aminopropyl) phenol is exerted antioxidant activities, free radical scavenger, O₂ scavenger, and NF-E2-related factor 2 (Nrf2) stimulant. Nuclear factor erythroid 2-related factor 2 (Nrf2) plays an important role in regulating the antioxidant proteins to protect against oxidative damages (Hwang and Jeong, 2010; Barajas et al., 2011; Ma, 2013; Cui et al., 2016). Under oxidative stress conditions, activated Nrf2 translocate into the nucleus and modulates antioxidant-related genes' expression at the transcriptional level (Johnson et al., 2008). Oxygen scavengers could prevent oxidation and quality foodstuff changes (Byun et al., 2011; Lalpuria et al., 2012; Mu et al., 2013). This compound has one phenol ring and one NH₂ bond. H atoms in the phenol and NH₂ ring act as electron donors to react as free radicalizers such as reactive oxygen species (ROS) (Lobo et al., 2010).

The present study has been found that *B. gymnorrhiza* fruit extracts possessed higher antioxidant potential and phytochemical content. The antioxidant potential is comparable with synthetic antioxidant vitamin E. The results revealed that *B. gymnorrhiza* fruit extracts exhibited strong antioxidant activity (IC₅₀ 70.45). This activity is higher than methanol extract of *B. gymnorrhiza* fruit extracts collected from the Odisha coast, India (Rout et al., 2015) and *B. gymnorrhiza* extract collected from Bangladesh (Hosen et al., 2020). Some factors are related to the different antioxidant activity of BG, e.g., the polarity of solvents (Jacob et al., 2013), the maturity of fruits (Sudirman et al., 2014), geographical locations (Rout et al., 2015; Hosen et al., 2020).

Our data showed a high level of phenolic content. This level is considerably higher than ethanol: methanol (1: 1) of *B. gymnorrhiza* fruit extracts (21.9 mg GAE/g powder) (Hosen et al., 2020). The extract contains 4-(2-Aminopropyl) phenol and -amino-2-beunzylbutanoic acid with an aromatic ring structure with a hydroxyl group (typical of phenolic compounds).

None of the selected chemical profiles functions as P-glycoprotein (P-gp) substrate and inhibitors. P-gp transporter acts as a protective barrier to keep toxins out of the body and as a cellular defence system to protect the accumulation of foreign materials in the human body (Marzolini et al., 2004; Szakács et al., 2008; Quan et al., 2012; Cloos et al., 2018). After absorption in the intestine, compounds flow to the blood then metabolized in the liver. The liver CYP450 enzyme family could break down the compounds and eliminate them from the human body (Natesh et al., 2021). According to the

ADMET analysis, 4-(2-Aminopropyl) phenol, isopimaradiene, and 3-amino-2-benzyl butanoic acid from the *B. gymnorrhiza* fruit extracts act as a substrate of CYP450 2D6, CYP450 3A4, and CYP450 2C9, respectively. Possibly, they will metabolize through those enzymes in the liver. Interestingly, all compounds are not showing any toxicity in the liver. Compound No. 1 – 3 showed class II toxicity, while compounds No. 4-5 exhibited class 3 toxicity. According to the GHS United Nations 2015, the toxicity of any compounds taken orally was classified into six categories based on the LD₅₀ ranges. Class 1- fatal (LD₅₀ ≤5), class 2- fatal (5 <LD₅₀ ≤50), class 3- toxic (50 <LD₅₀ ≤300), class 4- harmful (300 <LD₅₀ ≤2000), class 5- may be harmful (2000 <LD₅₀ ≤5000), and class 6, LD₅₀ >5000 is considered as non-toxic (Drwal et al., 2014).

5. Conclusion

It can be concluded that *B. gymnorrhiza* fruit extracts from Central Sulawesi could be developed as a folk medicine, nutraceutical, and functional food ingredient with antioxidant properties in a certain amount that is safe for consumption.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

The authors would like to thank the 6th International Conference committee on Tropical and Coastal Region Eco-Development (ICTCRED) 2020, who helped in the publication process. The authors would also like to thank Siti Khalimatu Sa'diah, S.Pi., Fatma, S.Pi., and Moh. Ahdia, S.Pi., which assists in sampling and extraction of mangrove fruit.

References

- Adesegun, S.A., Fajan, A., Orabueze, C.L. and Coker, H.A.B. (2009). Evaluation of antioxidant properties of *Phaulopsis fascisepala* C.B.Cl. (Acanthaceae). *Advanced Access Publication*, 6(2), 227-231. <https://doi.org/10.1093/ecam/nem098>
- Acharya, S., Patra, D.K., Pradhan, C. and Mohapatra, P.K. (2020). Anti-bacterial, anti-fungal and anti-oxidative properties of different extracts of *Bruguiera gymnorrhiza* L. (Mangrove). *European Journal of Integrative Medicine*, 36, 101140. <https://doi.org/10.1016/j.eujim.2020.101140>
- Aisiah, S., Olga, Tanod, W.A., Salosso, Y., Bambang and Riyadi, P.H. (2020). Computational analysis of ethyl acetate extract of *Nauclea subdita* (Korth.) Steud. leaves as peptidoglycan glycosyltransferase inhibitor in *Aeromonas hydrophila*. *IOP Conference Series: Earth and Environmental Science*, 584, 012022. <https://doi.org/10.1088/1755-1315/584/1/012022>
- Al-Maqtari, M.A. and Nagi, H.M. (2014). Screening of salt-stress, antioxidant enzyme, and antimicrobial activity of leave extracts of mangroves *Avicennia marina* L. from Hodaidah, Yemen. *Journal of Stress Physiology and Biochemistry*, 10(2), 190–199.
- Amin, M.N.G., Pebruwanto, D., Praleba, S.A., Hasan, M.N., Zakariya, Subekti, S., Pramono, H. and Alamsjah, M.A. (2021). Functional and rheological properties of mixed flour from mangrove fruit of *Bruguiera gymnorrhiza* flour and wheat flour. *Food Research*, 5(1), 167–173. [https://doi.org/10.26656/fr.2017.5\(1\).356](https://doi.org/10.26656/fr.2017.5(1).356)
- Analuddin, K., Septiana, A., Nasaruddin, Sabilu, Y. and Sharma, S. (2019). Mangrove fruit bioprospecting: nutritional and antioxidant potential as a food source for coastal communities in the Rawa Aopa Watumohai National Park, Southeast Sulawesi, Indonesia. *International Journal of Fruit Science*, 19 (4), 423–436. <https://doi.org/10.1080/15538362.2018.1555507>
- Arulkumar, A., Kumar, K.S. and Paramasivam, S. (2020). Antibacterial and invitro antioxidant potential of Indian mangroves. *Biocatalysis and Agricultural Biotechnology*, 23, 101491. <https://doi.org/10.1016/j.bcab.2019.101491>
- Armitage, D. (2020). Socio-institutional dynamics and the political ecology of mangrove forest conservation in Central Sulawesi, Indonesia. *Global Environmental Change*, 12(3), 203-217. [https://doi.org/10.1016/S0959-3780\(02\)00023-7](https://doi.org/10.1016/S0959-3780(02)00023-7)
- Barajas, B., Che, N., Yin, F., Rowshanrad, A., Orozco, L.D., Gong, K.W., Wang, X., Castellani, L.W., Reue, K., Lusia, A.J. and Araujo, J.A. (2011). NF-E2 – related factor 2 promotes atherosclerosis by effects on plasma lipoproteins and cholesterol transport that overshadow antioxidant protection. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 31(1), 58–66. <https://doi.org/10.1161/ATVBAHA.110.210906>
- Behgar, M., Ghasemi, S., Naserian, A., Borzoie, A. and Fatollahi, H. (2011). Gamma radiation effects on phenolics, antioxidants activity and in vitro digestion of pistachio (*Pistachia Vera*) Hull. *Radiation Physics and Chemistry*, 80(9), 963–967. <https://doi.org/10.1016/j.radphyschem.2011.04.016>
- Behbahani, E.A., Yazdi, F.T., Shahidi, F., Noorbakhsh, H., Vasiee, A. and Alghooneh, A. (2018). Phytochemical analysis and antibacterial activities extracts of mangrove leaf against the growth of some

- pathogenic bacteria. *Microbial Pathogenesis*, 114, 225-232. <https://doi.org/10.1016/j.micpath.2017.12.004>
- Blainski, A., Lopes, G.C. and De Mello, J.C.P. (2013). Application and analysis of the folin-ciocalteu method for the determination of the total phenolic content from *Limonium brasiliense* L. *Molecules*, 18 (6), 6852–6865. <https://doi.org/10.3390/molecules18066852>
- Byun, Y., Darby, D., Cooksey, K., Dawson, P. and Whiteside, S. (2011). Development of oxygen scavenging system containing a natural free radical scavenger and a transition metal. *Food Chemistry*, 124(2), 615-619. <https://doi.org/10.1016/j.foodchem.2010.06.084>
- Chen, I.F., Luo, D.D., Lin, Y.S., Liu, Y.H., Wu, J.Z., Yi, X.Q., Wu, Y., Zhang, Q., Gao, C.J., Cai, J. and Su, Z.R. (2020). Aqueous extract of *Bruguiera gymnorhiza* leaves protects against dextran sulfate sodium induced ulcerative colitis in mice via suppressing NF- κ B activation and modulating intestinal microbiota. *Journal of Ethnopharmacology*, 251, 112554. <https://doi.org/10.1016/j.jep.2020.112554>
- Clarke, G., Ting, K.N., Wiart, C. and Fry, J. (2013). High correlation of 145 2,2-Diphenyl-1-Picrylhydrazyl (DPPH) radical scavenging, ferric reducing activity potential and total phenolics content indicates redundancy in use of all three assays to screen for antioxidant activity of extracts of plants from the Malaysian rainforest. *Antioxidants*, 2 (1), 1–10. <https://doi.org/10.3390/antiox2010001>
- Cloos, J., Jansen, G., Jaramillo, A.C., Saig, F.A. and Peters, G.J. (2018). How to overcome ATP-binding cassette drug efflux transporter-mediated drug resistance?. *Cancer Drug Resistance*, 1(1), 6–29. <https://doi.org/10.20517/cdr.2018.02>
- Cui, T., Lai, Y., Janicki, J.S. and Wang, X. (2016). Nuclear factor erythroid-2 related factor 2 (Nrf2)-mediated protein quality control in cardiomyocytes. *Frontiers in Bioscience*, 21, 192–202. <https://doi.org/10.2741/4384>
- Dahdouh-Guebas, F., Mathenge, C., Kairo, J.G. and Koedam, N. (2000). Utilization of mangrove wood products around mida creek (Kenya) amongst subsistence and commercial users. *Economic Botany*, 54(4), 513–527. <https://doi.org/10.1007/BF02866549>
- Damanik, R. and Djamaludin, R. (2012). Atlas Mangrove Teluk Tomini. Makassar, Indonesia: SUSCLAM Sustainable Coastal Livelihoods and Management Program.
- Dewanto, D.K., Finarti, F., Hermawan, R., Ndobe, S., Riyadi, P.H. and Tanod, W.A. (2019). Antioxidant Activity of Soft Corals Extracts from Palu Bay, Central Sulawesi, Indonesia. *Jurnal Pascapanen Dan Bioteknologi Kelautan Dan Perikanan*, 14(2), 163–178. <https://doi.org/10.15578/jpbkp.v14i2.583>
- Dewanto, D.K., Tanod, W.A., Finarti, F. and Renol, R. (2018). Screening of Antiradical Activity From Some Central Sulawesi Mangroves. *Pharmaciana*, 8 (1), 155–168. <https://doi.org/10.12928/pharmaciana.v8i1.8187>
- Dong, J., Wang, N.N., Yao, Z.J., Zhang, L., Cheng, Y., Ouyang, D., Lu, A.P. and Cao, D.S. (2018). ADMETlab: a platform for systematic ADMET evaluation based on a comprehensively collected ADMET database. *Journal of Cheminformatics*, 10 (29), 1–11. <https://doi.org/10.1186/s13321-018-0283-x>
- Drwal, M.N., Banerjee, P., Dunkel, M., Wettig, M.R. and Preissner, R. (2014). ProTox: a web server for the in silico prediction of rodent oral toxicity. *Nucleic Acids Research*, 42(W1), W53–W58. <https://doi.org/10.1093/nar/gku401>
- Eswaraiah, G., Peele, K. A., Krupanidhi, S., Kumar, R.B. and Venkateswarulu, T.C. (2020). Studies on phytochemical, antioxidant, antimicrobial analysis and separation of bioactive leads of leaf extract from the selected mangroves. *Journal of King Saud University - Science*, 32(1), 842–847. <https://doi.org/10.1016/j.jksus.2019.03.002>
- Guan, L., Yang, H., Cai, Y., Sun, L., Di, P., Li, W., Liu, G. and Tang, Y. (2019). ADMET-score – a comprehensive scoring function for evaluation of chemical drug-likeness. *Medicine Chemical Communication*, 10(1), 148–157. <https://doi.org/10.1039/C8MD00472B>
- Filimonov, D. and Poroikov, V. (2008). Probabilistic Approaches in Activity Prediction. In Varnek, A. and Tropsha, A. (Eds.). *Cheminformatics Approaches to Virtual Screening*, p. 182-216. Cambridge: RSC Publishing. <https://doi.org/10.1039/9781847558879-00182>
- Hosen, M.Z., Biswas, A., Islam, M.R., Bhuiyan, M.N.I. and Hossain, S.J. (2020). Comparison of physicochemical and antioxidant properties of edible fruits in the sundarbans' mangrove forest, Bangladesh. *Bangladesh Journal of Botany*, 49(3), 671–678. <https://doi.org/10.3329/bjb.v49i3.50009>
- Hsiao, T.H., Sung, C.S., Lan, Y.H., Wang, Y.C., Lu, M.C., Wen, Z.H., Wu, Y.C. and Sung, P.J. (2015). New anti-inflammatory cembranes from the cultured soft coral *Nephthea columnaris*. *Marine Drugs*, 13 (6), 3443–3453. <https://doi.org/10.3390/>

- md13063443
- Huliselan, Y.M., Runtuwene, M.R.J. and Wewengkang, D.S. (2015). Antioxidant activity of ethanol, ethyl acetate and n-hexane extract from seswanua leaves (*Clerodendron squamatum* Vahl.). *Pharmakon*, 4(3), 155–163. <https://doi.org/10.1016/j.taap.2009.09.009>
- Hwang, Y.P. and Jeong, H.G. (2010). Ginsenoside Rb1 protects against 6-hydroxydopamine-induced oxidative stress by increasing heme oxygenase-1 expression through an estrogen receptor-related PI3K/Akt/Nrf2-dependent pathway in human dopaminergic cells. *Toxicology and Applied Pharmacology*, 242(1), 18–28.
- Jacob, A.M., Suptijah, P. and Zahidah. (2013). Chemical composition, bioactive component and antioxidant activity of large-leafed mangrove (*Bruguiera gymnorrhiza*) fruit. *Jurnal Pengolahan Hasil Perikanan Indonesia*, 16(1), 86–94.
- Jariyah, Widjanarko, S.B., Yuniarta, Estiasih, T. and Sopade, P.A. (2014). Pasting properties mixtures of mangrove fruit flour (*Sonneratia caseolaris*) and starches. *International Food Research Journal*, 21(6), 2161–2167.
- Johnson, J.A., Johnson, D.A., Kraft, A.D., Calkins, M.J., Jakel, R.J., Vargas, M.R. and Chen, P. (2008). The Nrf2–ARE pathway an indicator and modulator of oxidative stress in neurodegeneration. *Annals of the New York Academy of Sciences*, 1147(1), 61–69. <https://doi.org/10.1196/annals.1427.036>
- Karim, M.A., Islam, M.A., Islam, M.M., Rahman, M.S., Sultana, S., Biswas, S., Hosen, M.J., Mazumder, K., Rahman, M.M. and Hasan, M.N. (2020). Evaluation of antioxidant, anti-hemolytic, cytotoxic effects and anti-bacterial activity of selected mangrove plants (*Bruguiera gymnorrhiza* and *Heritiera littoralis*) in Bangladesh. *Clinical Phytoscience*, 6(8), 1–12. <https://doi.org/10.1186/s40816-020-00162-3>
- Kardiman, Ridhwan, M. and Armi. (2017). Lindur fruit (*Bruguiera gymnorrhiza*) as food for the Acehnese archipelago. *Serambi Saintia*, 5(2), 51–55.
- Kim, J. and Kim, S. (2009). Anti-inflammatory activities of *Illicium anisatum* essential oil. *Acta Pharmaceutica*, 59, 289–300. <https://doi.org/10.2478/v10007-009-0022-y>
- Kusmana, C. (2014). Distribution and Current Status of Mangrove Forests in Indonesia. In Faridah-Hanum, I., Latiff, A., Hakeem, K.R., Öztürk, M. (Eds). *Mangrove Ecosystems of Asia: Status, Challenges and Management Strategies*, p. 37-60. New York: Springer-Verlag. https://doi.org/10.1007/978-1-4614-8582-7_3
- Kusmana, C., Hidayat, T., Istomo and Rusdiana, O. (2018). Growth performance of *Bruguiera gymnorrhiza* derived from cut-propagule seedling. *Biodiversitas*, 19(1), 208–214. <https://doi.org/10.13057/biodiv/d190128>
- Lalpuria, M., Anantheswaran, R. and Floros, J. (2012). Packaging technologies and their role in food safety. In Demirci, A. and Ngadi, M.O. (Eds). *Microbial decontamination in the food industry: Novel methods and applications*, p. 701–745. Cambridge, United Kingdom: Woodhead Publishing Limited. <https://doi.org/10.1533/9780857095756.4.701>
- Lamuela-Raventós, R.M. (2017). Folin-Ciocalteu method for the measurement of total phenolic content and antioxidant capacity. In Apak, R., Capanoglu, E. and Shahidi, F. (Eds). *Measurement of Antioxidant Activity and Capacity: Recent Trends and Applications*, p. 107–115. New Jersey, USA: John Wiley and Sons Ltd. <https://doi.org/10.1002/9781119135388.ch6>
- Lin, Y., Zheng, X., Chen, J., Luo, D., Xie, J., Su, Z., Huang, X., Yi, X., Wei, L., Cai, J. and Sun, Z. (2020) Protective Effect of *Bruguiera gymnorrhiza* (L.) Lam. Fruit on Dextran Sulfate Sodium-Induced Ulcerative Colitis in Mice: Role of Keap1/Nrf2 Pathway and Gut Microbiota. *Frontier Pharmacology*, 10, 1602. <https://doi.org/10.3389/fphar.2019.01602>
- Lisna, M.A. and Toknok, B. (2017). Potential of mangrove forest vegetation in the coastal area of the equatorial village, Tinombo Selatan sub-district, Parigi Moutong district. *Warta Rimba*, 5(1), 63–70.
- Li, Q., Nengwang, Y., Yanping, W., Yanping, S., Kangping, L. and Wenling, G. (2013). Extraction optimization of *Bruguiera gymnorrhiza* polysaccharides with radical scavenging activities. *Carbohydrate Polymers*, 96(1), 148–155. <https://doi.org/10.1016/j.carbpol.2013.03.054>
- Liu, J.K., Lin, H., Dong, Z.J. and Hu, Q. (2004). DPPH radical scavenging activity of ten natural p-terphenyl derivatives obtained from three edible mushrooms indigenous to China. *Chemistry and Biodiversity*, 1(4), 601–605. <https://doi.org/10.1002/cbdv.200490050>
- Lobo, V., Patil, A., Phatak, A. and Chandra, N. (2010). Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacognosy Reviews*, 4(8), 118–126. <https://doi.org/10.4103/0973-7847.70902>
- Ma, Q. (2013). Role of Nrf2 in oxidative stress and toxicity. *Annual Review of Pharmacology and Toxicology*, 53, 401–426. <https://doi.org/10.1146/annurev-pharmtox-011112-140320>

- Marzolini, C., Paus, E., Buclin, T. and Kim, R.B. (2004). Polymorphisms in human MDR1 (P-glycoprotein): recent advances and clinical relevance. *Clinical Pharmacology and Therapeutics*, 75(1), 13–33. <https://doi.org/10.1016/j.clpt.2003.09.012>
- Mohandas, G.G. and Kumaraswamy, M. (2018). Antioxidant Activities of Terpenoids from *Thuidium tamariscellum* (C. Muell.) Bosch. and Sande-Lac. a Moss. *Pharmacognition Journal*, 10(4), 645–649. <https://doi.org/10.5530/pj.2018.4.106>
- Molyneux, P. (2004). The use of the stable free radical diphenylpicryl-hydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin Journal of Science and Technology*, 26(2), 211–219.
- Mu, H., Gao, H., Chen, H., Tao, F., Fang, X. and Ge, L. (2013). A nanosized oxygen scavenger: preparation and antioxidant application to roasted sunflower seeds and walnuts. *Food Chemistry*, 136(1), 245–250. <https://doi.org/10.1016/j.foodchem.2012.07.121>
- Natesh, J., Mondal, P., Penta, D., Salam, A.A.A. and Meeran, S.M. (2021). Culinary spice bioactives as potential therapeutics against SARS-CoV-2: Computational investigation. *Computers in Biology and Medicine*, 128, 104102. <https://doi.org/10.1016/j.compbimed.2020.104102>
- Noor, Y.R., Khazali, M. and Suryadiputra, I.N. (2006). Panduan Pengenalaman Mangrove di Indonesia. 2nd ed. Bogor: Wetlands International Indonesia Programme.
- Pandey, K.B. and Rizvi, S.I. (2009). Plant Polyphenols as Dietary Antioxidants in Human Health and Disease. *Oxidative Medicine and Cellular Longevity*, 2(5), 270–278. <https://doi.org/10.4161/oxim.2.5.9498>
- Putra, M.Y., Murniasih, T., Swasono, R.T., Wibowo, J.T., Saputri, A.N.C., Widhiana, M.R. and Arlyza, I.S. (2016). Secondary metabolites and their biological activities in Indonesian soft coral of the genus *Lobophytum*. *Asian Pacific Journal of Tropical Biomedicine*, 6(11), 909–913. <https://doi.org/10.1016/j.apjtb.2016.08.011>
- Quan, Y., Jin, Y., Faria, T.N., Tilford, C.A., He, A., Wall, D.A., Smith, R.L. and Vig, B.S. (2012). Expression profile of drug and nutrient absorption related genes in madin-darby canine kidney (MDCK) cells grown under differentiation conditions. *Pharmaceutics*, 4(2), 314–333. <https://doi.org/10.3390/pharmaceutics4020314>
- Riyadi, P.H., Romadhon R., Anggo A.D., Herawati V.E. and Setyastuti A.I. (2020) PASS and ADMET analyses for eight compounds from Nile tilapia (*Oreochromis niloticus*) viscera waste hydrolysate as anti-inflammatory nutraceutical. *AAFL Bioflux*, 13 (5), 2630–2638.
- Riyadi, P.H., Tanod, W.A., Wahyudi, D., Susanto, E., Fahmi, A.S. and Aisiah, S. (2020). Potential of tilapia (*Oreochromis niloticus*) viscera bioactive peptides as antiviral for SARS-CoV-2 (COVID 19). *IOP Conference Series: Earth and Environmental Science*, 584, 012004. <https://doi.org/10.1088/1755-1315/584/1/012004>
- Rodríguez-López, P., Lozano-Sanchez, J., Borrás-Linares, I., Emanuelli, T.; Menéndez, J.A. and Segura-Carretero, A. (2020). Structure–Biological Activity Relationships of Extra-Virgin Olive Oil Phenolic Compounds: Health Properties and Bioavailability. *Antioxidants*, 9(8), 685–702. <https://doi.org/10.3390/antiox9080685>
- Rout, P. and Basak, U.C. (2014). Antioxidant properties in leaf and root extracts of some medicinally important mangrove species of Odisha Coast. *American Journal of Pharmtech Research*, 4(4), 605–617.
- Rout, P., Singh, S., Kumar, N. and Basak, U.C. (2015). Nutritional and antioxidant potential of some selected edible mangrove fruits of Odisha coast. *International Journal of Advances in Scientific Research*, 1(9), 349–355. <https://doi.org/10.7439/ijasr.v1i9.2624>
- Rijo, P., Simões, M.F., Duarte, A. and Rodríguez, B. (2009). Isopimarane diterpenoids from *Aeollanthus rydingianus* and their antimicrobial activity. *Phytochemistry*, 70(9), 1161–1165. <https://doi.org/10.1016/j.phytochem.2009.06.008>
- Szakács, G., Váradi, A., Özvegy-Laczka, C. and Sarkadi, B. (2008). The role of abc transporters in drug absorption, distribution, metabolism, excretion and toxicity (ADME-Tox). *Drug Discovery Today* 13(9–10), 379–393. <https://doi.org/10.1016/j.drudis.2007.12.010>
- Situmorang, R.O.P. and Barus, S.P. (2015). Mangrove management as source of food alternative by the women fishermen group in Sei Nagalawan, North Sumatra, Indonesia, presented at International Conference of Indonesia Forestry Researchers III, Bogor, 2015. Indonesia, Ministry of Environment and Forestry-Research, Development and Innovation Agency.
- Sudirman, S., Nurjanah and Jacoeb, A.M. (2014). Proximate compositions, bioactive compounds and antioxidant activity from large-leafed mangrove (*Bruguiera gymnorrhiza*) fruit. *International Food Research Journal*, 21(6), 2387–2391.
- Suh, S.S., Hwang, J., Park, M., Park, H.S. and Lee, T.K. (2014). Phenol content, antioxidant and tyrosinase inhibitory activity of mangrove plants in Micronesia.

- Asian Pacific Journal of Tropical Medicine*, 7(7), 531-535. [https://doi.org/10.1016/S1995-7645\(14\)60089-4](https://doi.org/10.1016/S1995-7645(14)60089-4)
- Tanod, W.A., Dewanto, D.K., Ndobe, S., Riyadi, P.H., and Putra, M.Y. (2019). Screening of antibacterial and antioxidant activity of soft corals *Sinularia* sp. and *Sarcophyton* sp. from Palu Bay Central Sulawesi. *Squalen: Bulletin of Marine and Fisheries Postharvest and Biotechnology*, 14(2), 73–83. <https://doi.org/10.15578/squalen.v14i2.394>
- Tanod, W.A., Yanuhar, U., Maftuch, Wahyudi, D. and Risjani, Y. (2019). DPPH scavenging property of bioactives from soft corals origin palu bay, Central Sulawesi, Indonesia. *IOP Conference Series: Earth and Environmental Science*, 236, 012121. <https://doi.org/10.1088/1755-1315/236/1/012121>
- Tungcharoen, P., Wattanapiromsakul, C., Tansakul, P., Nakamura, S., Matsuda, H. and Tewtrakul, S. (2019). Anti-inflammatory effect of isopimarane diterpenoids from *Kaempferia galanga*. *Phytotherapy Research*, 34(3), 612-623. <https://doi.org/10.1002/ptr.6549>
- Van de Waterbeemd, H. and Gifford, E. (2003). ADMET *in silico* modelling: towards prediction paradise?. *Nature Reviews Drug Discovery*, 2(3), 192-204. <https://doi.org/10.1038/nrd1032>
- Wibawanti, J.M.W., Fadhiliya, L., Pamungkas, S. and Mudawaroch, R.E. (2018). Mangrove Processed Alternative Functional Food Production in Purworejo Regency. *Community Empowerment*, 3 (1), 27–33. <https://doi.org/10.31603/ce.v3i1.2450>
- Yamaguchi, T., Takamura, H., Matoba, T. and Terao, J. (1998). HPLC method for evaluation of the free radical-scavenging activity of foods by using 1, 1-diphenyl-2-picrylhydrazyl. *Bioscience Biotechnology Biochemistry*, 62(6), 1201–1204. <https://doi.org/10.1271/bbb.62.1201>
- Yu, L., Haley, S., Perret, J., Harris, M., Wilson, J. and Qiant, M. (2002). Free radical scavenging properties of wheat extracts. *Journal of Agricultural and Food Chemistry*, 50(6), 1619–1624. <https://doi.org/10.1021/jf010964p>
- Zhu, Z., Chen, J., and Zheng, H.-L. (2012). Physiological and proteomic characterization of salt tolerance in a mangrove plant, *Bruguiera gymnorrhiza* (L.) Lam. *Tree Physiology*, 32(11), 1378–1388. <https://doi.org/10.1093/treephys/tps097>