



OVERVIEW OF PREDATORS IN SHALLOTS PLANTATION IN PEATLAND, LANDASAN ULIN SOUTH BORNEO

MUHAMAMAD INDAR PRAMUDI*, SAMHARINTO, HELDA ORBANI ROSA,
SALAMIAH AND LYSWIANA APHRODYANTI

Departement of Plant Protection, Faculty of Agriculture, Lambung Mangkurat University

*Email: indar_pramudi@yahoo.com (corresponding author): ORCID ID 0000-0003-1085-5873

ABSTRACT

Species abundance and diversity of predatory arthropods are important information for developing biological pest control strategies. A study that aims to determine the abundance of predators (Coccinellidae and Araneae) in peatlands planted with shallots in Tegal Arum Village, Landasan Ulin District, Banjarbaru City, South Borneo has been carried out. The method used is purposive sampling method. Selecting plots with uniform growth and plant height of shallots in one plot or beds measuring 3 x 20 m and six plots were observed using three methods: direct sampling, swing nets, and pitfall traps. Observations on species abundance of spiders and predatory coccinellids were carried out to describe the number of species and the abundance of insects in shallot cultivation on peatlands. Predators from the order Coleoptera of the Coccinellidae family observed belong to (*Menochilus sexmaculatus*, *Micrapsis* sp, *Coccinella novemnotata* and *Scymnus* sp. with 143 individuals and the highest population was found by *Micrapsis* sp (128 individuals). Meanwhile, there were 6 species of spiders (Araneae) from 5 different families (Oxyopidae, Tetragnatidae, Lynphiidae, Thomisidae, and Lycosidae) with the most individuals being *Oxyopes variabilis* (Araneidae: Oxyopidae). The distribution index of species diversity (E) shows $E > 0.6$, indicating that the evenness of species on the land is uneven

Key words: Abundance, araneae, coccinellidae, coleoptera, diversity, families, peatland, predator, shallots, spesies

Plant pest population management is carried out by paying attention to the agro-ecosystem. Especially the biotic factors which consist of insect pests, natural enemies, both predators, and parasitoids (Chakravarthy et al., 2016). The approach to studying the structure of the agroecosystem is to study the biodiversity of the cultivated land, in this research the land used is peatland for shallot cultivation in South Kalimantan. Coleoptera predators are biological control agents that are found in various types of agricultural land. This insect has a large number of species with a cosmopolitan distribution (Efendi et al., 2016). These Coccinellidae is a well-known beetle family, consist of seven families, i.e. Epilachninae, Coccinellidae, Chilocorinae, Coccidulinae, Ortaliinae, Scymninae, and Sticholotidinae (Pope, 1988). Except for the mycophagous Coccinellinae (Halyziini and Tythaspis) and the phytophagous Epilachninae, all remaining coccinellids are predators of hemipteran insects from the suborder Sternorrhyncha (e.g. aphids, scales, psyllids and whiteflies), mites eventually other insect larvae and insect eggs (Amir, 2002; Dixon, 2000; Damayanthi, 2016). Fiaboe et al., (2007) stated that as many as 5200 species of predatory Coccinellidae have been identified. Foltz (2002) estimates that there are 5000 species of

predatory Coccinellidae worldwide, while according to Vandenberg (2009); Kundoo and Khan (2017) predatory Coccinellidae is 6000 species and are found in mountainous areas, agricultural areas, coastal areas to urban areas and 90% predatory. Coccinellidae has fairly high diversity, it is estimated that there are 5000 species worldwide, while in Indonesia it is estimated that more than 300 species are widespread (Nelly et al., 2015). Many types of predatory Coccinellidae in Indonesia have great potential in controlling the population of various types of plant pests (Aprila et al., 2019). The Coleoptera represent one of the most abundant animal groups (Sushko, 2018) and important natural enemies of many insect pests in terrestrial ecosystems such as peatlands. They are important members of food webs, providing protein for species at higher trophic levels (Sushko 2012; Sushko 2017). In Soraya's research (2016), nine species of predatory Coccinellidae were found in eggplant plantations, i.e. *Epilachna vigintioctopunctata*, *E. borealis*, *Illeis flava*, *Harmonia consformis* and *H. testudinaria*. Meanwhile, Efendi et al. (2016) found ten predatory Coccinellidae viz., *Chilocorus melanophthalmus*, *Coelophora maculata*, *C. inaequalis*, *C. reniplagiata*, *C. bisellata*, *Ropaloned a decussate*, *Verania discolor*. In eggplant and chili

three species were found *Coccinella transversalis*, *Menochilus sexmaculatus*, and *Verania lineata*. This study explores abundance of Coccinellidae predators and spiders as natural enemies.

MATERIALS AND METHODS

This research was conducted in Tegal Arum Village (-3°25'44" S, 11°44'69" E, 10.8 m, 284°), Landasan Ulin District, Banjarbaru City, South Borneo, Indonesia. The research was conducted from August to October 2020. Species diversity was analyzed using the Shannon-Wiener index (H'), the Simpson Domination Index (D), and Evenness (E) using the Magurran handbook (Magurran 1988). If the value of $H' < 1$ then the community is less stable, if the value of H' is between 1-2 then the community is stable, and if the value of $H' > 2$ is said to be very stable (Kent and Paddy, 1992). The criteria for the level diversity based on this index are high if the value of $H' > 3.5$, moderate if $H' = 1.5 - 3.5$, and low if $H' < 1.5$. Odum (1993) stated that the criteria for dominance if the C value approaches 0 (<0.5), then no species dominate and if the C value approaches 1 (≥ 0.5), then there is a dominant species. Observations of Predator. Using purposive sampling method, by selecting plots with uniform growth and plant height of shallots in one plot or beds measuring 3 x 20 m. The observed 6 plots. Observations were made using three methods: direct sampling, sweeping nets, and pitfall traps. The determination of the point for trapping is done deliberately. 3 traps were placed in the front, middle and back of the plot with a distance of 5 meters between the traps, placed in the center of the plot. Observations were made at the age of 0, 28, 42, 56 dd. Installation of pitfall traps for 24 hrs, at the same time as other observations. Sampling was done by direct collection. Hand picking on the sample plots. Sweeping net. Insect sweeping nets were used with 10 double swings across the plot area vertically and three captures were made for each observation. Pitfall trap were used to observe ground-level arthropods made of plastic cups with a diameter of 50 mm and a depth of 100 mm. This trap was filled with a solution of water and four drops of liquid detergent (10:1). This trap is filled with a solution of water and four drops of liquid detergent (10:1), the solution being filled with only one third of the total volume. Then placed in a hole that has been provided according to the size of the plastic cup, placed for 24 hours on the ground. The traps are cleaned and bottled for identification. Predator identification was carried out by observing insect specimens, especially the Coccinellidae, Coleoptera and spiders

(Family Araneidae, Araneae). Predatory Coccinellidae specimens obtained in the field were identified to species level by matching the specimens with pictures and descriptions from the identification key book. Identification of specimens using morphological characteristics of the wings, antennae, and thorax. Predatory Coccinellidae specimens obtained in the field were identified to species level using the identification key of Khan et al. (2006), Stephens and Losey (2004), Kalshoven (1981), McAlpine, et al. (1987), and Barrion and Litsinger (1995). To identify spiders using Wegner, (2009), Jocqué and Anna (2007), and Heisswolf, S., et al. (2010). Species composition data and the number of individual predators of Coccinellidae and spiders were used to analyze abundance and evenness. The measure of abundance used is the value of the Shannon-Wiener species diversity index (H') and Evenness (E) using Magurran's (1988) book.

RESULTS AND DISCUSSION

The land for shallot cultivation in the Tegal Arum is a peatland that has just been cleared and is starting to be planted with Citrus, sugar cane, chilies, and shallots. Based on the results of observations that have been made, 217 individual predators were collected. The results showed that the predators of the order Coleoptera family Coccinellidae were 4 species with 143 individuals and the highest population was found by *Micrapsis* sp (128). Meanwhile, there were 6 species of spiders with the most individuals being *Oxyopes variabilis* (22) Araneidae: Oxyopidae and the lowest was *Lycosidae* sp (Araneidae: Lycosidae) with 4 individuals (Table 1).

Predators from coccinellidae have more individuals than Aranea but the number of Aranea taxa are Coccinellids function in complex food webs as predators, as non-prey food consumers, and as prey or host of natural enemies. In the life stage of the coccinellids, the eggs are particularly susceptible to predation, and the coccinellids are behaviorally adapted to reduce egg predation from heterospecific predators (Weber and Jonathan, 2009).

Several papers have discussed the positive and negative effects of these two predators in new environments (Camacho-Cervantes et al., 2017; Kenis et al., 2017; Koch and Costamagna, 2017; Riddick, 2017). The second interaction affects community structure and predator-prey dynamics in various ecosystems, including agro-ecosystems. Thus, the expansion of the range of these predatory species has both negative and

Table 1. Diversity and abundance of predators in shallot peatland agroecosystems (Tegal Arum Village, Landasan Ulin District, Banjarbaru City)

No.	Taxon	Ordo	Family	Number of insects caught				Total
				Plant age				
				0	28	42	56	
1	<i>Menochilus sexmaculatus</i>	Coleoptera	Coccinellidae	8	1	0	0	9
2	<i>Micrapsis</i> sp	Coleoptera	Coccinellidae	43	45	30	10	128
3	<i>Coccinella novemnotata</i>	Coleoptera	Coccinellidae	2	1	0	0	3
4	<i>Scymnus</i> sp	Coleoptera	Coccinellidae	1	1	1	0	3
								143
1	<i>Oxyopes variabilis</i>	Araneidae	Oxyopidae	4	9	4	5	22
2	<i>Oxyopes</i> sp	Araneidae	Oxyopidae	3	2	0	0	5
3	<i>Tetragnatha</i> sp	Araneidae	Tetragnathidae	4	2	4	1	11
4	<i>Linyphiidae</i> sp	Araneidae	Lyniphiidae	10	6	2	0	18
5	<i>Vitia misumena</i>	Araneidae	Thomisidae	5	6	1	2	14
6	<i>Lycosidae</i> sp	Araneidae	Lycosidae	0	2	2	0	4
								74
Number of taxa				9	10	7	4	10
Total				80	75	44	18	217

positive ecological effects on biodiversity and levels of biological control. Although many studies examine the spatial and temporal patterns among predatory coccinellidae and predatory aranaea. However, the mechanisms involved in the coexistence of this species are still limited (Hongran et al., 2021). *Micrapsis* sp had the highest individual among the others. It is suspected that this species is a species whose natural habitat is in peatlands. This is supported by research by Soedijo et al. (2012) and Samharinto et al. (2011). *Micrapsis* sp was found in large numbers from three research sites on peatland planted with rice in South Kalimantan. When viewed from the length of the life cycle, *Micrapsis* sp were higher than the males in longevity and predation rate. Longer than other species, which ranged from 1-2 weeks to develop from eggs to adults. Chaudhary et al. (2015) in their research stated that female *Menochilus sexmaculatus* (Fabricius) consumed higher than other Coccinellidae. However, Hllaing et al. (2017) stated otherwise, the predation rate of 1st instar grub was 37.5 ± 4.6 and 4th instar grub was 51.7 ± 9.8 . The male predation rate (51.1 ± 16.1) and female predation rate (57.8 ± 16.4) were observed under the same conditions. It is possible that this is the basis for this species to dominate in onion fields. In observations this type is found in grasses, shrubs. Meanwhile, *Menochilus sexmaculatus*, *Coccinella novemnotata* and *Scymnus* sp were not found. According to Puspasari et al. (2016), *Menochilus sexmaculatus* is more commonly found in peanuts, soybeans, beans, corn, eggplant and vegetables.

Coccinella novemnotata in chili plants attacked by *Bemisia tabaci* and *Scymnus* sp was more commonly found in plants infected with mealybugs (*Paracoccus marginatus*) (Tairas et al., 2015). In the research area, eggplant, chili were planted, while banjar siam oranges and guava were attacked by mealybugs. It is possible that the three predators apart from this *Micrapsis* sp, have already gotten their food from these plants, so they are rarely found in shallot plantations.

Spiders are common predators that can eat a wide variety of prey (Foelix, 2011) and are very effective in managing pests and reducing crop damage. Spiders are very abundant in agriculture and if conserved and the population is plentiful, they can control insect pests (Basnet and Mukhopadhyay, 2014). At the initial observation, in the uncultivated land found various types of leafhoppers and grasshoppers. It is suspected that this is a supporting factor why many types of spiders are found. The ability of spiders to prey on the main insect pests of brown planthoppers, their ability to prey on nymphs and imago ranges from 5-15 tails/day and can also become cannibals for fellow species or different species (Soedijo and Indar, 2015). It is also suspected that the clearing of land for shallot plantations causes spiders to lose their habitat so that their population also decreases and food decreases. Each spider species has a different active time and prey species, so it is very important to manage the diversity of spider species in an area (Wayan and Hery, 2013). So that the time

of observation in the field also affects the number of individual spiders caught in the observation.

The highest individual in Araneae is *Oxyopes* sp. ratter (lynx spider) and *Lycosidae* sp. (wolf spider) was the lowest among the Araneae. Lynx spiders are roving hunters commonly associated with the shrubby and grassy understory and are small and slender. This is in contrast to the larger wolf spiders and fast-moving ground hunters. The research of Memah et al. (2018) states, the preying ability of *Oxyopes* sp. on imago is the highest compared to others. The families Agelenidae, Araneidae, Tetragnathidae, and Thomisidae are spiders that are nocturnal or nocturnal. On the other hand, Lyniphiidae is a family of spiders that are active during the day or diurnal. Costello and Daane (2005) also found that most members of the Araneidae and Lycosidae are nocturnal, while all Salticidae are diurnal. *Oxyopidae* is more in the field because this species was active during the day and night or 24 hrs.

There is a tendency for the predatory population of Coccinellidae and Araneidae to decrease with the increasing age of shallot plants. Successively, at 0 days to 56 days after planting the population was 80, 75, 44, 18 predators. There was a decrease in species diversity and the number of individuals along with the age of the shallot plants. The land used in this study is peatland, where this land is formed from organic materials, such as leaves, stems, branches, and plant roots. Likewise, according to Sugiyarto (2007), organic matter which is difficult to decompose will function as a ground cover so that it can be used as a good source of energy for the insect community. Therefore, trophic composition of predator assemblage can play an important role in determining the nature of the relationship between predator diversity and ecosystem function (Finke and Denno, 2005). Organic matter has an important role to support plants and soil organisms so that if the soil organic matter content decreases, the soil's ability to support the productivity of plants and soil organisms also decreases. Naturally, the number of insects in this study predators Coccinellidae and spiders is more at the beginning of the observation.

The population of pest predators at the beginning of planting (0 days) was more diverse and the numbers were more numerous plus there were still weeds and bushes. On the land, grasses, shrubs and shrubs were found such as *Leersia hexadra*, *Hymenachne acutigulma*, *Brachiaria* sp. *Pennisetum purpureum*, *Panicum* sp, *Paspalum* sp. *Stenochlaena palustris*, *Chromolaena*

odorata, *Xyris indica* and *Melastoma malabathricum*. Wild plants or weeds are important agro-ecosystem components because they can positively influence the biology and dynamics of natural enemies (Altieri and Nicholls, 2004; Puspasari et al., 2016). This plant is an alternative host, a place to hide if predators or natural enemies experience bad environmental conditions (Van Emdem, 1991; Altieri and Nicholls, 2004). The cropping pattern used also affects the diversity and population of insects in a land. This study used a shallot monoculture pattern. According to Semiun and Stefanus (2016) and Melhanon and Dewi (2020), the diversity and abundance of monoculture farming systems of arthropods are lower than that of polyculture systems.

Besides, the growth phase of the shallot also affects the abundance of these two predators. This is supported by Kaleb et al. (2015) which states that the height and low level of individual insects as natural enemies of the crown and soil surface indicate that the availability of available food sources is closely related, i.e. conformity to the growth phase of plants that provide a source of food for the growth and development of natural enemy insects, the reduction of pests that have the potential to serve as a food source for natural enemy insects.

At the beginning of soil cultivation or 0 days, the diversity index showed a moderate and stable category (1.57), along with land cultivation and cleaning of weeds and planting into monocultures (shallots), the diversity index decreased to a low (<1.5) (Table 2). This is thought to be due to land clearing and weed clearing, loss of housing for various types of insects, both pests and predators. Many types of planthoppers (zig-zag planthoppers, white-backed planthoppers, brown leafhoppers, green leafhoppers, leafhoppers, white and yellow rice stem borers and grasshoppers were found before land clearing. All these are food for spiders and coccinellid beetles. Changes occur in the food web or trophic transfer. Yang et al. (2018) observed the effects of predator diversity on prey biomass and

Table 2. Diversity index, evenness index of predatory (Coccinellidae and Araneida species) in shallot crop ecosystems on peatlands in South Kalimantan

Observation (days)	Index value	
	Diversity (H')	Evenness (E)
0	1.57	0.71
28	1.43	0.62
42	1.15	0.59
56	1.09	0.78

trophic transfer efficiency (using the predator/prey biomass ratio). Higher prey diversity increases predator diversity and biomass, as well as trophic transfer efficiency, which may result from a more balanced diet and/or increased niche complementarity due to higher prey diversity. This effect of prey diversity affects predator-prey interactions in natural ecosystems. Weber and Jonathan (2009) and Wilby et al. (2005) stated that the effects of species diversity appear in natural enemy assemblages depending on the context; they depend not only on the characteristics of the predatory species but on the identity of the species they prey on. This effect of prey diversity affects predator-prey interactions in natural ecosystems.

ACKNOWLEDGEMENTS

The Dean, Faculty of Agriculture Lambung Mangkurat University and the Head of the Plant Protection Study program are acknowledged.

FINANCIAL SUPPORT

No financial support received.

AUTHOR CONTRIBUTION STATEMENT

Salamiah and Helda conceived and designed research. Samharinto and Muhammad conducted experiments. Pramudi and Lyswiana contributed analyzed data. All authors read and approved the manuscript.

CONFLICT OF INTEREST

No conflict of interest.

REFERENCES

Altieri M A, Nicholls C I. 2004. Biodiversity and pest management in agroecosystem. Second Edition. New York: Chapman and Hall. 236 p.

Amir. 2002. Kumbang Lembing Pemangsa Coccinellidae (Coccinellinae) di Indonesia. Bogor. Puslit Biologi-LIPI.

Aprila, Meli, Rover R, Siska E. 2019. Diversitas Coccinellidae predator pada ekosistem Pertanian Cabai di Tiga Kecamatan Kabupaten Kuantan Singingi. Jurnal Agronomi Tanaman Tropika (Juatika) 1(1): 32-41.

BP2LHK. 2017. Makrofauna indikator Perubahan ekosistem Lahan Gambut. Artikel. 8 Juni 2017. <http://foreibanjarbaru.or.id/archives/3244>.

Barrion A, Litsinger J. 1995. Riceland spiders of South and South East Asia. Entomology Division, International Rice Research Institute, Los Banos, Philippines.

Basnet K, Mukhopadhyay A. 2014. Biocontrol potential of the lynx spider *Oxyopes javanus* (Araneae: Oxyopidae) against the tea mosquito

bug, *Helopeltis theivora* (Heteroptera: Miridae). International Journal Tropical. Insect Science 34(4): 232-238.

Camacho-Cervantes M, A Ortega-Iturriaga, E Del-Val. 2017. From effective biocontrol agent to successful invader: the harlequin ladybird (*Harmonia axyridis*) as an example of good ideas that could go wrong. Peer J 5: 3296.

Chakravarthy A K, Kammar V, Shashank P R. 2016. Arthropods: Evolution and ecology. Chakravarthy A, Sridhara S. (eds) Economic and ecological significance of arthropods in diversified ecosystems. Springer, Singapore. 1-16 pp. https://doi.org/10.1007/978-981-10-1524-3_1

Chaudhary D, Kumar B, Mishra G, Omkar. 2015. Resource partitioning in a ladybird, *Menochilus sexmaculatus*: function of body size and prey density. Bulletin of Entomological Research 105(1): 121-128.

Costello, M J, Daane, K M. 2005. Day vs Night sampling for spiders in grape vineyards. Journal of Arachnology 33:25-32.

Damayanthi E. 2016. Keanekaragaman Coccinellidae predator Pada Pertanaman Padi di Dataran Rendah dan Dataran Tinggi di Sumatera Barat. (Skripsi). Universitas Andalas. Padang.

Dixon A F G. 2000. Insect predator-prey dynamics. Lady birds beetles and biological control. London, Cambridge University Press. 257 pp.

Efendi S, Yaherwandi, Novri N. 2016. Analisis keanekaragaman Coccinellidae predator dan Kutu Daun (*Aphididae* spp) pada ekosistem Pertanian Cabai. J. Bibiet 1(2): 67-80.

Fiaboe K K M, Gondim M G C, de Moraes, G J, Ogotland, C K, Knapp M. 2007. Bionomics of the Acarophagous Ladybird Beetle *Stethorus Tridens* Fed *Tetranychus evansi*. Journal Appl. Entomology 131: 355-361.

Finke D L, Denno R F. 2005. Predator diversity and the functioning of ecosystems: the role of intraguild predation in dampening trophic cascades. Ecology Letters 8: 1299-1306. <https://doi.org/10.1111/j.1461-0248.2005.00832.x>

Foelix R F. 2011. Biology of Spiders. New York: Oxford University Press Incorporation.

Foltz J L. 2002. Coleoptera: Coccinellidae. Dept of Entomology and Nematology. University of Florida. <http://entomology.ifas.ufl.edu/Coleoptera/Coccinellidae.html>.

Heisswolf S, Walsh B, Bilston L. 2010. Identification of insects, spiders and mites in vegetable crops: trainer's handbook. (Teaching Resource). The State of Queensland (Department of Employment, Economic Development and Innovation). 78 p.

Hllaing N Z, Thiha, W T, Weine, N N O. 2017. The biology of *Micraspis discolor* and study of its predation Efficiency on black aphid. Proceedings of 105th. The IIER International Conference, Bangkok, Thailand, 5th-6th June 2017.

Hongran L, Baoping L, Gábor L L, Timothy J K, John J O. 2021. Interactions among native and non-native predatory coccinellidae influence biological control and biodiversity. Annals of the Entomological Society of America 114(2): 119-136. Doi: 10.1093/aesa/saaa047.

Jocqué R, Anna S D S. 2007. Spider families of the world. Royal Museum for Central Africa 13, Leuvensesteenweg 3080 Tervuren (Belgium). www.africamuseum.be. 338 p.

Kaleb R, Flora P, Nur K. 2015. Keanekaragaman Serangga Musuh Alami pada Pertanaman Bawang Merah (*Allium ascalonicum* L) yang Diaplikasi dengan Bioinsektisida *Beauveria bassiana* (Bals.-Criv.) Vuill. J. Agroland 22(2): 114-122.

Kalshoven L G. 1981. The pest of crops in Indonesia. Van Hoeve, Jakarta. P. T. Ichtiar Baru.

- Kenis M, Adriaens T, Brown P M J A, Katsanis G, San Martin, Branquart E, Maes D, Eschen R, Zindel R, Van Vlaenderen J. 2017. Assessing the ecological risk posed by a recently established invasive alien predator: *Harmonia axyridis* as a case study. *BioControl* 62: 341-354.
- Khan AA, Zaki F A, Khan Z H, Mir R A. 2016. Biodiversity of predaceous ladybird beetles (Coleoptera: Coccinellidae) in Kashmir. *Journal Biol. Control* 23(1): 43-47.
- Koch R L, Costamagna A C. 2017. Reaping benefits from an invasive species: role of *Harmonia axyridis* in natural biological control of *Aphis glycines* in North America. *BioControl* 62(3): 331-340. DOI: 10.1007/s10526-016-9749-9
- Kundoo AA, Khan AA. 2017. Coccinellids as biological control agents of soft bodied insects: A review. *Journal of Entomology and Zoology Studies* 5(5): 1362-1373.
- Puspasari L T, Martua S S, Sri H. 2016. Komposisi Komunitas Serangga Aphidophaga dan Coccidophaga pada Agroekosistem Kacang Panjang (*Vigna sinensis* L.) di Kabupaten Garut. *Jurnal Agrikultura* 27(1): 30-37.
- McAlpine J F, Peterson B V, Shewell G E, Teskey H J, Vockeroth J R, Wood D M. (1987). *Manual of Nearctic Diptera*. Agriculture Canada, Research Branch.
- Magurran A E. 1988. *Ecological diversity and its measurement*. Princeton University Press, Princeton. Scientific Research an Academic Publisher.
- Melhanah Lilies S, Dewi S. 2020. Struktur Komunitas Arthropoda Nocturnal pada Jagung Manis dan Kacang Panjang Organik dan Konvensional di Lahan Gambut. *Daun: Jurnal Ilmiah Pertanian dan Kehutanan* 7(10):11-22. <https://doi.org/10.33084/daun.v7i1.1603>.
- Memah V V, James B K, Trina E T. 2018. Preying ability of spiders *Oxyopes* sp. and *Pardosa* sp. on four prey types of insect pests under laboratory conditions. *Bioscience Research* 15(1): 145-151.
- Nelly N, Yاهرwandi and Muhammad S E. 2015. Keanekaragaman Coccinellidae Predator dan Kutu Daun (*Aphididae* spp.) pada Ekosistem Pertanian Cabai. *Pros Sem Nas Masy Biodiv Indon* 1(2): 247-253. Doi: 10.13057/psnmbi/m010213
- Pope R D. 1988. A revision of the Australian Coccinellidae (Coleoptera): Subfamily Coccinellinae. *Invertebrate Taxonomy* 2(5): 633 -735.
- Riddick E W. (2017). Spotlight on the positive effects of the ladybird *Harmonia axyridis* on agriculture. *BioControl* 62(3): 319-330. DOI:10.1007/s10526-016-9758-8.
- Samharinto, Abdul LA, Bambang TR, Hakimah H. 2011. Keanekaragaman Arthropoda pada Persawahan Irigasi di Kalimantan Selatan Studi Kasus di Desa Sungai Rangas. *Prosiding Seminar Nasional Biologi XXII Perhimpunan Biologi Indonesia. Peran Biologi dalam Pendayagunaan Bioresources Indonesia untuk Meningkatkan Daya Saing Bangsa*. 31 Agustus – 1 September 2013.
- Semiun C G, Stefanus S. 2016. Kelimpahan dan Keanekaragaman Arthropoda Tanah pada Lahan Pertanian Monokultur dan Polikultur di Desa Labat Kupang. *BioWallacea* 2(2): 154-161.
- Sugiyarto. 2007. Konservasi Makrofauna Tanah dalam Sistem Agroforestri. *Puslitbang Bioteknologi dan Biodiversitas LPPS. UNS Surakarta*.
- Soedijo S, Abdul LA, Bambang TR, Hakimah H. 2012. The increase of arthropods biodiversity in paddy field ecosystem managed by using integrated pest management at South Borneo. *Journal of Tropical Life Science* 2(3): 72-76.
- Soraya I. 2016. Jenis-Jenis Coccinellidae (Coleoptera) pada Tanaman Terung (*Solanum melongena* L.) di Nagari Paninjauan Kecamatan X Koto Kabupaten Tanah Datar. *Skripsi. Program Studi Pendidikan Biologi STKIP PGRI Sumatera Barat*. Padang.
- Stephens E J, Losey J E. 2004. Comparison of Sticky Cards, Visual and Sweep Sampling of Coccinellid Populations in Alfalfa. *Environmental Entomology* 33(3): 535-539. 1 June 2004. <http://doi.org/10.1603/0046-225X-33.3.535>.
- Sushko G. 2012. *The Insect Fauna of Yelnia Peat Bog (North-west Belarus)*. Lambert Academic Publishing, Saarbrücken, Germany, 113 pp.
- Sushko G G. 2017. Diversity and species composition of beetles in the herb-shrub layer of a large isolated raised bog in Belarus. *Mires and Peat* 19(10): 1-14. DOI:10.19189/MaP.2017.OMB.266.
- Sushko G G. 2018. Effect of vegetation cover on the abundance and diversity of ladybirds (Coccinellidae) assemblages in a peat bog. *Biologia (Bratislava)* 73(4): 371-377.
- Tairas R W, Max T, Jantje P. 2015. Musuh Alami Kutu Putih *Paracoccus marginatus* Williams & Granara de Willink (Hemiptera: Pseudococcidae) Pada Tanaman Pepaya di Minahasa Utara. *Eugenia*, 21(2): 62-69.
- Van Emdem H F. 1991. Plant diversity and natural enemy efficiency in agroecosystem *in*: Mackkauer M, Ehler L E, Roland J, editor. *Critical Issue in Biological Control*. Great Britain: Atheneum Press.
- Vandenberg N J. 2009. The new world Genus *Cycloneda* (Coleoptera: Coccinellidae: Coccinellini): Historical Review, New Diagnosis, New Generic and Specific Synonyms, and An Improved Key to North American Species. *Entomological Society of Washington* 104(1): 221-236.
- Wayan S, Hery H. 2013. Diversity of spiders and their potentials as natural enemies of cashew pests. *Indonesian Journal of Entomology* 10(1): 24-30. DOI: 10.5994/jei.10.1.24.
- Weber D C, Jonathan G L. 2009. Assessing the trophic ecology of the coccinellidae: Their roles as predators and as prey. *Biological Control* 51(2009): 199-214. <https://doi.org/10.1016/j.biocontrol.2009.05.013>.
- Wegner W S. 2009. *Spider Identification Guide*. BASF. 60 p.
- Wilby A, Villareal S C, Lan L P, Heong K L, Thomas M B. 2005. Functional benefits of predator species diversity depend on prey identity. *Ecological Entomology* 30(5): 497-501. Doi:10.1111/j.0307-6946.2005.00717.x
- Yang J W, Wenxue W, Chin C H, Kuo P C, Gwo C G, Chih H H. 2018. Predator and prey biodiversity relationship and its consequences on marine ecosystem functioning—interplay between Nanoflagellates and Bacterioplankton. *The ISME Journal* 12: 1532-1542. <https://doi.org/10.1038/s41396-018-0111-3>

(Manuscript Received: September, 2023; Revised: December, 2023;

Accepted: December, 2023; Online Published: January, 2024)

Online First in www.entosocindia.org and indianentomology.org Ref. No. e23598