

Identification and abundance of fruit fly species (*Bactrocera mcgregori*) in melinjo plant (*Gnetum gnemon*) in Damit Village, Batu Ampar Sub-district, Tanah Laut Regency

by Noor Aidawati

Submission date: 13-Aug-2024 11:39PM (UTC-0400)

Submission ID: 2431806576

File name: n_Damit_Village,_Batu_Ampar_Sub-district,_Tanah_Laut_Regency.pdf (502.34K)

Word count: 2726

Character count: 14037



RESEARCH PAPER

6
International Journal of Biosciences | IJB |
ISSN: 2220-6655 (Print), 2222-5234 (Online)
<http://www.innspub.net>
Vol. 14, No. 2, p. 146-152, 2019

OPEN ACCESS

2
Identification and abundance of fruit fly species (*Bactrocera mcgregori*) in melinjo plant (*Gnetum gnemon*) in Damit Village, Batu Ampar Sub-district, Tanah Laut RegencyMuhammad Rahmani¹, M. Indar Pramudi^{2*}, Noor Aidawati³¹Agroecotechnology Study Program, Program in Plant Protection, Faculty of Agriculture, Lambung Mangkurat University, Indonesia²Plant Protection Study Program, Faculty of Agriculture, Lambung Mangkurat University, Indonesia**Key words:** *Bactrocera mcgregori*, *Gnetum gnemon*, Abundance, Sex Ratio.<http://dx.doi.org/10.12692/ijb/14.2.146-152>16
Article published on February 12, 2019**Abstract**

Research on fruit flies in melinjo plants in Indonesia has been carried out but will only be limited to locations on the island of Java, so it is necessary to prove whether fruit fly species that attack melinjo in South Kalimantan are the same or different from the results of previous studies. The locations chosen for this study were Damit Village, Batu Ampar Sub-district, Tanah Laut Regency has many melinjo plantations which are attacked by fruit flies; therefore, it is necessary to identify the species of fruit flies and to find out the abundance and sex ratio of the species population. The study was conducted by collecting the attacked fruit from five different points and putting the fruit in jars filled with wood sawdust. The results of the identification showed that the fruit flies that attacked melinjo fruit was *Bactrocera mcgregori*. The highest abundance of *B. mcgregori* was 75.6% and the lowest was 8.4% with an average of 37.06%. The highest sex ratio of *B. mcgregori* was 0.88 and the lowest was 0.61 with an average of 0.78. The attack of *B. mcgregori* fruit flies on melinjo fruit was the first report in South Kalimantan.

***Corresponding Author:** M. Indar Pramudi ✉ indar_pramudi@yahoo.com

Introduction

The production of melinjo in Indonesia in 2014 was 197.647 tons and the harvested area was 15.383 ha with an average production of 12.85 tons/ha (Kementan / Ministry of Agriculture, 2015). The production of melinjo in Tala Regency in 2015 was 213.3 tons from the harvested area of 63.856 ha with an average production of 33.40 tons ha⁻¹ (BPS, 2016) although melinjo plants were often attacked by various pests.

One of the pest problems encountered in the cultivation of melinjo plants (*G. gnemon*) is fruit fly pests from the *Bactrocera mcgregori* species (Bezzi). This fruit fly is found in Indonesia, Singapore and the Philippines. The *B. mcgregori* (Bezzi) is a species of fruit flies that attacks melinjo plants (Siwi *et al.*, 2006). The presence of the fruit fly larvae on melinjo fruit peels has not been considered to cause economic losses because the utilization and selling price of melinjo peels are quite low. The melinjo peel is useful as an antioxidant which is efficacious to increase endurance and natural antimicrobials because it contains phenolic proteins and flavonoids (Santoso *et al.*, 2010).

The attack of *B. mcgregori* fruit flies only causes damage to the fruit peel tissue. Presently the *B. mcgregori* fruit flies are known to be monophagic and only attack melinjo plants. Studies on this species are still very small (Saragih, 2017). Research about fruit flies on melinjo plants in Tanah Laut Regency which is the largest producer of melinjo in South Kalimantan has not been conducted and it is not known what kind of fruit flies attacking the melinjo. The purposes of this present study are to identify the species of fruit flies attacking melinjo plants in Damit Village, Batu Ampar Sub-district, Tanah Laut Regency, and to find out the abundance and sex ratio of the species population.

Materials and method

The study was conducted at the Entomology Laboratory of the Faculty of Agriculture, Lambung Mangkurat University, Banjarbaru, in March-April

2018. The sampling of melinjo fruit was carried out in 5 different points (Indriyanti *et al.*, 2014) with the distances adjusted to those in the location of study (Damit Village). The sampling of 250 pieces of melinjo fruit was conducted in each observation with three replications at the same location, with a 1-week lag from the first sampling.

At each point, the attacked fruit showed the symptom in the form of black spots on the melinjo fruit either falling from the tree or not. The symptomatic melinjo fruit was put into 5 plastics, each containing 10 pieces of fruit, labeled and then taken to the Entomology Laboratory and then moved into plastic jars with the bottom filled with wood sawdust and covered with gauze, and tied using raffia or rubber bands. Rearing was carried out for 22 days until the fruit fly imagoes were obtained (Larasati, 2012).






The fruit fly imagoes were preserved by putting them into a glass bottle containing alcohol. The imagoes were then identified for the type, abundance and sex ratio. The identification was carried out by observing the morphology of the wings, thorax and abdomen of fruit flies using a microscope and books (Siwi *et al.*, 2006).

Results and discussion

The fruit flies attacking the melinjo fruit were only one species. The results of the identification of fruit flies attacking melinjo fruits along with their morphology can be seen in Table 1.

The fruit flies found in melinjo fruit in Damit Village were only one species, namely *B. mcgregori*. According to Larasati *et al.* (2016), *B. mcgregori* is found only in melinjo fruit and is monofag. *B. mcgregori* is found in the melinjo varieties with large fruit (Yong *et al.*, 2014). The fruit flies are found in the Philippines, Singapore and Malaysia, and Nicobar Island (Hardy, 1973; Yong, 1994; Ranganath and Veenakumari, 1999; Yong *et al.*, 2014). This current report is the first report of the presence of *B. mcgregori* fruit flies attacking melinjo fruit in South Kalimantan.

Table 1. Morphological characteristics of *B. mcgregori* fruit fly.

Morphology	Note
<p>a</p> 	<p>On the wing there is a brown band on the costa line from the base of the upper wing to the tip of the wing passing through R₄₊₅ and the brown band on the anal cell.</p>
<p>b</p> 	<p>On the thorax there are yellow lines on the right and left sides of the back, and the scalp is also yellow.</p>
<p>c</p> 	<p>The abdomen has a yellowish brown color. There is a black line extending sideways on the second segment, and there is no black line extending to the next segment, but in the fourth segment there is a black line extending downward and there is a bump on the third segment.</p>
<p>d</p> 	<p>There is no facial spot on the face.</p>
<p>e</p> 	<p>The body size of this fruit fly is smaller compared to other <i>Bactrocera</i> species.</p>

Morphologically, *B. mcgregori* has distinctive characteristics that differentiate it from other types of *Bactrocera*. Its body size is smaller compared to *B. albistrigata*, *B. carambolae*, and *B. papayae* (Isnaini, 2013).

The size of its body, if measured from the head part to abdomen, is 5 mm. (Table 1e). In addition to the body

size, the color and the pattern on the abdomen and thorax of *B. mcgregori* does not have a special pattern, only in the form of a black line which is broken on the third segment and there are 7 pectan in the third section. Thorax is dominated with yellow, while in the abdomen it is pale reddish (Siwi *et al.*, 2006).




Fig. 1. Percentage of abundance of *B. mcgregori* fruit flies in Damit Villange, Tanah Laut Regency.

Abundance

The abundance obtained from each observation was very different. The highest abundance was seen in the second observation (P2M) which was 75.6% and the

lowest abundance was found in the first observation (P1M), 8.4% (Figure 1). The average abundance of all observations was 37.06%.




Fig. 2. Number of males and females of *B. mcgregori* in melinjo plants with the attacked fruit sampled three times in Damit Village

¹⁴ The number of fruit flies in each observation was strongly influenced by natural enemy factors namely parasitoid and not hatched pupae.

The high abundance in P2M observation was because the number of parasitoids found in these observations was very low with the ratio of fruit fly to parasitoid, 1: 3, while the low abundance in P1M observation was due to the large number of parasitoids with a ratio of

1: 7. Parasitoid larvae and pupae develop inside fruit fly larvae so that larvae or pupae of fruit flies die before becoming imagoes. Sota and Mogi (1996) state that the population of fruit flies is influenced by biotic factors especially feed and its natural enemies, and physical factors such as climate, bulk rain and temperature. According to Hasyim *et al.* (2008) the population of fruit flies on the land will increase due to an increase in rainfall and temperature.




Fig. 3. Comparison of sex ratio of *B. mcgregori* fruit flies on each attacked fruit sampling and its average in Damit Village.

Based on the observations, an increase in the number of the fruit fly population will be followed by a decrease in the number of parasitoids. This is contrary to Laba and Kartohardjono's (1998) statement that if the population of fruit flies increases, the parasitoid population will increase as well as, and vice versa.

Melinjo fruit that is ripe and deciduous has a thin peel texture and is easily attacked by fruit flies. The abundance of *B. mcgregori* in melinjo is categorized into the high category with an average abundance of 37.06%. This is stated by Krebs (1989) in Ismawan *et al.* (2015) that this abundance is included in the high range which is greater than 20%. However, this abundance does not affect the economic value of melinjo fruit because the utilization of melinjo peel is still less and only the seeds have the high economic value (raw materials for making salted or sweet

chips/emping). Melinjo trees on the research land were intentionally planted in a row with an area of \pm 0.5 ha. The number of trees, either large or small, did not affect the ability of *B. mcgregori* to attack these plants (direct observation in the field).

Sex Ratio

In the study, there were more female fruit flies compared to male fruit flies.

The male and female fruit flies emerging from the attacked melinjo fruit showed the highest number in the second observation (P2M) with the number of males and females of 87 and 102, respectively. Meanwhile the lowest number of fruit flies was in the first observation (P1M) with the number of males and females of 8 and 13, respectively. The average number of males and females found in the observations 1, 2 and 3 were 42 and 50, respectively (Figure 2).

The comparison between the sex ratio of ⁴ male and female fruit flies from the first observation to the third observation was very different (Figure 3).

The lowest sex ratio was 0.61 at the first observation, which increased to 0.88 at the third observation. If all is averaged, the sex ratio of male to female fruit flies is 0.78.

In this case the number of female flies is more than male flies because *B. mcgregori* can experience parthenogenesis (females can produce eggs without being fertilized by males and can hatch them into new individuals), so the population will not be affected if there is a lack of males.

The more the number of females is, the more likely the population of *B. mcgregori* will increase. This result is consistent with Isnaini's (2013) study in Demak Regency revealing that the most found fruit flies were females, so the more females, the more the population would increase.

It is suspected that the more eggs were placed on the attacked fruit. According to Indriyanti *et al.* (2014) the comparison between the sex ratio of *Bactrocera* spp. showing more females than males indicates that the population will increase. Meanwhile, the relationship of mating in *Bactrocera* spp. is polygamy, so the males can fertilize more than one female.

If it is seen from the contents of melinjo fruit such as high carbohydrate and protein, the two contents, especially carbohydrate which is a source of energy for fruit flies and sugar such as sucrose can stimulate the eating desire of the fruit flies (Susrama, 2017).

Due to its small body size compared to phytophagic insects (orthoptera, lepidoptera and coleoptera), the fruit fly requires higher carbohydrates compared to proteins and amino acids; therefore, melinjo fruit strongly supports the proliferation of *B. mcgregori* fruit flies. Generally, the fruit flies are found to attack when the fruit is nearly ripe or matured.

⁷ Conclusion

The species of fruit flies found in melinjo fruit is only one species, namely *Bactrocera mcgregori*, while the abundance category of fruit flies is categorized into high categories with an average abundance of 37.06% and an average sex ratio of males to females 0.78 (More female than male fruit flies).

References

- Badan Pusat Statistik.** 2016. Kabupaten Tanah Laut dalam Angka. Pelaihari.
- Hardy DE.** 1973. The Fruit Flies (Tephritidae: Diptera) of Thailand and Bordering Countries. Honolulu : Entomology Dept., Bernice P. Bishop Museum, 1973. Pacific insects monograph **31**, 353.
- Hasyim A, Muryati, Kogel WJ.** 2008. Population Fluctuation of Adults Males of the Fruit Fly, *Bactrocera tau* Walker (Diptera:Tephritidae) in Passion Fruit Orchards in Relation to Abiotic Factors and Sanitation. Indonesian Journal of Agricultural Science **9(1)**, 28-33.
<http://dx.doi.org/10.21082/ijas.v9n12008.p29-33>
- Indriyani DR, Isnaini YN, Priyono B.** 2014. Identifikasi dan Kelimpahan Lalat Buah *Bactrocera* pada Berbagai buah Terserang. Biosaintifika: Journal of Biology & Biology Education **6(1)**, 38-44.
<https://doi.org/10.15294/biosaintifika.v6i1.2933>
- Ismawan A, Rahayu SE, Dharmawan A.** 2015. Kelimpahan dan Keanekaragaman Burung di Preval Taman Nasional Kutai Kalimantan Timur. Jurusan Biologi FMIPA, Universitas Negeri Malang: Malang.
- Isnaini YN.** 2013. Identifikasi Spesies dan Kelimpahan Lalat Buah *Bactrocera* spp di Kabupaten Demak. Skripsi, Jurusan Biologi FMIPA, Universitas Negeri Semarang: Semarang.
- Kementerian Pertanian.** 2015. Statistik Produksi Hortikultura. Direktorat Jendral Hortikultura. Jakarta.

- Laba IW, Tartohardjono A.** 1998. Pelestarian Parasitoid dan Predator dalam Pengendalian Hama Tanaman. *Jurnal Penelitian dan Pengembangan Pertanian* **17(4)**, 122-129.
- Larasati A.** 2012. Keanekaragaman, Persebaran dan Kunci Identifikasi Lalat Buah (Diptera: Tephritidae) di Kabupaten Bogor dan Sekitarnya. Tesis. Pascasarjana Institut Pertanian Bogor: Bogor.
- Larasati A, Hidayat P, Buchori D.** 2016. Kunci Identifikasi Lalat Buah (Diptera: Tephritidae) di Kabupaten Bogor dan sekitarnya. *Jurnal Entomologi Indonesia* **13(1)**, 49-61.
<http://dx.doi.org/10.5994/jei.13.1.49>
- Purnomosidhi P, Suparman, Roshetko JM, Mulawarman.** 2002. Perbanyak dan Budidaya Tanaman Buah-buahan. International Centre for Research in Agroforestry (ICRAF) dan Winrock International. Bogor, Indonesia.
- Ranganath HR, Veenakumari K.** 1999. Notes on the Dacine Fruits Flies (Diptera: Tephritidae) of Andaman and Nicobar Island. *Journal Raffles Bulletin of Zoologi* **(1)**, 221-224.
- Santoso M, Naka Y, Angkawidjaja C, Yamaguchi T, Matoba T, Takamura T.** 2010. Antioxidant and Damage Prevention Activities of the Edible Parts of Gnetum gnemon and Their Change upon Heat Treatment. *Journal Food Science and Technology Research* **16(6)**, 549-556.
<http://dx.doi.org/10.3136/fstr.16.549>
- Saragih RB.** 2017. Komposisi Parasitoid Lalat Buah Melinjo (*Bactrocera Mcgregori*) di Kabupaten Bantul. Universitas Gadjah Mada: Yogyakarta.
- Siwi SS, Hidayat P, Suputa.** 2006. Taksonomi dan Bioekologi Lalat Buah Penting di Indonesia (Diptera: Tephritidae). Balai Besar Penelitian dan Pengembangan Bioteknologi dan Sumberdaya Genetik Pertanian: Bogor. 65 p.
- Sota T, Mogi M.** 1996. Species Richness and Altitudinal Variation in the Aquatic Metazoan Community in Bamboo Phytotemata from North Sulawesi. *Researches on Population Ecology* **38(2)**, 275-281.
<http://dx.doi.org/10.1007/BF02515737>
- Susrama I GK.** 2017. Kebutuhan Nutrisi dan Substansi dalam Pakan Buatan Serangga. Artikel Ulasan, Program Studi Agroekoteknologi, Universitas Udayana: Bali.
- Yong HS.** 1994. The Gnemon Fruit Fly. *Nature Malaysia* **19**, 37-40.
- Yong HS, Lim PE, Tan J, Sauna IW.** 2014. Gentum Gnemon (Gnetaceae): A New Host Plant of Carambola Fruit Fly *Bactrocera Carambolae* (Insecta: Tephritidae). *Journal of science and technology in the tropics* **10**, 39-44.

Identification and abundance of fruit fly species (*Bactrocera mcgregori*) in melinjo plant (*Gnetum gnemon*) in Damit Village, Batu Ampar Sub-district, Tanah Laut Regency

ORIGINALITY REPORT

15%

SIMILARITY INDEX

5%

INTERNET SOURCES

13%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to Higher Education Commission Pakistan **2%**
Student Paper

- 2** Herry Marta Saputra, Tasya Dwi Nanda, Rion Apriyadi, Henri Henri, Fahri Setiawan. "KEANEKARAGAMAN HAMA LALAT BUAH PADA TANAMAN SAYURAN BUAH DI KABUPATEN BANGKA DAN KUNCI IDENTIFIKASINYA", Jurnal Agrotek Tropika, 2023 **2%**
Publication

- 3** Jeffij V. Hasinu, John A. Patty, Gratiana N. C. Tuhumury. "MORPHOLOGICAL IDENTIFICATION AND POPULATION OF FRUIT FLY (*Bactrocera* sp.) (Diptera: Tephritidae) IN CHILI FIELDS, SAVANAJAYA VILLAGE BURU DISTRICT", Jurnal Hama dan Penyakit Tumbuhan Tropika, 2020 **2%**
Publication

4

Agus Susanto, Wahyu Daradjat Natawigena, Luciana Djaya, Tohidin Tohidin, Fauza Saputra. "The Effect of Methyl Eugenol Block Plus on Bactrocera dorsalis Complex Total Captured in Chili Plantation", Jurnal Biodjati, 2021

Publication

1 %

5

S Sulaeha, A H Bahtiar, M Melina. " Identification fruit fly species associated with watermelon plants ((Thunb.) Matsum. & Nakai) in South of Sulawesi, Indonesia ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

1 %

6

businessdocbox.com

Internet Source

1 %

7

Herry Marta Saputra, Budi Afriyansyah. "DISTRIBUTION AND IDENTIFICATION OF FRUIT FLIES (DIPTERA: TEPHRITIDAE) ATTRACTED ON METHYL EUGANOL AND CUE LURE IN CENTRAL BANGKA REGENCY, BANGKA BELITUNG", Jurnal Hama dan Penyakit Tumbuhan Tropika, 2021

Publication

1 %

8

Rahayu Supratiwi, Rion Apriyadi, Euis Asriani. "FRUIT FLIES (DIPTERA: TEPHRITIDAE) DIVERSITY IN HORTICULTURAL FARM OF MERAWANG SUB-DISTRICT, BANGKA DISTRICT, BANGKA BELITUNG ISLANDS",

1 %

Jurnal Hama dan Penyakit Tumbuhan Tropika, 2020

Publication

9

Agus Susanto, Tohidin, Toto Sunarto, Luciana Veronica Sinaga, Ariyo Nugroho, Muhamad Basuki, Luciana Djaya, Andi Fadillah. "Effect of Trap Height Level on The Capture of Fruit Fly (*Bactrocera* spp.) on Crystal Guava Field", IOP Conference Series: Earth and Environmental Science, 2023

Publication

1 %

10

Submitted to Lambung Mangkurat University

Student Paper

1 %

11

Nan Zarchi Win, Khin Mi Mi, Thi Tar Oo, Kyaw Kyaw Win, Jinyoung PARK, Jong Kyun PARK. "Occurrence of Fruit Flies (Diptera: Tephritidae) in Fruit Orchards from Myanmar", Korean Journal of Applied Entomology, 2014

Publication

<1 %

12

etd.repository.ugm.ac.id

Internet Source

<1 %

13

idoc.pub

Internet Source

<1 %

14

jhpttropika.fp.unila.ac.id

Internet Source

<1 %

twj.ulm.ac.id

15

Internet Source

<1 %

16

, Jean Bedel Fagbohoun. "Thermal stability of two xylanases from Macrotermes subhyalinus little soldier: kinetic and thermodynamic analysis", AfricArXiv, 2022

<1 %

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Identification and abundance of fruit fly species (*Bactrocera mcgregori*) in melinjo plant (*Gnetum gnemon*) in Damit Village, Batu Ampar Sub-district, Tanah Laut Regency

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/0

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
